

- Tanaka, Y., 1958: Seasonal changes occurring in the gonad of *Stichopus japonicus*. — Bull. Fac. Fish. Hokkaido Univ. 9: 29-36.
- Théel, H.J., 1876: Note sur l'*Elpidia*, genre nouveau du groupe des Holothurides. — Bih. K. svenska Vetensk. Akad. Handl. 4, 4: 1-7.
- 1877: Mémoire sur l'*Elpidia*, nouveau genre d'Holothurides. — K. svenska Vetensk. Akad. Handl. 14, 8: 1-30.
- 1879: Preliminary report on the Holothuridae of the Exploring voyage of H. M. S. "Challenger", I. — Bih. K. svenska Vetensk. Akad. Handl. 5, 19: 1-20.
- 1882: Report on the Holothurioidea, I. — Rep. scient. Results Voyage Challenger, Zool. 4, 13: 1-176.
- 1886a: Report on the Holothurioidea, II. — Ibid. 14, 39: 1-290.
- 1886b: Report on the Holothurioidea. Reports on the results of dredging... "Blake", 30. — Bull. Mus. comp. Zool. Harv. 13: 1-21.
- Thorson, G., 1946: Reproduction and larval development of Danish marine bottom invertebrates. — Meddr Kommn Danm. Fisk.- og Havunders., Ser. Plankton 4: 1-523.
- 1952: Zur jetzigen Lage der marinen Bodentier-Ökologie. — Zool. Anz., Suppl. 16: 276-327.
- Ursin, E., 1960: A quantitative investigation of the echinoderm fauna of the central North Sea. — Meddr Danm. Fisk.- og Havunders., N. S. 2, 24: 1-204.
- Ushakov, P. V., 1952: Investigations on the deep-sea fauna. — Priroda, Mosk. 1952, No. 6: 100-102. (In Russian).
- Vaney, C., 1906: Holothurides. — Expédition Antarctique Française (1903-1905). 30 pp.
- 1908: Les Holothurides de l'Expédition Antarctique Nationale Écossaise. — Trans. R. Soc. Edinb. 46: 405-441.
- 1914: Holothurides. — Deuxième Expédition Antarctique Française (1908-1910). 54 pp.
- Vasseur, E., 1952: Geographic variation in the Norwegian sea-urchins, *Strongylocentrotus droebachiensis* and *S. pallidus*. — Evolution, Lancaster, Pa. 6: 87-100.
- Verrill, A. E., 1884: Notice of a remarkable marine fauna occupying the outer banks off the southern coast of New England, No. 9. — Am. J. Sci. 28: 213-220.
- Verrill, A. E., 1885: Results of the explorations made by the steamer "Albatross" off the northern coast of the United States in 1883. — United States Commission of Fish and Fisheries, Part XI. Report of the Commissioner for 1883: 503-699.
- Vinogradova, N. G., 1959: The zoogeographical distribution of the deep-water bottom fauna in the abyssal zone of the ocean. — Deep Sea Res. 5: 205-208.
- 1962a: Vertical zonation in the distribution of deep-sea benthic fauna in the ocean. — Ibid. 8: 245-250.
- 1962b: Some problems of the study of deep-sea bottom fauna. — J. oceanogr. Soc. Japan, 20th Anniversary Volume: 724-741.
- Walsh, J. H. T., 1891: Natural history notes from H. M. Indian Marine Survey steamer "Investigator". No. 24. List of deep-sea holothurians, collected during seasons 1887-91, with descriptions of new species. — J. Asiatic Soc. Beng. 60: 197-204.
- Wolff, T., 1956a: Isopoda from depths exceeding 6000 meters. — Galathea Rep. 2: 85-157.
- 1956b: Crustacea Tanaidacea from depths exceeding 6000 meters. — Ibid. 2: 187-241.
- 1960: The hadal community, an introduction. — Deep Sea Res. 6: 95-124.
- 1961: Animal life from a single abyssal trawling. — Galathea Rep. 5: 129-162.
- 1962: The systematics and biology of bathyal and abyssal Isopoda Asellota. — Ibid. 6: 1-320.
- 1970: The concept of the hadal or ultra-abyssal fauna. — Deep Sea Res. 17: 983-1003.
- 1971: "Archimède" dive 7 to 4160 metres at Madeira: Observations and collecting results. — Vidensk. Meddr dansk naturh. Foren. 134: 127-147.
- Zenkevich, L. A., 1963: Biology of the seas of the USSR. — London. 955 pp.
- & Ya. Birshtein, 1956: Studies of the deep water fauna and related problems. — Deep Sea Res. 4: 54-64.
- — 1960: On the problem of the antiquity of the deep-sea fauna. — Ibid. 7: 10-23.
- Z. A. Filatova, G. M. Belyaev, T. S. Luk'yanova & I. A. Suetova, 1971: Quantitative distribution of zoobenthos in the world ocean. — Byull. mosk. Obshch. Ispyt. Prir. (Biol.) 76: 27-33. (In Russian, English summary).

## VI. INDEX TO NAMES OF GENERA AND SPECIES


Main entries in *italics*. Generic names preceded by \* refer to non-Elasipoda. Specific names in keys, diagrams of bathymetric distribution, and maps of geographic distribution are not included (see main entry to generic name).

* <i>Abyssocucumis</i> , 216	130, 150, 155-156, 159,	<i>Amperima</i> , 128, 129, 130,	186, 226, 235
<i>abyssorum</i> , 216, 234,	184, 187, 188, 190, 191,	158, 184, 186, 189, 192,	<i>tui</i> , 158
241	192, 213, 229	213, 229	<i>velacula</i> , 161, 185, 186,
* <i>Acanthotrochus</i> , 216	<i>ecalcareia</i> , 155, 156-157,	<i>furcata</i> , 159, 186, 226	215, 234
<i>mirabilis</i> , 216	185, 186, 187	<i>insignis</i> , 162	<i>Amphideima</i> , 15, 38
* <i>Acaudina</i> , 216	<i>lactea</i> , 156, 157	<i>naresi</i> , 159-161, 184,	<i>investigatoris</i> , 39, 43,
* <i>Achiridota</i> , 217	<i>monactinica</i> , 155,	185, 186, 192, 196, 204,	44, 45
<i>inermis</i> , 216	157-158, 187	242	* <i>Amphigymnas</i> , 215
<i>profunda</i> , 216	<i>paradoxa</i> , 156, 157	<i>robusta</i> , 161-162, 189	<i>Apodogaster</i> , 47, 75, 190,
<i>Achlyonice</i> , 127, 128, 129,	<i>tui</i> , 158, 187, 215	<i>rosea</i> , 158-159, 185,	192, 210, 215

- alcocki*, 75, 234  
 \**Astichopus*, 204  
   *multifidus*, 204  
*Bathygone*, 52  
   *papillatum*, 67, 69, 70  
 \**Bathyplotes*, 186, 215, 216  
   *moseleyi*, 234  
   *natans*, 204, 234  
   *reptans*, 234  
*Benthodytes*, 10, 76-77,  
 190, 191, 192, 194, 206,  
 211, 225, 226  
   *abyssicola*, 76  
   *assimilis*, 106, 108  
   *browni*, 76  
   *gigantea*, 76  
   *glutinosa*, 89  
   *gotoi*, 77, 79, 80  
   *hystrix*, 84, 85, 86, 87  
   *incerta*, 76, 77-80, 87,  
   191, 203  
   *janthina*, 76, 80, 82, 89,  
   93  
   *kerhervei*, 102, 105  
   *lingua*, 76, 80-82, 87,  
   190, 191, 203, 214, 215  
   *mamillifera*, 76  
   *papillifera*, 89, 92, 93  
   *plana*, 76, 84, 87-89,  
   235  
   *recta*, 99  
   *regularis*, 77, 80  
   *salivosus*, 75  
   *sanguinolenta*, 76, 93,  
   94-96, 190, 191, 194,  
   203, 226  
   *selenkiana*, 76  
   *sibogae*, 76, 84-87, 203,  
   220  
   *sordida*, 76  
   *spuma*, 76  
   *superba*, 76, 89, 190  
   *typica*, 76, 89-93, 187,  
   191, 194, 202, 203, 204,  
   205, 223, 227, 237, 238  
   *valdiviae*, 76, 82-84  
*Benthogone*, 47, 48, 52,  
 190, 206, 210, 224  
   *abstrusa*, 51-52, 220  
   *fragilis*, 49-50, 191  
   *quadrilineata*, 48, 49  
   *quatrolineata*, 67  
   *rosea*, 48-49, 50, 187,  
   191, 202, 221  
*Benthophyces*, 48, 50  
   *fragilis*, 49, 50  
 \**Benthothuria*, 215  
 \**Capheira*, 215  
 \**Caudina*, 216  
   *chilensis*, 189  
 \**Ceraplectana*, 216  
   *trachyderma*, 216  
 \**Chiridota*, 216  
   *abyssicola*, 216  
*Cryodora*, 52  
   *spongiosa*, 58  
 \**Cucumaria*, 189  
   *echinata*, 189  
*Deima*, 15, 16, 183, 190,  
 191, 193, 205, 206, 210,  
 222  
   *atlanticum*, 18, 20, 22  
   *blakei*, 17, 20, 21  
   *fastosum*, 17, 20, 21,  
   22, 23, 194  
   *mosaicum*, 18, 20, 22, 23  
   *pacificum*, 20, 24  
   *validum*, 16-24, 190,  
   191, 194, 198, 199, 215,  
   223, 227, 228, 237, 241,  
   245  
   *v. pacificum*, 17, 22, 23,  
   24  
   *v. validum*, 17-23, 204  
 \**Dendrothuria*, 215  
 \**Ekmocucumis*, 216  
   *turqueti*, 234  
*Ellipinion*, 128, 129, 130,  
 131, 162-163, 184, 186,  
 189, 192, 213, 230  
   *bucephalum*, 163-164  
   *delagei*, 163, 215  
   *facetum*, 166, 189  
   *galathea*, 165, 192, 241  
   *kumai*, 162, 166, 189,  
   215  
   *molle*, 165, 166, 169  
   *papillosum*, 165-166,  
   189  
   *solidum*, 163, 164-165  
*Elpidia*, 9, 10, 127, 128,  
 130, 172-174, 184, 189,  
 190, 192, 196, 205, 207,  
 213, 227, 231, 232, 241,  
 243  
   *adenensis*, 172, 175, 232  
   *ambigua*, 151  
   *antarctica*, 174  
   *atakama*, 172, 181, 182,  
   232, 241, 242  
   *birsteini*, 172, 181, 232,  
   242, 243  
   *chilensis*, 172, 175, 232  
   *glacialis*, 8, 172, 173,  
   175-181, 185, 186, 190,  
   196, 204, 205, 215, 223,  
   232, 234, 237, 242, 245  
   *g. glacialis*, 175,  
   176-178  
   *g. kermadecensis*, 180  
   *g. kurilensis*, 180-181,  
   243  
   *g. solomonensis*, 180,  
   181  
   *g. sundensis*, 178-180  
   *g. theeli*, 174  
   *g. uschakovi*, 180  
   *globosa*, 167  
   *hanseni*, 172, 181, 242,  
   243  
   *h. hanseni*, 181, 232  
   *h. idzubonensis*, 181,  
   232  
   *incerta*, 143  
   *javanica*, 178, 180  
   *kermadecensis*, 180  
   *kurilensis*, 180, 243  
   *longicirrata*, 172, 181,  
   232, 241, 243  
   *minutissima*, 172, 175,  
   232  
   *mollis*, 165  
   *murrayi*, 167  
   *nana*, 170  
   *papillosa*, 165  
   *purpurea*, 151  
   *rigida*, 136  
   *solomonensis*, 180  
   *sundensis*, 178  
   *theeli*, 172, 174-175,  
   185, 186, 232  
   *uschakovi*, 180  
   *verrucosa*, 147  
   *willemoesi*, 144  
*Elpidiogone*, 128, 129, 131  
   *dubia*, 144  
*Eynpniastes*, 127, 128  
*Euphronides*, 99  
   *anchora*, 102, 103, 104,  
   105  
   *auriculata*, 106, 108  
   *bifurcata*, 102, 112  
   *cornuta*, 106, 109, 110  
   *depressa*, 106, 108, 109,  
   110  
   *dyscrita*, 116, 124  
   *kerhervei*, 102, 105  
   *scotiae*, 111  
   *talismani*, 106, 108, 109,  
   110  
   *tanneri*, 106, 108, 109,  
   110  
   *verrucosa*, 112  
   *violacea*, 106, 108, 109,  
   110  
 \**Eupyrgus*, 216  
   *pacificus*, 216  
   *scaber*, 216  
*Euriplastes*, 128, 132  
 \**Filithuria*, 215  
 \**Galatheathuria*, 185, 204  
   *aspera*, 185, 204, 205  
 \**Gastrothuria*, 215  
 \**Gephyrothuria*, 216  
 \**Hadalothuria*, 195, 216  
   *wolffi*, 242  
 \**Hedingia*, 216  
   *albicans*, 216  
   *californica*, 216  
   *fusiforme*, 216  
   *ludwigi*, 216  
   *planapertura*, 216  
 \**Heteromolpadia*, 216  
   *marenzelleri*, 216  
   *pikei*, 216  
   *tridens*, 216  
 \**Holothuria*, 184  
*Ilyodaemon*, 48, 52  
   *abstrusus*, 48, 51, 190  
   *fimbriatus*, 64, 67, 69  
   *ijimai*, 67  
   *maculatus*, 63  
   *miurense*, 63, 65, 66  
*Irpa*, 8, 127, 128, 130, 131,  
 171, 186, 189, 192, 194,  
 213, 230  
   *abyssicola*, 171, 172,  
   232  
   *ludwigi*, 171, 172, 185,  
   186, 189, 215, 235  
 \**Karenilla*, 215  
*Kolga*, 8, 127, 128, 130,  
 131, 170, 186, 187, 188,  
 192, 194, 213, 230  
   *foliacea*, 152  
   *furcata*, 159  
   *hyalina*, 170-171, 187,  
   188, 205, 225, 232  
   *ludwigi*, 172  
   *nana*, 170, 171, 188  
   *obsoleta*, 134  
*Laetmenoeus*, 52  
   *scotoeides*, 52, 61  
*Laetmogone*, 47, 48, 52-  
 53, 191, 210, 224, 225  
   *biserialis*, 52, 53, 69,  
   70-71, 191, 220, 238  
   *brongniarti*, 58  
   *enisus*, 64, 65  
   *fimbriata*, 52, 53, 67-  
   70, 71, 191, 202, 215,  
   221, 228, 238  
   *ijimai*, 52, 66, 67, 192,  
   215  
   *interjacens*, 47, 52, 53-  
   54, 191, 220  
   *jourdaini*, 58  
   *maculata*, 52, 63-67,  
   191, 202, 215, 228, 238  
   *neglecta*, 70, 71  
   *parva*, 67, 69, 70  
   *scotoeides*, 52, 61-63,  
   220  
   *selenkai*, 67, 69, 70, 71  
   *spongiosa*, 58  
   *theeli*, 52, 57-58, 61,  
   191, 215, 246  
   *violacea*, 52, 57, 58-61,

- 185, 191, 201, 202, 215, 221, 234, 238  
*wyvillethomsoni*, 52, 53, 54-57, 58, 61, 70, 71, 191, 215, 246  
*Laetmophasma*, 72  
*fecundum*, 72, 74  
\**Leptosynapta*, 189  
*inhaerens*, 189  
\**Meseres*, 215  
\**Mesothuria*, 215  
*bifurcata*, 234  
*candelabri*, 241  
*intestinalis*, 235  
*verrilli*, 235  
\**Molpadia*, 216, 246  
*antarctica*, 234  
*bathybia*, 216  
*blakei*, 216, 241, 246  
*granulata*, 216, 246  
*musculus*, 216, 235  
*parva*, 216  
\**Myriotrochus*, 10, 210, 216, 217  
*bathybius*, 216  
*bruuni*, 242  
*giganteus*, 216  
*kermadecensis*, 242  
*kurilensis*, 241  
*longissimus*, 241  
*mitis*, 242  
*theeli*, 216  
*zenkevitchi*, 242  
*Nectothuria*, 99  
*translucida*, 116, 124, 125  
*Oneirophanta*, 15, 16, 24, 183, 191, 205, 206, 210, 215, 222  
*affinis*, 32, 36  
*alternata*, 25, 31  
*conservata*, 36, 222  
*mutabilis*, 24-36, 187, 190, 193, 194, 198, 214, 223, 226, 231, 232, 238, 241, 245  
*m. affinis*, 32-36, 191, 199, 200, 201, 204, 222  
*m. mutabilis*, 24-32, 199, 201, 222  
*setigera*, 36-38, 47, 186, 194, 198, 222  
*Orphmurgus*, 15, 16, 38, 191, 206, 210, 215, 220, 223  
*asper*, 16, 38-39, 221  
*glaber*, 16, 39-46, 183, 190, 194, 198, 220, 228, 238, 245  
*insignis*, 16, 39, 43, 44, 45, 46, 47  
*invalidus*, 39, 43, 45  
*protectus*, 16, 47  
*rigidus*, 39, 43, 45, 46  
*vitreus*, 16, 46-47, 220  
\**Paelopatides*, 76, 204, 215  
*aspera*, 185  
\**Paelpidia*, 208  
*norica*, 208  
*Pannychia*, 47, 72, 192, 210, 224  
*glutinosa*, 80, 82  
*moseleyi*, 72-75, 202, 220, 238  
*m. mollis*, 72, 73  
*m. virgulifera*, 72, 73, 74  
*multiradiata*, 72, 74  
*pallida*, 72, 74  
\**Paracaudina*, 216  
*Paelpidia*, 127, 128, 129, 131, 190  
*anamesa*, 132  
*cylindrica*, 147  
*elongata*, 147  
\**Paroriza*, 241  
*grevei*, 241  
*Pelagothuria*, 128  
*Peniagone*, 127, 128, 129, 130, 131-134, 187, 188, 192, 194, 195, 205, 207, 212, 215, 228, 229  
*affinis*, 139, 142-143, 187  
*atrox*, 132  
*azorica*, 138-142, 146, 187, 204, 206, 215, 242  
*bispiculata*, 132  
*challengeri*, 146  
*convexa*, 135-136, 152, 187  
*diaphana*, 152, 153-155, 187, 192, 205, 238  
*discrepans*, 131  
*dubia*, 144-145  
*ecalcareia*, 131  
*elongata*, 147, 148, 187  
*expansa*, 152, 153  
*ferruginea*, 151  
*foliacea*, 152, 153  
*gracilis*, 152, 155  
*horrifer*, 132  
*humilis*, 138  
*incerta*, 139, 142, 143-144  
*incondita*, 145  
*intermedia*, 152-153  
*islandica*, 150, 191  
*japonica*, 147, 187, 215, 233  
*lacinora*, 151  
*lugubris*, 148, 151  
*mossmani*, 134  
*mus*, 132  
*naresi*, 159  
*nybelini*, 134  
*obscura*, 132  
*obsoleta*, 134-135, 136, 152  
*papillata*, 145-146, 187  
*piriei*, 132  
*porcella*, 134  
*purpurea*, 151-152, 153, 155, 186, 187, 205  
*rigida*, 136-137, 138, 152  
*setosa*, 148, 149  
*stabilis*, 132  
*theeli*, 146, 147, 185  
*vedeli*, 138, 139  
*verrucosa*, 147-148  
*vexillum*, 132  
*vignoni*, 146-147, 215, 233  
*vitrea*, 143, 148-150, 187  
*willemoesi*, 138, 139, 142, 144, 187  
*willtoni*, 134  
*wyvillii*, 131, 150-151  
*Periamma*, 128, 158  
*furcata*, 159  
*kumai*, 166  
*ludwigi*, 172  
*naresi*, 159  
*roseum*, 158  
*tetramerum*, 159, 160  
\**Praeuephronides*, 209  
*multiperforata*, 209  
\**Protankyra*, 217  
*abyssicola*, 217  
*brychia*, 217  
\**Pseudostichopus*, 215  
*mollis*, 234  
*occultatus*, 235  
*villosus*, 242  
\**Pseudothuria*, 215  
\**Psolidium*, 216  
*disjunctum*, 216  
*panamense*, 216  
\**Psolus*, 216  
*phantapus*, 189, 196  
*pourtalesii*, 216  
*Psycheotrepes*, 76, 96, 191, 192, 211, 227  
*exigua*, 96, 98, 99, 191, 203  
*magna*, 96, 97-99, 194, 241  
*recta*, 96, 99  
*Psychrelpidia*, 128, 129, 130, 131, 187, 190, 192, 194, 195, 206, 207, 212, 227  
*discrepans*, 131  
*Psychropotes*, 76, 99, 191, 192, 196, 204, 211, 226, 227  
*belyaevi*, 100-102, 194, 203, 204, 235, 244  
*brucei*, 116, 122, 123, 124, 125  
*buglossa*, 116, 122  
*depressa*, 106-111, 187, 191, 192, 194, 203, 204, 214, 215, 223, 234  
*dubiosa*, 116, 124  
*fucata*, 116, 122  
*grimaldii*, 116, 122  
*kerhervei*, 102, 105  
*laticauda*, 116, 122, 123, 124, 125  
*longicauda*, 99, 102, 115-126, 127, 182, 187, 190, 191, 194, 203, 205, 214, 223, 226, 232, 238, 241  
*loveni*, 126-127, 191  
*minuta*, 106  
*mirabilis*, 102, 111-112, 115  
*raripeis*, 116, 124  
*scotiae*, 111  
*semperiana*, 102-105, 106, 192, 194, 203  
*verrucosa*, 102, 112-115, 194, 203  
*Rhipidothuria*, 128, 129, 130, 131, 190, 192, 194, 212, 227  
*racowitzai*, 131  
*Scotoanassa*, 127, 128, 129, 131, 152  
*diaphana*, 153  
*gracilis*, 155  
*incerta*, 132, 228  
*translucida*, 153, 154  
*Scotodeima*, 15, 38  
*parvispiculatum*, 15  
*protectum*, 47  
*setigerum*, 36  
*vitreum*, 46  
*Scotoplanes*, 9, 128, 129, 130, 131, 166, 168, 184, 186, 189, 190, 192, 205, 213, 230, 234  
*albida*, 162  
*angelicus*, 165, 166  
*clarki*, 166, 169-170, 184, 189, 196, 204, 234, 246  
*delagei*, 162, 163  
*facetus*, 166  
*galathea*, 165  
*gilpinbrowni*, 155  
*globosa*, 166, 167-169, 170, 184, 189, 196, 204, 215, 223, 232, 234,

- |                                   |                                 |                               |                            |
|-----------------------------------|---------------------------------|-------------------------------|----------------------------|
| 242, 246                          | * <i>Stichopus</i> , 186        | <i>discoidalis</i> , 215      | <i>triplax</i> , 215       |
| <i>insignis</i> , 162             | <i>japonicus</i> , 197          | <i>gilberti</i> , 216         | <i>viridilimus</i> , 215   |
| <i>mollis</i> , 165               | <i>regalis</i> , 215, 235       | <i>heteroculus</i> , 215      | * <i>Synaptula</i> , 189   |
| <i>murrayi</i> , 167, 168, 169    | <i>tremulus</i> , 215           | <i>horridus</i> , 216         | <i>hydriformis</i> , 189   |
| <i>papillosa</i> , 165            | * <i>Synallactes</i> , 215, 216 | <i>ishikawai</i> , 215        | <i>Triconus</i> , 99       |
| <i>robusta</i> , 161              | <i>aenigma</i> , 15, 216        | <i>mollis</i> , 215           | * <i>Trochoderma</i> , 216 |
| <i>theeli</i> , 167, 168, 169     | <i>alexandri</i> , 215          | <i>monoculus</i> , 215        | <i>elegans</i> , 216       |
| * <i>Sphaerothuria</i> , 216      | <i>anceps</i> , 216             | <i>multivesiculatus</i> , 215 | * <i>Trochostoma</i> , 246 |
| <i>bitentaculata</i> , 216        | <i>carthagei</i> , 234          | <i>nozawai</i> , 215          | <i>granulatum</i> , 246    |
| * <i>Staurocucumis</i> , 216      | <i>challengeri</i> , 215        | <i>profundus</i> , 216        | <i>Tutela</i> , 172        |
| <i>liouvillei</i> , 182, 208, 234 | <i>chuni</i> , 215              | <i>rigidus</i> , 216          | <i>echinata</i> , 175      |
|                                   | <i>crucifera</i> , 216          | <i>robertsoni</i> , 216       |                            |



#### Additions and corrections

P. 76, right column, line 6 from below: Koehler & Vaney, 1905 ..... Read Vaney, 1908.  
P. 127, left column, line 3 from below: Théel, 1879 ..... Read Théel, 1882.  
P. 235, left column. With regard to the penetration of species into the Mediterranean deep sea the following information from E. Tortonese 1965 (Fauna d'Italia, 6) should be noted: (1) *Mesothuria verrilli* may not occur in the Mediterranean, the recorded specimens probably belonging to *M. intestinalis*. (2) *Molpadia musculus* penetrates to at least 820 m. (3) The molpadonian *Hedingia mediterranea* is known from the type specimen only (Tyrrhenian Sea: 800–1005 m). The species is likely to be identical with *H. albicans* (North Atlantic, Bay of Bengal, southwestern Pacific: 491–3200 m).