

Achras

obras

Sayer  
Lula  
essi

lo

vel

aii

Frage

macrophylla Achras foliis lanceolatis - ellipticis alternis, pedunculis axillaribus  
aggregatis

Habitat in Montserrat ad Ryan 5

Rami teretes, crassitie digiti minimi, cortice griseo albido, glabri.

Folia petiolata, alterna, semi-pedalia, pollices tres et ultra lata, integerrima, basi versus sensim parum angustiora, apice obtusiuscula glabra, venosa, plana, supra nitida, subtus pallidiora: costa et nervi purpurascens.

Petioles unguiculares, triquetri, supra plani, ~~sub~~ glauci.

Pedunculi axillares, inferne copiosiores, superiores pauciores, petiolo parum longioris, uniglori.

Calycis foliola quinque, <sup>regio</sup> duo extenora vellosiuscula rotundiora, tria interna oblonga, concava

Corolla monopetala, tubulosa, unguiculans, diametro penna cygneo, b. 2. vel 3.

Stigma: laciniis oblongis, rotundatis, erectis

Nectarium squama totius laciniis castro, ~~pubulata~~ inter singula lacinia, ~~castro~~ <sup>castro</sup> breviora laciniis dimidio breviora

Stamina filamenta 6. 7. 8 in medio singula laciniis ~~antibus~~ <sup>antibus</sup> oblonga acuta erecta, corolla breviores



~~microphyllum~~ ~~Adiantum~~

Pest. German superum villosum, hemisphaericum. Stylus subulatus; teres, longitudine  
corolla, purpurascens, nitidus. Stigma simplex

(Semina plura adesse, nisi si recte memini, retulit Dr. Ryan, hinc potius Aethras  
quam Bumelia, et eodem vno lat. illustr: addit <sup>quod</sup> non raro semina omnia  
propter unicum in Aethras abortientia abortiunt.


⊕ Semina 2-3. nucis juglande majores flavo basi nitidi longissimi hinc longi  
-tudinali



a



b



c




d



e

a flor magnitudinis naturali  
 b - corolla  
 c - eadem dispecte  
 d calyx cum pistillo  
 e Pistillum

de hinc macro  
 phylla



*Achras macrophylla*



*Desfeita Aikmar*

8088

Habitat herb. Mus.

Rami glabri, crassitie pennis anferis, ciliati. Folia in apice, petiolata, bipollucina, cuneata, emarginata, supra nitida, subtus tenuissime venosa, subconacca, utraque glabra. Petioli purpureascentes. Peduncululi ad apices ramorum, reflexi, purpurei, pollicares. Calycis foliolo, uti petala lanceolata, glabra, extra tomentosa.

*Volata Achnar*

*Habitat en Martinique la Manche*

*habes*



Achras

Habitat in <sup>China</sup> ~~Cena~~ Dombey herb. Guss.

Arbor 40 pedalis. Rami nutantes, apice ferrugineo tomentosi. Folia versus apicem  
4. 5. pollicaria, elliptica, subacuminata, glabra supra nitida, subtus pallidiora  
submembranacea, venosa, basi versus apicem parum angustata. Petioli tomentosi  
ferruginei, vix semipollicares. Pedunculi axillares, solitarii, uni flori, uti Calyx  
et ferrugineo tomentosi.

Calyx 5-phyllus: foliola oblonga, duo exteriora longiora. Corolla 5-fida: lacina  
obtusè lanceolata. Stamina 5. Corolla affixa. Squama 5: filamenta cu  
stata asperitantes. Stylus corollae longior. Stigma obtusum. Perium 1 obli  
care. Semina 1. 2. 3. 4. figura Castanea Dombey.



*Cecropia*

*Balata* Aubl.

Habitat in Guiana herb. Jusq. 5 Richard

Rami crassitie digiti intermedii, glabri. Folia versus apicem, petiolata, spithamea et ultra, elliptica, glaberrima, coriacea, avenia, enervia, acuminata brevi terminata, subtus pallida, costa subtus extans, striata. Petioli sesqui pollicares, veniosissime subtomentosi.

aliud specimen ab Aubletis ipse, acumina emarginata. Folia, 2-3 pollicaria, supra nitida. Peduncululi brevissimi, uniflori, tomentosi.

1 mammosa Aethras floribus solitariis, foliis cuneiformi - lanceolatis Lin. I. V. p. 347.  
Malus persica maxima, foliis magnis integris longis, fructu maxime oblongo  
scabro: ospeculo patina nigro patina glabo Sloan hist. 2. p. 124. t. 21 P.  
Arbor americana pomifera, frondosis ramulis, foliis amplis longioribus obtusis  
dunis et venosis margine aequali Pluk. alm. 39. t. 260. fig. 2.  
Fructus oblongus utrinque acuminatus s. conicus lavis splendens pediculus  
Raj. hist. 1000

~~fructibus paucis oblongis turgidis~~  
Aethras fructu maximo ovato (Brown jam. 201)

Aethras floribus pentadonis facque Amer. 56. t. 102. f. 14

Habitat in America meridionali

Ramuli ad folia tomentosa, tuberculati et crenatifati e capite foliorum florum et foliorum. Folia oblonga, basi attenuata, ac uti in stipula nitida, venosa, petiolata, spollie ana ad apices ramulorum. Pedunculi breves infra folia in ramusculis nudis, copiosi. Semina plurima abortientia



terminalis Aethras ramulis apice floriferis, foliis longis terminalibus

Habitat in America meridionali

Rami crassitie pennis cygnis ciliatis mucronati et ciliatis  
foliorum deciduorum. Ramuli sparsi approximati versus apicem  
ramorum obtusi. Folia petiolata, usque versus apicem ramulorum  
bifida sive tripollicaria longissima, supra nitida, vix venosa, obtusa  
emarginata, basi acuta. Pedunculi ramuli floriferi ternales apice  
triflori quinquiflori. Flores pedunculati. Pedicelli semper  
culores domentis. Coleola calycina parva, obtusa, tomentosa.

virginicum Medeola



3 cuneata Achnas floribus axillaribus aggregatis, foliis alternis obovatis retusis

Habitat in America meridionali

Rami teretes glabri. Folia petiolato, remota, ad tripolliciana supra  
coriacea, avenia emarginata, Petioli angulati, pollicares.  
Pedunculi axillares, 6-7. pollicares superne uncinatis, longitudi-  
ne petioli, uniflori. Calyx ut in reliquis

a Bumelia retusa Schw. differt. flor foliis alternis

4 Sapota Aehras <sup>2 axillaris</sup> floribus solitariis, foliis ellipticis nitidis.

annona foliis laurinis glabris viridi-fusis, fructu minore Sloan hist. 2.  
p. 171. t. 230. Cat. Carol. 2. p. 87. t. 87.

~~Aehras <sup>gula</sup> trichotoma diffusis, fructu subtortido cicutrifolius matrona brevi~~  
~~Brow. jam. 200~~

Aehras sapota: floribus solitariis, foliis <sup>lanicolato ovatis</sup> ovato oblongis Lin. S. V. pag. 342.  
Jacq. Strep. select. p. 73

Aehras fructu elliptico featro majori, floribus solitariis axillis  
cicutrifolia feminis ultra maturam porrecta Brow. jam. 1. p. 200  
t. 19. fig. 3

Sapota fructu ovato majori Plum. gen. 43

Aehras Loef. 106.

Aehras floribus hexandris Jacq. Am. 57. t. 41

Habitat in America meridionali

Rami teretes. Folia petiolata sparsa, bipollicaria et ultra, utrinque  
acuta, coriacea, tenuis sine venosa. Petioli pollicari. Peduncululi  
axillares solitarii, uniflori superne crassiores tomentosi. Calycis  
foliolo ovata, acuta, tomentosa. Stylus corolla longior

an Sapotilla Brow. hujus varietas Syn. confusenda



5 *Dissecta* *Achras* floribus <sup>confertis</sup> *solidariis*, foliis cuneiformibus emarginato-retusis Suppl.  
210.

Mangrove ana Rheed. herb. Malab. P. 3. p. 53. T. 25.

*Achras dissecta*: floribus confertis corollis octodecimfidis, foliis obovatis emarginato-retusis Boiss. pl. exs. 13. prodr. p. 25.

Habitat in Anfila Americam;

glabra. Boliâ petiolata ad apices ramorum <sup>bipollicaria</sup> conferta, crassezima, concava  
tenuifera venosa patentissima. peduncululi ad apices ramorum foliis  
interpersi; plures pollicares apice crassiores uniflori

balata A. foliis ovato-oblongis subtus incanis, pennis oblongis ovato  
Aubl. pl. Ju. 1. p. 308.

Rheed. Malab. 4. T. 25

non id.