

Alsea

1. spicata A. racemis ovato, fructibus baccatis foliis supra decompofitis
2. mucronata A. racemis longis formis, fructibus baccatis, foliis decompofitis
3. japonica A. spicis longis formis, foliolis compofitis, foliolis marginali
matis

Actea

Habitata America septentrionali herb. Desfont.

glabra. Tolia superne ternata vel snata pinnata: foliola intermedia
subspiciata, oblonga, infima saepe bipartita, terminale trifidum, venosa
subtus glauca, sesqui pollicaria, intermedia superne utroque dente
unico. Racemus subcompositus.

palmata Aitka

Habitat in America septentrionali Richard
Folia simplicia palmata fere Staphisagno.

apera Aetou foliis lanceolatis apertis, spicis interruptis; caule scandente
Laur. fl. Cochinch. pag. 332.

Habitat prope Cantonem 5

Folia subserrata alterna. Racemi axillares. Spicis lanceolatis Cal et
Cotolla 4-fidi. Baccis exsertis uniloculari.

non vidi

2 Spulata A racemo ovato, fructibus baccatis L.

fl. Dan tab 490

nigra Christophoriana Clus. hist. 2. p. 86.

Albidum bacc. ferum Bauh. pin. 103.

albop. Albidum bacc. is niveis Corn. canad. 76. t. 77.

Christophoriana americana racemosa baccis niveis et rubris
Mont. hist. 2. p. 8. f. 1. v. 2. fig. 7.

Habitat in Europa nemoribus; β America 4

Caules erectus, pedalis, simplex, β rarius glaber. Folia supra
decomposita ternata alterna alt ^{caulis} paucis altitudinis
caulis foliola petiolata ovata lanceolata attenuata
ternata subtus pallidiora subpelosa terminale trifidum.
Peduncululus terminalis elongatus flores in apice pedicella
& approximati in ^{racemo} ~~racemo~~ oblongo. Pedicelle inferiores unguic
ulares perenne superius minores pubescentes

2-mucosa A racemis longis, fructibus siliis Len. Spec. pl. pag. 722 v

Christophoriana facie herba spicata Pluk. amalth. 54. t. 383. fig. 3.

Christophoriana americana proenoi & longis spicata Dell. Elbth.
79. t. 87. fig. 70.

Habitat in Florida, Virginia, Canada &

Caules teres corvi bif. tripedati erecti. folia decomp^a ~~partia~~ ternata ^{a duo} partiale pinata inferum ternata
intermedium foliola pinna ovato lanceola
la attenuata glabra subtus pelis minuti inaequaliter
ferrata terminale incipit. Pedunculus caulem ^{pubescens}
termina racemi 3. vel 5 in singulo elongati. Flores
approx. mat; pedicellis vix fimi inqueiculanibus
bractea pubulata ad singulum.

3 japonica A. spec. longifloris, foliis ovatis palmatis indivis. Thunb. fl. jap.
pag. 221 + V. p. 400

Habitat in Japonia

Caulis herbaeum, uti dicta pelantia, glaber. Folia petiolata ternata
foliolo petiolata simplicia, cordata, indivisa 5:7 lobis lobis
indivisi. Petiolus striatus. Flores spica spica saepe spitha
moa Thunb.

pentagyna A. floribus solitariis heptapetalis Walt. fl. car.
pag. 157.
Habitat in Carolina

non ordi

Dicella A. floribus paniculatis, caule suffruticoso Walt. fl. Carol.
pag. 153
Habitat in Carolina

non vidi

monogyne *A. racemis longisimis radiatis* Walt. flor. Car. pag 157
Habitat in Carol.

non radi