

Aphynomene

Aphynomene vero gaudet filamentis s que in singulis corpore Aethas

pubescens. *Caule herbaceo floribus racemosis*
folioli ^{linearibus} *subtus* *subsericeis*, caulis, *superne* *villosis* *aristatis*
caulis mucronati.
Habitat in Guineae Thonung.

Caulis herbaceus, bipedalis, erectus, teres, inferne glabrescens. Folia
petiolata, remota, bipinnata, pinnata cum pan, 16-24 pinnis: foliola
linearia, ^{oblonga} bipinnata, mucronata, subtus glauca et villosa subsericea
supra pilosiuscula. Stipulae subulatae, et pediculi villosi, caducis.
Racemi axillares, bipollicares, laevi, 7-10 flori: pedicelli remoti
bracteae minutae ad basin utraque ad basin calycis, caducis. Calyx
complanatus, 5que dentatus. Corolla minor quam in herbis
lutea: vexillum extus nigro punctatum. Legumen breve pedicellatum
filiforme, congestum, teres, tomentosum, mucronatum, pungens, multi-
loculare, dehiscens: loculi macrocarpi.

herbaria Lyana folii ~~et~~ *occurrit in* *folioli* *subsericeis*
pediculi *subsericeis*
pediculi *subsericeis*

Seneca
musfonda
sericea

G. foliis ovatis attenuatis subtus mollescens, corymbis termina-
libus dichotomis, corollis ~~hyssopifolius~~ ~~sericeis~~ utraque extrusis

Habitat in *Isabela* *Borbonis* *Commeson*

Rami ~~et~~ villosi densi teretes. Folia petiolata opposita
ovata, quadragonelliana acumine longo terminata obliqua
nervis supra pilis minutis adpressis proposita in costis ner-
vis et margine subtus nervis protuberantibus et pilis subveo-
si villosi concipitibus in venis et nervis can qui eade
conspicentes. Petioli ^{vellosi} tomentosi stipula utrinque gemina
uncularum valde attenuata decidua. Corymbus
terminalis pedunculis parvis oppositis bis vel
ter dichotomis villosis. Bractea parva ovata acuminata
ubi calycis et corolla extrusis villosis concipitibus densis
subsericeo tecto. Calyx 5 partitus laciniis brevibus subova-
tis acutis intus hispidis Corolla fere tripolliciana
laciniis lanceolatis attenuatis ^{micromatis} margine ~~ma~~ extrus
et supra intus villosis tenuissimis tomentosis. Tubus intus
hispidus. Gemen oblongum. Anthera 5 lineares sespiles
in tubo corollae. Stylis linearis per hispidus crispus
stigmatata duo altera parum longior

Ophegnomene longifolia caule fruticoso glabro, foliis ovato-lanceolatis, leguminibus
lobis linearibus-acutis Ort. Dec. pag 70

Habitat in nova hispania 5

glabra. Caulis simplex, erectus, teres, purpurascens, nebula glauca aspersus, quadrangulis
datis, ramosus. Folia longe petiolata, dependens, abrupte punctata 7-10 nervis: foliola
cum nervis brevi, glauca, breviter petiolata. Stipulae subulatae decedens. Racemi axillares,
solitarii, foliis dimidio breviores, multiflori. Bractea duae in pedicellis minimis
tubulatae. Corolla lutea. vexillum marginibus involutum. Legumen torulosum
lineare, purpurascens Ort.

Ophiogonoma platycarpus Ruben
Habitat in Carolina Dedit Boppe
Fructus com. praesens, lanceolata tur

pubescens Aphynomene caule herbaceo foliis que lineariis micronatis villosis,
leguminibus compressis marginatis filidos ovibus

Habitat in America meridionali St. v. Prodr.

Caulis teretiufculus, superne subangulatus; crassitie penno columbino, villo
sus, rariopis.

Folia petiolata, alterna, pennata cum impari, pithamboa, cincta
10, 14 piza: foliola linearia, breviter petiolata, unguis vix pollicis
na, interiora minoria, alterna, linearia, apice micronata, basi acu
ta, supra glabra, subtus villosa-subpericea.

Petoli canescentes; teretiufculi, villosi.

Racemi axillares, foliis brevioribus 4-5 flori. Pedielli remoti ungui
culares

Calyx quinque-dentatus: dentibus acutis

Legumina pithamboa, angustissima, glabra, apice subulata aciu
nata: acuminata unguicula ^{sub} pennata

(Racem Aph. lasban differ folioli ^{intermedia} ~~obpatis pithamboa~~ ~~apice~~ ~~subulata~~
foliola glabra differ vero caule folioli que alternis subtus
villosa subpericea

India Aschynomene caule herbaceo, glabro foliolis obtusis enerviis integerrimis, bracteis
ovatis 2 lineis serrulatis, articulis leguminum quadratis
Aschynomene sess indica caule herbaceo lori, leguminibus loricibus hinc forosis obtu-
ris, foliolis obtusis her P.V. pag. 671.

(ab Asch. americana differt caule glabro, foliolis lineari oblongis utraque obtusis, subtus
minime nervosis, integerrimis, stipulis triplo latioribus, enerviis, integerrimis
minime nervosis. Pedunculis terminalibus cum ramulo tandem axillares
triplo latioribus. Leguminibus pollicaribus, disco convexo ad depressum,
glabris; articulis quadratis hinc convexis bracteis lanceolatis, serratis
Petio lori subtus longitudinaliter ad presens punctis minutis solitaneis

Americana Aphynomene caule herbaceo hispido, leguminibus articulis semicordatis, foliolis
acutis ~~nervi~~, bracteis cordatis ciliatis

Aphynomene ~~indica~~ ^{Americana} caule herbaceo laseo hispido, leguminum articulis semicor-
datis, foliolis acuminatis, bracteis ciliatis Len. I. V. pag. 674.

Caulis teres, subtilis sine striatui, adpressis tuberculi minutissimi, cui inf^{er}
peli pilosari, flavescientes, nitidi. Ramuli axillares.

Folia brevia ^{ter} pedicelata, remota, pennata, 27 jugis: foliola linearia, semun-
quivalaria, alterna, sessilia, superne laeva, subtus nervosa: nervis quatuor,
versus apicem ramuli, altero latere angustiori, altero latiore, basi
nervum longiora, versus apicem serrulata. Candcolata,

Stipula utraque semisagittata, membranacea, nervosa, serrata
ciliata, utraque ac candcolata, utraque acuminata
^{in ramulis sive terminalibus}

Racemus axillaris, bipartitus, ~~in~~ sive sive sex florus. Flores remoti.

Pedunculus flexuosus, teres, bipollicari, villosus viscosus: pedicelli
periculi flexuosus, vel uniflorus, villosus viscosus, uti pedunculi

Bractea cordata, nervosa, acuminata, ciliata, pedicello qua-
druplo brevior.

Calyx quadrupartitus: lacinis candcolatis, ciliatis, duobus inter
medios longioribus.

reuecllatum
Legumen populicium, 7. D articulatum: articulis semiovdatis, medis
gibbosis, et siabis

Spinulosa Oschynomene caule herbaceo petiolisque aculeatis, leguminibus cylindricis
ovoidis aequalibus.

Habitatio India Orientali Roing.

Caulis erectus, terres, ramosus, glaber, uti rami et petioli dispersi aculeoli
minutis. Folia tripolliciana, circa 13 juga: foliola opposita vel alter-
na, linearia, glauca viridia, obtusis, mediumata. Stipula linearis lanicola
et attenuata, caduca. Peduncululi axillares, breves, triflori. Flores minores
quam in Oschynomene sesban. Calycis dentis ovatis. Corolla sub purpurata.
Legumen filiforme.

prope sesban herb. hort

Brevifolia *Alphynomena* (Chebucica) *superne* *hispida* *scabra*. Leguminibus hinc repen
di articuli hinc

Habitat in Insula Bourbonis Commerçon herb. Musf.

Pauli erectus, sesquipedalis, inferne teres glaber. Folia semipollicaria: foliola
linearia, oblonga, parva. Stipula lanceolata, attenuata. Pedunculi qua
driflori: Flores parvi. Legumina semipollicaria: articuli gibbosi

rumula Q. caule herbae coarcti, leguminibus hinc serratis medio scabris,
foliis acuminatis Lin. Spec. pl. p. 1061 *

Hedysarum annuum minus ceylanicum, mimosa foliis T. fl. rept. 551.

Habitat in India Q

semipedalis inferne ramosa foliola lanceolata linearia

herb
Guss } Hedysarum ceylanicum minus foliis et sensu minus
Onobrychidis minor ceylanica mimosa foliis articulatis apertis
Mimosa non spinosa minor ceylanica Breyer. cent. 57.

Hedysarum humifusum hirsutum } Plum.
Terram equorum frutescens hirsutum }

scriptura & caule fruticoso glabro, foliis obtusi, leguminibus caribus
obtusis, stipulis acutis deciduis. Sw. prodr. p. 107

Hedyranum arboreum foliis mucosum Plum. Cat. 1. v. 149. fig. 2.
Habitat in India occidentali

Caulis ramisque laxis ramuli pubescentes. Folia petiolata
foliola parva pedunculi biflorum legumina subtripollicia

^{na}
Caulis tenues piperis punctati laxis vel scabri. Foliola parva
legumina articulata: articuli subquadrangulares, ultimo apice ro-
tundato infimo subobovato herb. Kunj

margine externae rorando

Cannabina Q. herbacea, foliis obtusi acuminati pedunculis solita
ris unifloris, leguminibus compressi Corbis Rete. obs.
s. p. 26.

Habitat in India orientali

michi ignota

indica & caule herbaceo lani, Leguminibus lani bus hinc torosis obtusis,
foliolo obtusis. Lin. 1. V

Neli-tali Rheed. mal. 9. p. 31. t. 18.

Cajatis Rumph. amb. 4. p. 64. t. 24.

Habitat in India

Caulis teres lani glaber folia breiter petiolata foliola
numerosa obtusa subtus glauca glabra stipula lanceola
la membranacea pedunculata est apice ramulorum solita
ni triflori legumen pollicare

arborescens & caule arborescens lavi, leguminum articulis semiovdatis glabris
Roxb. Leydb. 304. f. v.

Habitat in India

Specimen sub hoc nomine in herb. Gesf. simile erat grandiflorum. Rami
apice terminati - canescentes. Folia abrupte pennata: foliola 20 puga
obtusae. Scallum floris palmare. Legumina longa compressa.

~~Urtica~~
codiceina \varnothing caule arboreo foliis pennatis. foliolis numerosis lineari-
obtusis pulverulentis, leguminibus compressis aequalibus Lin.
Suppl. p. 330. P.V. p. 671.
Habitat in Otaheite et Ayaheina $\frac{1}{2}$

non vidi

appena il caule herbaeoscibus, leguminum artu uli medio scabis
An. S. V.

minosa non spinosa major zeylanica Raj. hort. 982. Breg. cart. 51.
t. 52.

Aphyomene caule scabro, folio infimo quadrupenni Fl. zeyl. 2904

Habitat in India

Caule inferne ^{adpresso} tuberculis elevatis planis confertis superne
loris prorepentibus Folia breviter petiolata foliola copiosissima
parva linearia obtusa stipula lanceolata ^{attenuata} firmis antheris
pedunculis axillaris bif. Inflori legumina parva ^{bif. 3 pollinibus}
parallela pediculis bif. ~~tripollinibus~~ infimo et extremo
apice rotundato pedunculi pubescente

inferne ^{bus} munitate leguminibus ^{bus} subequalibus

punctata *Clypeomene* caule ^{spithameo} ~~herbacea~~ villosa, foliis obovatis serratis leguminibus punctato

Liabris
Habitat in Lima de Nouis

Rami teretes villosi

Folia petiolata pennata cum impari quadryga foliola ^{terlinearia} opposita obovata
extrosam serrata pilosissima remota

Petiolus unguicularis. Bractea ovata acuta parva

Racemus terminalis erectus tripollicaris. Pedicelli remoti pubescentes
~~Bractea ovata minuta~~

Calycis villosi

Stellum striae purpureis

Legumen unguiculare pendulum articulis subrotundis punctis numerosis
fusis adpressis, pilosissimis

amenciana \mathcal{L} caule herbario hispido, leguminum articulis semicordatis, foliis
acuminatis bracteis ciliatis *Lin. S. V. p.* *Ait. botrop.*
1763. p. 321. 412.

Hedyarum caule hispido, memoso foliis alatis: pinnis acutis mi
nemis graminis *Loan. jam. bot. 1. p. 106. t. 110. fig. 3.*

Hedyarum amencianum ^{humif. fi}
^{sum} *hispidum* \mathcal{L} *Habitat in Jamaica*

Plin. Tournef. Caulis inferne glaber superne hispido subtricipis. Scaber
ins. v. h. rug. Folia presensim petiolata multijuga foliola linearia
articuli ^{minuti} glabr. altero latere scila integerrima altero apicem
hispida
annata serrulata subtus glauca Innervia, seminaque
icularia. Stycula ardo lanceolata valde attenuata
latere altero fetaceo subciliato. pedunculi axilla
longitudine folio axilla pedicello ad pedicellum
flexuosi scabri terti glutinosi bractea semicorda
ta serrulata ad basin singuli pedicelli. Pedice
li uniglori margini articulorum leguminis mucinata
Legumina pollicaria

herba la Manth ill. tab 6 a. g. 121. } credo quod est herb. Lus. Descripta per
Nicos a Ste Bourbon Dutaer

Specimen ex insula Bourbonis dedit Luspan. Caule ramoso pedali. Ramis
terribus, remotis, patentibus.