

Allophyllus

1. regius. A. foliis simplicibus denticulato-serratis, floribus racemosis
2. seylanicus A.
3. cominus A. foliis ternatis, floribus paniculatis
4. nicotianus A. foliis ternatis, subtus tomentosis floribus racemosis

An hujus generis. Hedera trifolia sinuatis et dentatis foliis americana fructu
baccatam costas racemosis per maturitatem albecans. Strato succus indur
americanis dicta Plum. alm. 101

Folia ternata: foliola oblonga, integerrima et remote serrata, subtus villo
herb. Mus.

huc refer Annetia Torr.

medicinalis Allophyllus foliis ternatis, foliolis subsessilibus pilosis, racemis simplicibus folio longioribus, ramis subtomentosis

Habitat in Guinea Thonning 5

Trinex oxygala, ramosus: ramuli teretes. Folia petiolata: foliola ovata, acuta, denticulata: Venella tomentosa, uti petioli. Racemi superne, axillares, filiformes. Flores parvi, glomerati. Bractea minutissima. Flores polygami, hermaphroditi et masculi in eodem vel distincto racemo. Calyx 4-phyllus, compressus, persistens: foliola 2 opposita obtusata concava, arcuata, duo minora, ovata, patula. Petala


4: 4: obovata, intus villosa, gemini opposita. Nectarium glandula 4: compressa, receptaculo cupaeque petali inserto, inter media majores. Filamenta 8 longitudine calycis, basin geminis inserta, marcescentia. Antherae steriles. Germen minimum, villosum omnium. Stylus calyce duplo longior, arcuatus, compressus. Stigma bifidum. Baico binae altera 3 spius abortians: singula magnitudine ribeos, globosa, pilosissima, coriacea. Semen unicum, subglobosum, intra anillum fragile. Flores et Cor: hermaphroditi. Stamina corolla longiora, thalamo villosa inserta, patentia, secunda Thonning

~~an gustifolia Murr. foliis lanceolatis, ^{conymbo} elliptica ebracteata~~

~~Habitat in India Orientali~~

~~Rami superne pubescentes. folia lato lanceolata, petiolata
ita utrinque attenuata supra glabra subtus villosa, utraque
juniora cuneata. Stipulae utrinque duo folia supra
et infra interfoliatae. Corolla exserta villosa, pollinibus
non venis bracteis et calycis fere subulata~~

an lupini generi non modo
varietas frondosa



a corolla hermaphr: magnit: natural; b eadem aucta
 c. flori masculini d petalum. intus e extus f stamen
 g. pistillum
 allophyllus micropus

racemosus Allophyllus foliis ternatis, floribus racemosis. Sw. prod. pag. 62.

Habitat in Montserrat in Ryan's.

Rami aeterni, teretes, cortice cinerascens inferne glabri, superne villosi-subtomentosi, subangulati.

Folia petiolata, aeterna, ternata. Folium intermedium breviter petiolatum, latocuneolatum, subcuneatum 4-5 pollicare: lobus lateralis sessilis, latolobatus, lobis interioribus versus angustiora, intermedio parum breviora; omnia venosa, apice attenuata, utrinque acuta, venis serrata, basi versus integerrima, supra glabra, costa et venis exceptis majoribus exceptis, subtus villosi-subtomentosa, mollis cinerascens.

Petiolus bipollicaris tomentosus.

Rami laterales et axillares, 3-4 pollicares, erecto-patentes, simplicissimi. Pedunculus filiformis villosus; pedicelli breves uniflori. Braea parva linearis ad basin singulorum pedicelli.

Hermafroditici flores

Calyx tetraphyllus: foliis subrotundis concavis, corolla parum brevioribus; lobis duobus parum majoribus.

Corolla petala quatuor, ad aeternum latas, calyce parum longiora, cuneiformia villosa, ciliata. Nectarium diffusum carnosum, obtusum; 5-6 dentatum.

Stamina filamenta 8. corolla breviora. Anthera obtusa
Petalum ~~ne tale, oppositum ab altero calyce,~~
Germen ~~superum~~ Ovium. Stylus ~~reflexi,~~ Corolla ~~longiora~~ et ~~unicui,~~ bifida, corolla
longior. Stigmata simplicia

masculi flores in diffinito individuo
Calyci. Corolla ut in hermaphrod to.
Stamina corolla longiora
Petalum 0.

Pentacarpum globosum, magnitudine pepi, glabrum

caudatus, albophyllus folijs ternatis ^{foliis} glabris serratis ovato ellipticis acuminatis
1 racemosa Schmedelia Lin. Mant. 67x. 514. S.V. p. 375

~~Ursaria longispinis~~
~~Ursaria triphylla Burm. ind. Og. t. 32. fig. 6~~

Habitat in India Orientali

Arbor ramis teretibus glabris canescentibus, albido punctatis. Folia
petiolata alterna, remota, ternata. Folia latera petiolata, ovata
bipollicata bi f. 3 pollicaria, lateralia minora, venosa, acuminata
exteriora subserrata, subtus pallidiora petioli pubescentes supra
canaliculati. Racemi axillares, solitarii spithamoi, filiformes.
Flores minuti brevissime pedunculati plures ex eodem puncto. Bractea
glaber magnitudine pif.

Schmedelia racemosa Lin. Mant. 67x 514. S.V. p. 375
Ursaria triphylla Burm. ind. Og. tab. 32. fig. 7

~~apocorteri~~ alloph. foliis ternatis folioli ovati utraque glabris
~~ternata~~ meliopo foliis Forst. gen. 20.
~~indeggifolius~~ Habitat in nova zelandia ^{h. ang. velipul}
Rami teretes flexuosi ~~pubescentes~~. Folia petiolata, alterna, ternata
Foliola ovata, indegemma, glabra venosa ses qui pollicitari ^{in fest}
Racemi axillares solitarii longitudine folii flores ~~proplei~~ ^{proplei}
Solitarii minuti pubescentes
Arbor facie Rhos

~~Habitu accophylli diversa videtur stylo bene indurata~~
Aponetia ternata Forst

Zeylanicus Al. foliis ternatis oblongo ovatis serratis ramis que superne villosi racemus
simplicibus lateralibus subsessilibus
Rhois in foliata frutex salvaicensis, floribus juliformibus Petrus in
670. Ray. Deendr. 50

Allophyllus zeylanicus in fl. zeyl. 1404 S. V.

Rhus cobbe foliis ternatis: folioli ovatis acuminatis serratis pedunculis
et tomentosi in S. V. pag. 293.

Habitat in India orientali

Rami teretes superne villosi nec tomentosi. Folia petiolata
alternata ternata: foliola oblongo ovata seque pollicem latera
lata ^{subsessilibus} de medio minora basi indentata latera angustiora et obliqua
omnia serrata remote serrata serraturis mucronatis acutis
puncta supra saepe villis raris adpressis pubescentia intus
glabrata laeviora, subtus villosiora tomentum non molle, pallidius
ra - petioli angustulato villosi Racemi axillares folio
parum longior. Peduncululi filiformes villosi a medio ad
apicem obtusi floribus copiosis confertis brevibus magnitudine
superi villis raris adpressis

Polygama dicia deserv. partem fructific. vide in specimine herb.

Coble A.

ternatus Allophyllus foliis ternatis serratis, racemis terminalibus
longis Lour. fl. Cochinch. p. 232

Habitat in Cochinchina

arborescens spatulata. folia magna inaequaliter serrata
flores parvi albi. Petala⁴ minima pilosa Nectarum glan
Dialo 4. Stylus bifidus. Germinum didymum

non vidi

aequalis A. foliis ternatis foliis petiolatis ellipticis serratis subtus tomentosi basi
subaequalibus, micis axillaribus

Molago maram Rheed mal. 5. pag. Tab. 25.

Habitat in India orientali &

Rami teretes inferne glabri punctati, superne lentes tomentosi. Folia per
petiolata alterna ternata. foliola petiolata, ~~et per pollicaria elliptica~~
^{ut nunc} acuta ~~intermedium basi~~ versus magis attenuatum et acutius
serrata serraturis micronatis, veposa, basi aequalia supra glabri nervis
tantum villosa subtus pallida villosa mollia villosa crenatifida
venella utraque. Micis axillares petiolo parum longiores.

inferna herb. M.
subcomp. p.

Petoli bipollicares parvales ut semper unguiculares intermedium
parum longior omnes tomentosi. Flores copiosi conferti masculorum
tantum vidi cum reliquis speciebus convenienter in oracibus

Seylanicus Al.

4 racemosus Allophyllus foliis ternatis, ^{subtus tomentosis} floribus racemosis Rich. prod. p. 62.

Habitat in Hispaniola ~~et adriati coaste, censeo~~
 inferne partes radii et ~~sub~~ ^{sub} ~~opomeno~~ ^{opomeno}
 Ramis superne angulatis, villosis. Folia petiolata, alterna. ^{attenuata} Foliola sesplea

lanicolato elliptica, ^{venosa} ext. opposita ^{vel parvis utrinq. superne} dilatata, utrinque ^{attenuata} acuta
 remote denticulata, ^{venosa} ~~subtus~~ mollesima bif. ^{ext.} Incolliquis. Racemi

latensia ^{boerha.} axillares petiolo duplo longiores. Flores brevissime pedicellati: cernui
 basi interiori angustiori

ellip. folioli ellipticis

sesplebis, racemis sim

reliqua vide inter descript.

an Cerasus trifolia Plum. herb. Guss.

Unifoliata Ornithopis

Habitat in Insula Francia Fusca ^{mont. de Fusca} ^{monti elevati parvo conidentibus de fusca}
Rami teretes, glabri, cortice cinereo. Folia petiolata, alterna,
ternata. Foliola elliptica, petiolata tripollicaria, utrinque glabra
venosa, subrepanda, ^{si per apicem fere perpendicularia uny vel altero puncto} ^{dele 4.} ^{interiore} ^{obliqua} ^{tamno uno vel altero} ^{per quibus decidua minutis} ^{acuminata basi acuta, lateralibus parvis}
minoribus, altera latere breviora. Peduncululi axillares, inter
dum gemini quadrupollicares, longitudine petiolonum, e pediculis
persistentibus brevibus reflexis squamosi. Flores in racemo simpli
ci vel composito

Allophyllus ornithopis

Folia ternata: foliola elliptica glabra ut subintegerrime
petiolata micis subcompositis.

an Schmedelii generis

angustifolius ^{ternatis} ~~foliis lanceolatis~~ ^{foliolis} lanceolatis extrosum serratis utrinque
glabris, racemis simplicibus

Habitat in Cap. b. p. i. & Dr. Buloni

Rami alternis, tereti rugidi inferne ~~et~~ glabri inferne cortice purpureo
siccitate fusco cinereo punctato, superne cinerei ~~ovato~~ ^{velis} ~~et pilis~~ ^{minutus}
adpressi angulati. Folia per versus apic. summitatem rannonum
alternata, ternata: foliola pollicaria, sessilia, utrinque glabra

basi subobovata

utrumque extrosum ^{obtusis basi acuta} versus serraturis utrinque vel 2 saepe in
gemina intermedium lateralebus paulo longior ex utroque adde
mentem in primo basi lateralia interdum ^{curvatura} obtusum. Laterale
paniciforme ^{paniciforme} racemi axillares longitudine foliorum simpliciter
Peduncululi villosuli pedicelli parsi approximati breves

angustifolius A. Folio Ternatis

(nigrus Allophyllus foliis simplicibus denticulato-spinosis, floribus racemosis
Schw. prodr. p. 62.
Habitat in Hispaniola }

axillaris Cellophyllum foliis ternatis glabris. foliolis obovatis integris dentatisque.

Habitat in Cap. b. sp. 17

Arbor facie Arboris ramis purpureo-fertibus griseis apice angulatis cinereis. Folio-
petiolato tripartito in foliola sessilibus basi attenuata, obtusiuscula integra
interdum superne dente uno alterove obtuso angulata pollicaria
lateralia minora subtus pallidiora. Racemi axillares solitarii
folio brevis. Pedunculis pubescentibus. Flores conferti parvi. Calyx
tetraphyllus foliolis suborbundis concavis glabris. Petala obovata longi-
tudine calycis. Polygama Stylus bifidus.