

Anabasis

1. aphylla A. aphylla, arboribus emarginatis
2. foliosa A. foliis subulatis, floribus axillaribus confertis
3. tamaris ^{posita} A. foliis subulatis, ~~penicillis~~ ^{confertis} floribus axillaribus
4. spinosissima A. frutescens ramis nudis spinosis fere

Arabasis

Funosopia Arabasis Len. l. v. pag. 264. Symb. bot.

Salpola echinur: fruticosa glabra, foliis subulatis, multivis, spinis divan-
cati floriferis Billard. Ic. dec. 2. pag. 10. t. 5

herb. Desfont. Specimen a Billard.

Rami sempredales: ramuli patentissimi & pinnatifidi

integri folia areolla foliis simplicibus ^{integerrimis} oblongis petiolo inferne dilatatis ciliatis ~~vaginantibus~~

~~Vetus~~
Aroella ^{filamentosa} umbellata L. Martio Encyclop.

Terra del fuoco
Habitat ad praedum magellanicum ¶

Caule inferiori ~~veteri~~ basi stipularum vaginantium persistente. subramosis
folia alterna petiolo ^{amurquialana} superne ^{approximata}, inferiora subovata acuta
superiora angustiora fere linearia ^{semit} glaberrime crassiuscula revoluta
nervone albo terminata subtis seta una alterave bilineari adpressa
pleniusque in costa folii. petiolo superne ^{semit} inferne dilatato
ovatis ^{ter} vaginantibus imbricatis margine ciliis aequot. ligulis albis.
umbellae laterales pedunculatae simpliciter. Pedunculi pollicares
Involucrum pentaphyllum foliolis ovatis acutis setaceo ciliatis. Pedicelli
unquiculus involucrio duplo longioribus, 12-13

Chorombella simpliciter involucriata Petala integra Capsula bicocca. Semina solita
na

aphylla A aphylla, articulis emarginatis Lin. S. V

Salpola baccifera falcicornis Jacq. Gmel. hb. 3. p. 101. t. 10.

Troali bacciferum, falcicornis Jacq. Burb. cent. 1. p. 11. t. 10.

Habitat ad mare Capri littora et circa Trapanum

Caulis erectus paniculae ter glaber ~~aphyllus~~ ^{ramis, ramulique oppositi.} articulatissimus
flexuosis articulis pollicibus in ramulis paucis minoribus, folia
nulla nisi squamula minuta utrinque ad singulum articulum. Flores in
ramis ~~et ramulis~~ ad articulos in ramis remotioribus in ramulis ultimis im-
bricati oppositi. Petala suborbiculata

Cretacea Anabasi aphylla, caulis numerosi cespitosi simplicissimi
nisi articulatis basi floriferis: articulis emarginatis Willd
Spec. pl. pag 1370.

Anabasi cretacea Pallas et. 1. app. n. 109. tab n
Habitat in Siberia 7

foliosa a foliis subclavatis

Salsola foliis uncinatis obtusis, supra planis, infra convexis Gmel. pb.
n. p. 99.

Trali baicifera, foliis clavatis Burth. cent. 1. p. 19. t. 19. fig. 1.

+ Habitat inanis Cappi littora ☉

+ *Salsola baicifera Pallas*

Caulis erectus teres ^{semipedalis} purpurascens glaber ramifer. rami alterni frequentes
nollicares simplicis ^{compositi} spiciferi, foliis ^{compositi} serotina apice uncinata
carnosa unguiculata. flori axillares ^{compositi} sessiles plures. Petala 5 ungue
oblonga ^{sub} ^{robenda} ^{membranacea}

tamanziifolia Anabasi foliis subulatis, Pericarpis exsertis Linn. Spec. p. 324* S. V. p. 264
Vulgariter ofum Hispanicum, tamen Eufolia Tournef. 4. Pl. 4. 247.

Habitat in Hispania,

Trunc ramulis albis, glaberrimis. Folia subulata, triquetra, Tamanzi. Flores
subsiccati, solitarii, axillares. Calyx triphyllus, major. Petala 5. ovata, calycis
minora. Stamina 5. corolla longiora. Geromen conicum, desinens in stylum
subulatum. Stigmatibus 3 bus obtusis Linn. Spec.