

Cnrorogon

pedunculato
compressis sed
flo. tubulorum
dilatis

Pulmoni neti Di glabra tener

in loquax

Simplex.

Andropogon

Habitat in Guinea Thonung,

Pedmi pedales, ²Varices, filiformes, ~~parva~~ inferne parum ramosi: Rami
simplices, recti. Folia ^{remi} perfollicaria, linearia, basi ^{remi} raris pilosa, margine ^{remi} scabro
pula. Pedunculi ^{remi} inferiores duo ^{remi} tres a singula vagina ^{remi} superior eo solitarii
subsesquialter vel brevissime pedicellati. Involucrum ^{remi} lanceolatum; ^{remi} pro
pollinaria ^{remi} sub ferrugineo-rufoescens, glabrum. Spica solitaria, basi involucri
in partae apice ^{remi} que longitudinis ^{remi} ~~per~~ ^{remi} flores ^{remi} ~~quatuor~~ ^{remi} panum. Rachis sulcos pilos a

quatuor panum
glumae calycinae
hermaphroditi in
Sedam attenuatae

~~remi~~ ^{remi} flores lanceolati: ~~remi~~ ^{remi} ~~per~~ ^{remi} ~~quatuor~~ ^{remi} panum. Rachis sulcos pilos a
hermaphroditi vix pollicari, geniculata, glabra
masculis mixtus. Andra

canicosum A. spica solitaria imbricata, seminibus hirsutis: anthesis nudis
contortis Lin Spici. pl. pag. 14004

Gramen canicosum Rumph. amb. 5. p. 17. t. 7. fig. 2. A.

Habitat in India.

culmi filiformes. folia vaginis brevioribus pilis raris adpressa.
Spica terminali bipollicari spicula hirsuta nitida antheris
spicula duplo longior.

Andropogon panicula pedunculis simplicissimis 9 floris, fasciculo herma-
phrodito aristato, rachide piloso

Habitat

Isent

Culmus ramosus: rami semiteretes. Vagina lanceolata, valde attenuata, spi-
culis longiores, spathacea. Pedunculi superiores pilosi, flori 9 flori: pedicelli
triflori, alterni. Flores intermedius sessilis, hermaphroditus, aristatus, aristo
longa teretis, vane curvata, villosa: laterales pedicellati, steriles, vane mi-
tici. Valvula glabra, lineares.

undatum Andropogon

Andropogon undatum Jacq. coll. 3. pag. 237

Habitat in insula Mauritii 4

Caules plures ramosissime tripedales, graciles, teretes, ad Centrum villosuli
repentes, ex omnibus nodis radice antea, gemisuli inaequaliter hirsuti
folia lanceolata acuta, et versus scabra margine, undata, bipollicaria
ad Centrum villosula. Rachis 4-5 uncialis, ad singulam spicam hinc con
serta illius plana. Spica alterna 2-3. subsessile, patens, fere
unicolor flavo-rosea. flos perianthi unica per gemini flos ^{obivalvis} hypogynus
superior infimus inaequaliter pedicellati gluma calycis ^{obivalvis} torato ^{obivalvis} spica villosula
dorso venter areolata infra apicem aristata acuta colla brevior est
non arista longa. valvula collae exterior aristata maribus herme
phrod.
facies una ut panic: all.

Cymbanum A. panicula sparsa, bracteis cymbiformibus, floribus trans-
versis aristatis Terni Lin Mant 302. & P. V. pag. 903.

Habitat in India orientali

Culmi teretibus, teresibus, laxis, erectis, maximam partem secti
foliorum vaginis, crassitie pennae. folia grammæ pedalis, mar-
gine striata, plana ^{base teres pilosa.} vaginis laxis, ore barbato. Panicula
ex utriusque foliorum superiorum in unam dispersa, plurima, pedi-
cellis capillaribus, unde tota panicula pedalis. Bractea tam
quam Calyx communis: speculis ^{erectis} geminis alter sessilis quadriflorus
alter pedicellatus biflorus, cymbiformis sive ovata adnata
tonicava membranacea purpurascens, transversa posita,
et oppositi subula & ma' her maphiditi sessiles aristati subsessili
masculi angustiori glabrispecula basi barbata

comparatur cum aristata gigantea Parva

Bladhii A-^{pr}ici febriculis: flosculo hermaphrodito sepli an stato, neutro pedun
culato, diluato mutuo Retz. of. 2-pag. 27.

non vid.

Cymbachne Andropogon, spicis conjugatis valvula exteriori linearis flosculi
hermaphroditi linearis interiori cymbiformi anisata
Cymbachne ciliata Retz. obs. 6. pag. 36.

Habitat in India orientali Nonny

Gramen Tenue, culmi plumbis fm plumbis ramosisque. Folia
brevia tenuia, oramine supra vaginam pelis longis difixitis
albis ciliata. Vagina truncata, ore pallido fuso, tubulata.

Spica gemina e vaginis supremis, sequi pollicares. Rachis
flexuosa. flosculi bini alter flosculi hermaphroditi alter
pedunculatus muticus basi barbati. Pedicellus singularis ^{antheris} ~~complanatus~~
latus latius superne truncatus, compressus altero latere convexo ^{exteriori}
ultratus flos valvula exteriori flos hermaphroditi cymbiformis dorso
ciliata, latere rachis obvertens, ^{emarginatus} supra interiori in linea pro
compressa ^{annata} fundata exteriori dorso opposita recurvata. valvula flosculi
attenuata concava convexa lanceolata

tenuiflorum A. specis conjugatis, flokulis hispatis

Habitat in India orientali

Facies omnino Schouanthis et forte tantum varietas

racemosum And. spec. i racemosi flosculis basi villosi.

Habitat in India orientali

Culmi adscendentes basi ramosi. Folia lanceolata bipolliciana.

Pedicelli pro capillares oppositi floris parvis.

provincialis a. fructu fasciculatis loricibus, floribus alternis subsplebeis
secundis aristatis Retz. obj. 3. pag. 43.

descriptum a La Martini provinciale
arundi-

incurvatum. A. spicis filiformibus subdigitatis, floribus gemmatis: foemineo
pedicellato antherato, masculo sessili multico, calycibus univalvibus
Pet. obs. 5. p. 21.

Habitat juxta flumina. Torquebaria

non vidi

geminata. ^{lobis} plicis conjugati divaricati flocculis basi barbati

Habitat in cap. b. pui

Fasci And. distachyi vel Schonanthi an eadem. At flocculi
multo longioribus
rachis vultu barbata valvula glumae non subtilissime striata

afes A. species subternis Terminalibus, flospulis omnibus anfractis Brui.
A. 5. app. pag. 47. lam. 1. cone

non vid.

22-23. annulatum Andropogon species digitatis plurimis, flosculis obtusis sessilibus
an statis, pedicellatis mutis, geniculis barbatis

Andropogon annulatus, species ad apicem culmi confertis, alternis,
floribus gemmatis; an statio lespli hermaphrodito, pedicellato
mutico inani Boiss. descr. pag. 173

Ex India Orientali misit ^{Dr} Königius

(Statura et summa affinitas A. Schoeni et facile eandem ducere
Differt autem geniculis valde barbatis, ligula foliorum membra
natae nuda, nec vagina; ore piloso, flosculi minus pilu-
glabros, glumis calycinis obtusis nec laniceo latius acutis

ramosus A. spicis terminalibus conjugatis spiculis geminis: macula mutica
pedicellata, culmo ramoso Forst. descr. pag. 173

multi haud satis notum

brevisfolium A. fructis lateralibus solitariis, flozulis alternis remotis arista
tis, cu' hmo geniculato compresso Sw. prodr. pag. 26.
Habitat in Jamaica.

mibi ignotum?

~~infernum~~ ^{imbriatum} A. spica simpliciter imbricata, sepalis binis florum anfractis septicim
binis multo pedicellatorum, calycibus univalvibus Retz obs. 5 pag
21. sub nomine Ad: semati
Habitat in India orientali.

michi ignotum

Dulcia Andr. spica folitana imbricata, flosculis muticis Burm. in
pag. 219.

Cypripedium dulci Rumph. amb. 6. p. 7. tab. 3. fig. 1.
Habitat in Andia orientali

varietas β Burmanni alia planta

non vidi

municatum And. panicula contracta glomerulis geminis mucronatis
muticis

Phalaris zizanioides panicula simplicissima, filiculis mucronatis, altero sepli Lin Mant. 103

Andropogon municatum

Andropogon squarrosus

Habitat in India orientali

Calamus bipedalis et ultra folia angusta cannata retroflexa
scabra, glabra. Glomeruli subulata margine utraque
aculeis

Diphysa florum Andropogon

Habitat

Ded. Richard

Distachyum A. spicis binis terminalibus, culmo indiviso Lin. Spec. pl. p. 14814
Ger. fl. Gall. prov. 106. t. 3. fig. 2.

Gramen bicorne f. Distachyophoron Boic. sic. 20

Habitat in Europa australi, Barbaria

Culmi plus quam pedales. Folia plana. Nervi nulli. Spicis duo
aequales, oblongis, exserti in fiula, culmum terminantes: singula
ex flospulis viginti pluribus. Pedicelli aut brevissimi, planiusculi:
basi pilosi superne latiores. Flores solitarii. Calycis unisp
gluma exterior flore fere longior latiorque apice bifido:
anfra longa, articolata, patens. Gluma calycina interior
minima Lin.

Figura geradi.
aut mala aut
a planta mea
diversa
Sub nomine Gram. Andr. provinciale gramen a Thounio accipi
tunc forte pertinens Culmi et folia ut in Lin. Spicis gemino
tripliciter. florei gemini alter pedicellatus alter sessilis.
glumae biflorae foeminae (forte hermaphrodite) anfrata anfr
longa tortili gluma exteriora calycis uti pedis ractu pedu
culi et ractu pedis longis
and. v. prozom provinciale
Ritney affinis barbato
Inciis 15-20 a mea prozom.
Drochun
floribus geminis caly

Andropogon rupestris

Habitat in Puerto Rico dedit G. Don

pelosum Andropogon ^{Digitatus} specie fasciculatis hirsutis, floribus omnibus aristati herme-
phroditi sessilibus, foemineis pedicellatis
Habitat in India Orientali Rottler

Culmi teretes, glabri, in ferne radiantes, an basi repentes. Folia ex-
guata, striata, glabra. Vagina striata, folio longior, supra spathe-
ica, terminata folio convulso, quadrilobum, glabro, striato. Lig-
ula membranacea. Spica terminalis quadrifida, fere 3 pedunculata: ca-
va duo, ubi glumae ^{catyicis} ^{2^{ae}} villis albidis hirsuta. Calycis glumae, bivalves
uniflorae: salvalte altera conica lanceolata, altera conica corollam formans
altera plana angustior. Glumae corollae longitudinae fere calycis, gla-
bra, altera dorso ^{2^a} arista recta; arista recta longitudinae flosculi pediculis
longior. Styli duo apice penicillati.

Schoenanthus a paniculo spicis conjugatis ovato-oblongis, rachis pubescente, flo-
lis sessilibus anthera totuosa L. V

Junci odoratus aromaticus. Baub pin. 11. Theatr. 163.
gramen ad juncum accedens aromaticum majus sijnacum
mon. hist. 3. p. 229. f. O. V. g. fcy 25.

Gumen dactylon aromaticum, multiplici panicula, spicis
brevibus. Yomento candidantibus ex eodem pediculo binis Plin
alm. 175. V. 190. fcy 1.

Habitat in India, Arabia

~~in mea a Linnao spicis enim lineari nec ovato-oblongo~~

^{non}
^{olea}
Sagena ^{supernona} convolutata spicis solitariis vel geminis ^{lineari}
ses viri pollicari, flori gemini alteri sessilibus ^{hermaphroditi} antheris
Cana alba, spiculis flosculis glabri

alternatus
nascitur

Folia plana lanceolata attenuata

Rumph. 5. V. 72

lineariibus flosculis ^{glabri} geminis alteris
sessilibus antheris

Schoenus niger Andropogon ^{Andropogon} paniculato spicis conjugatis ovato-oblongis
rachis pubescente, flocculis sespibus anfracta tortuosa Lin. L. V. pag.

904
Andropogon beornae Forst. descr. pag. 173.

Castrogratum A. panicula spica solitaria, pedunculis elongatis subcastanea
tis, rachis lanata, flocculis aristatis, masculo sterili Sw. p. 102 p. 26
Habitat in Samalia

non vidi

bicornis A. panicula spicis conjugatis, pedunculis ramosis, rachis lanata,
flosculis anthera adula, masculis tabescente *Lin. l. V. Sw. obs. p. 302*
gramen dactylen bicorne tomentosum maximum, spicis numerosis
Sloan hist. 1. p. 42. t. 15.

Habitat in Brasilia, Jamaica.

gramen altum culmo crassius penno asperno superne subtusque
lavi glaberrimo. folia vixnanta longa, convoluta, margine
Panícula termi ^{triginta} semulato fiabra glabra. Pedunculi ^{numerosi} atri rami pedunculorum
natis fastigiat ^{foliis} ^{ovariis} ^{saepius} ^{partiales} ^{alterni} ^{filiformes}. Spicula
he Dale ^{solitaria vel glomerata} ^{venues} ^{rachis} ^{filis} ^{longis} ^{barbata}. Spicula ^{angustissima}
Flosculi ^{polygami} ^{herma} ^{vellum} ^{longum} ^{occultati} ^{herma}
phrod. ^{subspiculis}. Masculi 12. ad latera, pedicellati, multo mino
20

glaucom A. panicula foliacea: involuclis calycibusque bifloris, flo-
rum fere omnium calycibus trivalvibus, pedicello tanquam bivalvibus retru-
so. *obf. 5. pag 20*
Habitat in India orientali.

alopecuroides α panicula laxa, nichi lanata, glomulo utroque anxia tor
taosa lin l. v. p. 904. *

Gramen Dactylon alopecuroidis facie, panicula longissima & spicis
plurimis tomentosis constante Sloan bot. 1. p. 113. ~~44~~ 470
fig. 1.

Habitat in America septentrionaliore.

Canis glomulis longior ^{etiam} in ipsis glomulis

non vidi

undatus A panicula erecta ramosa: spicis alternis subternis, & costatis
ternis, mapulo inferiore proae laequ. coll. 3. pag. 230

non vidi

Aurum Q. spicis fasciculatis quibus herpetiformis flosculis geminis omnibus
anctis

Habitat in insula Bourbonia

Culmi proque pedales; erecti; basi ramosi. vaginati. Sagina ore ciliata
Pedunculus superne pilis adpressis. Spica tripolliciana e
pilis longis fulvis mollibus herpetiformis. Flores gemini
alter sessilis alter pedunculatus ambo hermaphroditi anthera
4. anthera longa. Pistilla penicilliformia flosculo longiora

Nardus A panicula ramis supra decompositi prolesens fl. reyl. 457

P. V

Calamus odoratus mathioli Bauh. pin. 17. theatr. 263.

Habitat in India

magnitudo anandi, Culmi Tenues tenuissimi articuli inferne vaginati glabri vagina longitudine foliorum uti folia, glabra. Sola angusta. Panicula terminalis composita inferne magis composita palmari. Bractea vaginans ad basi ramorum et pedicellorum hinc unguiculatae floris imbricatae glabuli ad basi barbatae altera aristata

Gryllus Andropogon

Aegilops bromoides, suba purpurascens Baub. hist. 2. p. 436. Scheuchz.
gram. 267.

Gramen spartum festucum seu Aegilops spartea villosa Barr. ii.
10. f. 2.

Andropogon gryllus; paniculae pedunculo simplicissimis trifloris:
flosculo hermaprodito sessili aristato ciliato basi barbato Lin.

Amoen. acad. 4. p. 1332. Syst. veg.

Habitat Helvetia, Verona, Montpelie &

Culmi aggregati, simplicissimi, pedales, bi- & tri-nodi, glabri. ^{infracto vix} ~~infracto~~ geniculis inferioribus ^{infracto} ~~infracto~~
radicalia, duo vel tria in culmo eoque breviora, angusta, pilosa. Pedun-
culi sessilibus, simplicibus, interdum unus vel alter ex inferioribus bifidus; ca-
pillares, raris pilis, apice subflosculo pilis helveticis barbati. Glomeruli
tuberculati: vix unguiculares. Glumae calycinae laeviflosculorum masculinorum
laeves, in unam brevem attenuatae, glabrae, superiore subciliatae; pedi-
cella flosculo foemineo duplo breviora. Corollae hermaproditae salivula
exterior duriuscula, flava, margine tuberculata, tuberculis piliferis, inter
dum linea tuberculorum quoque in dorso. ⁱⁿ ~~in~~ Arista longior major flosculo
duplo longior ab infra geniculum villosa; salivulae aeternae brevior, recta

in interior hyalinum
ciliatum

aciculatum And. panicula contracta erecta, pedunculis trifloris, floribus majoribus
bis pedicellatis acuminatis, foemineo sessile aristato Retr. off. 5. p. 22.
Gramen aciculatum Rumph. Amb. 6. p. 14. V. 15. fig. 1.

Habitat in India orientalis fertilebus & nederatis

Culmi pedales basi radicanter ascendentes, simpliciter flavescente
tri-quadrinerviati geniculis fuscis. Folia plurima inferne
bi-tripollicaria linearia glabra obtusiuscula margine & scabra
vaginae inferiores & internodiis longiores compressi. Pedunculi
verticillati, capillares flosculi minores quam in Gryllo, basi ^{vallis} pubescentibus
natis flavescens brevibus, an ~~se~~ hermaphr. Valvula arista anistae
brevior ^{major} p flosculo duplo longior flavo ore lentis denticulatae
valvulae margine ciliatae pedicelli superne flavosca, helvolo
vellosi