

Apium

1. graveolens A. foliis pinnatis. folioli cuneiformibus subtrifidis incisi
2. antarcticum A. foliis pinnatis. pinnis omnibus subtripartitis incisi  
triplois utraque acutis
3. petroselinum A. foliis pinnatis pinnis radiatis  
caulisem lineantibus, involuclis minutis

Clp. antarcticum } Banks  
decumbens }  
microphyllum }

*Apium*

*Apium creticum*, minimum, Anisi facie Tournef. Por. pag. 21.

Habitat in Creta herb. Tournef. O

Caules plures, teretes, ramosi, tenues, glabri, laeves, semipedales, erecti. Folia radicalia longe petiolata, cordato pinnatifida, unguiculata, crenata, glabra: caulina ternata, inferiora laciniis lateralibus lanceolatis, intermediis obovata, extorsum serrata superiora linear-lanceolata, integerrima, unguiculata, acuta, suprema pinnulata. Involucra et involuella nulla. Flores parvi, radiati, fertiles.



nodiflorum Apium

Habitat in monte vides Comnes-herb Guss.

Caulis teres, striatus. Folio glabra, uti tota planta, radicalia bipinnata: pinnis pinnatis sed & inciso; lacinis linearis: caulina folioli 3 partiti: lacinis linearis inter gemmas. Umbellae laterales, sessiles: foliolum parvum, tripartitum sub umbella umbello <sup>uniformes</sup> fertiles; - Semina globosa, membrana tenui alata

Australe Apicum

Habitat ad fretum majellanicum Comest. herb. huj.

Caulis teres, striatus, crassitie pennis an senis. Folia superora petiolata, pennata, 3-4  
paga: foliola petiolata, 3fida, incisa, glabra, venosa, a cutu, semi-pollicaria.  
Petioli basi membrana aucti. Pedunculi e dichotomia. Umbella radii 9 fer  
tiles, uniformes

Apium

Apium sativum minus sicut Lametula Tournef. Cor.

Apium creticum minus crassifolia Tournef. Cor.

Habitat in Creta herb. Mus.

Caulis interdum subfistulosus. Semina laevia.

habes



bipennatum A. folius caulicis bipinnatis, folioli crepso serratis Walt  
fl. car. pag. 115.  
Habitat in Carolina

non vidi

anartheum Q. Folis pinnatis, pennis omnibus subtripartitis inis  
umbellae pedunculatis

Habitat ad fretum <sup>glaber</sup> magellanicum <sup>sepiotupm</sup>

Caulis striatus. Folia ~~pedunculata~~ <sup>glabra</sup> foliolaque petiolata  
folia petioli <sup>sepiotupm</sup> striatus vagina latens  
membranacea foliola <sup>sepiotupm</sup> ~~operta~~ petiolata glabra lobis  
subtrifidis <sup>sepiotupm</sup> laciniis mucronatis venosa. Petioli <sup>sepiotupm</sup> ~~longi~~  
umbellae laterales et terminales multiflorae <sup>sepiotupm</sup>  
affinis graveolenti, at foliola bipartita trifida nec  
trifido <sup>sepiotupm</sup> ~~feratis~~ quadripartita nec <sup>sepiotupm</sup> ~~minor~~  
cuneata <sup>sepiotupm</sup> utrinque <sup>sepiotupm</sup> ~~excisa~~



gracile A. foliis caulinis cuneiformibus Len L.V

Fl. Dan. 790

Apium palustre f. apium officinarum Bauh. pin. 154.

β. Apium dulcis Celeni Italorum Tournef. 4. R. 4. 305.

Habitat in Europa humectis, prope mare maritimi &

Caulis erectus vel decumbens ~~fruticatus~~ glaber interdum altitud.  
homini subcatus. Folia ternato vel quinato pinnata  
folioli inferioribus petiolatis superioribus sessilibus basi  
acutis ~~exterioribus~~ ~~latius~~ ~~incis~~ ~~infusis~~ ~~umbellis~~  
~~petiolatis~~ ~~terminalibus~~ & ~~lateralibus~~ ~~sessilibus~~ ~~pedunculatis~~  
latis &c.

pinnati <sup>fol</sup> caulinis cuneiformibus incisus umbellis latera-  
libus sessilibus

*Petroelinum* A. foliis caulinis linearibus, involuicellis minutis Lin. S. V

*Apium hortense* Dod. pempt. 694.

*Apium hortense Petroelinum* vulgo Bauh. pin. 153.

β. *Apium* vel *Petroelinum crispum* Bauh. pin. 153.

Habitat in Sardinia juxta Scaturigines ♂

Caulis erectus terribitalis sulcatus. Folia radicalibus  
pinnatis pinnis oppositis petiolatis ovatis serratis  
in medio sesquipedalibus caulinea linearia laniceo  
tri sesquipedalibus paucis serratis. umbello terminalibus  
et oppositifolia

radicalibus pinnatis pinnis ovatis serratis

an folia radicalia quoque bipinnata