

Biscutella

1. *aureolata* B. Calycibus nectaris et nique quibus, filiculis in stylum  
locantibus <sup>quamops</sup>
2. *lorigata* B. filiculis glabris a stylo distinctis, foliis lanceolatis <sup>sema</sup>
3. *apula* B. filiculis scabris, foliis lanceolatis sessilibus <sup>sema</sup>
4. *lyrata* B. filiculis scabris, foliis lyratis, caule subaphyllo
5. *coronopifolia* B. filiculis glabris, foliis dentatis hertis
6. *sempervivens* B. filiculis glabris, foliis lanceolatis tomen  
tosis, caulibus suffruticosis.

peruviana Bisulcula

Habitat in Peru herb. Gusf.

Caules herbarius, erectus, teres, crassitie fere penno asperino. Solia frequentia, inferiora petiolata, superiora sessilia, sesquipollicaria, oblonga, basi angustata, <sup>venosus</sup> <sup>subtus</sup> ~~venosus~~ <sup>subtus</sup> subnervosa, areola, acuta, <sup>ex</sup> <sup>superior</sup> <sup>dentata</sup> basi versus integerrima, <sup>subtus</sup> pallida, suprema lanceolata. Racemi superne, alterni, distantes, cum pedunculo tripollicares. Pedunculi pubescentes. Siliqua duo distincta, orbiculata, membranacea, venosa, glabra, orbiculata, integerrima.

Cynati B. filiculis scabris, foliis bipartitis Lin. Mant. 254. & L. V. p. 592.

*Rhaphani bipartitum raphani s. inonis folia* Boerh. ju. 45. t. 23.

Habitat in Hispania, Sicilia, Barbana C

Folia plurima radicealia, vix ulla in caule nisi lanicolatum  
breve integerrimum scabriusculum sub ramificationibus, petiolata  
lacinis inferioribus alternis paucis oblongo vel ~~inreg~~ <sup>inreg</sup> ~~triang~~ <sup>triang</sup> ~~inreg~~  
lacis subdentata, terminali maxima oblongo <sup>ob</sup> ovata dent  
bipollicari dentata venosa, pilosa. petioli ~~pauci~~ dilatati  
amplexici caules valde pilosi proferunt margine. Caules sere  
plures <sup>ramosi</sup> ~~ramosi~~ rami subdivisi. Racemi terminales. Silicula  
rotunda ad singulum latius styli diffo elevatione margineque  
villosa scabriuscula

caule subephyllis

lect. mus. | *Thlaspidium raphani folia* Tournef Inst. v. h.  
*Biscutella raphanifolia* la Martk

peninsular B

Habitat in Penn. hies. Guss.  
Solia ex Troscum serrulata

Corrigata B. filiculis glabris, foliis lanceolatis serratis Lin. Mant.  
255. S.V. p. 592. Paigu. fl. austr. 4. V. 339

Leucopium alyssoides umbellatum montanum Col. ceph. 1. p. 203  
d. 204.

Habitat in Italia & Helvetia.

Folia radicalia lanceolata, basi attenuata, obtusiuscula  
dente uno vel altero obtuso bipolliciana <sup>in</sup> ~~marginibus~~ <sup>marginibus</sup> ~~in~~ <sup>in</sup> ~~caule~~ <sup>caule</sup> ~~linearia~~ <sup>linearia</sup> ~~caulis~~ <sup>caulis</sup> ~~superne~~ <sup>superne</sup> ~~ramis~~ <sup>ramis</sup> ~~pedunculo~~ <sup>pedunculo</sup> ~~superne~~ <sup>superne</sup>  
margine scabro ~~in~~ <sup>in</sup> ~~caule~~ <sup>caule</sup> ~~linearia~~ <sup>linearia</sup> ~~caulis~~ <sup>caulis</sup> ~~superne~~ <sup>superne</sup> ~~ramis~~ <sup>ramis</sup> ~~pedunculo~~ <sup>pedunculo</sup> ~~superne~~ <sup>superne</sup>  
filula ~~in~~ <sup>in</sup> ~~caule~~ <sup>caule</sup> ~~linearia~~ <sup>linearia</sup> ~~caulis~~ <sup>caulis</sup> ~~superne~~ <sup>superne</sup> ~~ramis~~ <sup>ramis</sup> ~~pedunculo~~ <sup>pedunculo</sup> ~~superne~~ <sup>superne</sup>  
altis duobus pollicibus, pedalis. Siccitate in facie B. auricula  
lata squamose margine membranaceo per stylos a stylo tamen  
distinctum

Folia linearia, extusum dentata, superiora linearia, integerrima  
uti caulis hirsuti. Pedunculi glabri.

auniculata B. calycibus tertio utraque guttis, filiculis in stylum  
occurrentibus.

*Thlaspidium bifidatum* villosum, flore calicari donato Bauh.  
prodr. 49. pin. 107.

*Thlaspidium alpestris* lutea angustifolia Barr. ii. 236

*Leucosium montanum*, flore pedato Col. eiphr. 2. p. 59. t. 61.

Habitat in Italia, Galloprovincia. O. petali albi nigrescunt pellucid.  
Caulis erectus ramosus, pedalis et ultra, pilosus. Folia semamplexicaula  
creta bipollicaria vel ultra ovato lanceolata remote dentato-  
subsinuata, obscure simpliciter venosa vili ramoribus caulinis  
sinuatis adpressa margine scabra. Racemi terminalis ramorum  
et axillares erecti. Filicula latiori quam longa margine membra-  
naceo per stylum ascendente squamis albidis copiosis adperso. Stylus  
filicula longior

coronopifolia B. filiculis glabris; foliis dentatis hirtis Lin. Mant. 255. ✓

S. V. p. 592.

Habitat in Italia, Hispania ☉

Radix superne divisis. Solia ad radicem aggregata fere  
Pl. coronopi dentibus utraque 23. valde & pilis copiosis valde  
hirta, <sup>vollicana</sup> & vel 3 in caule lineari lanceolata amplexicaulis  
angusta inferne dente uno vel altero parvo superora integer  
rima. Caulis plures palmatis fructi fere filiformes  
basi hirti superne glabres, ut ramofi. Filiculis omnino  
laevi

apula B. siliculis scabris, foliis lanceolatis sessilibus serratis Lin. Man.  
p. 254 L. V. p. 592.

Gondrubra alyssoides apula spicata Colum. aeghr. 1. p. 203.

t. 204. fig. 1.

Thaspi clypeatum Plin. hist. 2. p. 133.

Habitat in Italia

Tota planta caule foliisque hirta, pilis sparsis. Caulis erectus  
pedalis, ramis duobus vel tribus. Folia lanceola sessilia  
remotissime serrata, scabra. Silicula orbiculata glab. lora  
et mea, Lin. scabra

planta sub nomine B. aegyptiaci mesa ab Allionio in nullis  
aliis a B. apula differre videtur quam siliculis lobatis  
quod et fuit secundum partem sanati pili scabrusculi et  
lobatis

sempervirens B. foliis scabriusculis, foliis lanceolatis tomentosis  
Linn. Mant. 253. L.V. p. 2924

*Thlaspi bifidellatum* Tutin. anethifolium Rou:  
Mus. 267-4. 122. Barr. u. 841.

Habitat in Hispania et oriente S

Caules suffruticosi, palmari, prostrati, simpliciter, crispiter penna  
anethifolium, apice foliosi. Folia conferta, lanceolata, densa  
tomentosa (Cheranthi micari) canaliculata, dentato-serrata, mar-  
gine reflexo. Caulis floriferus inter folia enatus, simplex  
subaphyllus. Racemus <sup>terminalis</sup> terminalis, compositus, minus  
elongatus. ~~Alpula~~ <sup>Alpula</sup> magnitudine et figura B. apulo.  
Linn.

non vidi

montana *Besleria* *filicularis* glabris, foliis <sup>ob</sup> ovatis dentatis tomentosi  
Cavan. in: pag. 59. Tab. 177.

Habitat in Hispania

Caulis pedalis tomentosi etate aevi. Folis radicalibus nu-  
merosis, dentatis caulibus alternis simplicibus basi subnatis  
Calyx luteo feno foledis ortis. Corolla lutescens. Cavan.

apetala B. floribus apetalis, calycibus marginatis Walt. fl. Carol

pag 174

Habitat in Carolina

mhi ignota