

Brunnicia

emorsa. Brunnicia Godn. de fruct. 1. p. 213. t. 49. f. 2.

Rapana ovata; foliis ovatis venosis petiolatis, caule volubili
Walth. fl. car. 247.

Habitat in Bahama, in Carolina Rose.

Salic. Polygoni. Caulis volubilis, striato-angulatus, glaber. Folia
petiolata, distantia, alterna, bi- & tripolliciana, ovato-oblonga,
basi saepe subtransversa, integerrima, acuta, venosa, utrinque
glabra, simpliciter venosa. Racemi axillares et terminales; foliis
longiores: pedunculi partiali alterni, florigeri bractea brevioris;
fructiferi longiores, compressi: pedicelli 3-5 in apice pedunculi par-
tiali, uniflori, capillares, superne articulati. Bractea membra-
nacea, lanceolata, pedunculo partiali longior, eumque in volvens;
hinc pedicelli in planta florente aggregati apparent.
Calyx persistens, inferus, quinquefidus: laciniis linearibus oblongis
obtusis, margine membranaceis; fructiferis tubulosis; semi-
pollicaris, coriaceis: laciniis linearibus obtusis, inferne et la-
verte

~~25~~

gatus in pedunculo ses semi pollicari, versus articulum pedicelli filiformi
 attenuato, compresso, margine altero convexiore linea utrinque elevato
 marginato, altero subarcuato tenuiore, superne subfructu dilatato in
 receptaculo puberoso. Corolla nulla. Filamenta 10. fundoca
 levis inserta, eoque longitudines. Germen unicum, ovatum, attenua
 tum. Stili tres, longitudine calycis, divergentes. Stigmata bifida, diver
 gentia. Capsula e calyce ante inclusa aique brevior, hinc primo
 introitu fructus. calyce corollata videtur, crustacea, ovalis,
 trigona, monosper. Semen ovatum, acutum, sulcatum, fusco
 purpureum.

Carludovici
 Dr. folia ovata venosa pectinata, caule reticul. Max
 fr. carol. pag. 247
~~Floribus in Canadum~~