

Bradloa

umbellata B.

glochidium Forster. hue a mon

rami florum *Glochidion* Torft.

Habitat herb. Mus.

Rami teretes, glabri. Folia petiolata, oblonga, integerrima, glabra. Pedunculi  
breves, plures, uniflori.

~~foliis  
myrsinoides~~ ~~Phyllanthas~~ ~~foliis~~ ~~pinnatis~~ ~~oblongis~~ ~~pedunculis~~ ~~unifloris~~ ~~solanis~~, caule arboreo  
~~ramnoides~~ ~~fructu~~ ~~capitellari~~ ~~ovato-~~  
foliis ~~oblongis~~ ~~ovalebus~~ ~~obtusis~~ ~~glabris~~ ~~alternis~~, ad foliorum ~~orium~~ ~~frunis~~ ~~etiquis~~ ~~exappicatur~~  
~~bifurcatis~~ ~~ramulis~~ ~~pluth.~~ ~~phyt.~~ ~~t. 122. fig 4~~  
pedunculis ~~terminalibus~~ ~~articulatis~~ ~~t. 4~~  
tertiis floribus ~~Rhamnus ceylonicus~~ folio subrotundato glabro, caulis hispida, frunis  
exgau, ad foliorum orium exappicatis, glauculo pedunculo, basica

~~Rhamnoides Roten.~~ ~~negra~~ ~~Burm. fl. Ceyl. 190. t. 88.~~ forte modo quod in Burmanni  
obs. 5. pag. tanai hispida in nostra glabra est igitur Rhamnus utriusque

~~Habitat in India orientali.~~

~~Burm. flor. India eadema planta~~

Arbor ramis terribus, glabris cortice purpurascente, superne apice  
compresso-piceo Ramuli saxe gemino

Folia alterna, saxe gemina palmata foliolis plerunque II. petiolis

utrumque lata ~~altissimis~~ de strobilis, ungue ulana oblonga vel subovata  
(strobilo subulata) integririma obtusaveneris venosa, pubescens  
subrugosa Petiolis linearis. Squama nasuta ad petiolis basi rugido ovata  
ovata parvo subsparsa inter petiolas squama vel squamulae et squamulae

ut nique. ~~Hipulò binò f. subulato ad bofia foliòtonum subulato, petiolato~~

~~binario~~

<sup>3 nigris binos magis</sup>  
Pedunculi ex axillis folio <sup>non</sup> longiori solitarii un. flor. breviores  
<sup>in foras locules de cedentia</sup>  
superne foliis <sup>longiora</sup> <sup>longiora</sup> <sup>longiora</sup> magnitudine perennis  
Cappula globosa utr. depresso glabra & lineata, Cerasa loculari. Loculamen-  
tis obtus. convexis latibus con praesiso margine intenore acutis, disper-  
mi

Rhamnus reylanicus, folio subrotundo glabro etc. Burm. fl. Reyl: pag.

190 Tab. 88 plantam nostram optime representat, cum vero rasoni his  
justi in Burm: quo in mea sit tota planta glaberrima forte diversa  
est, si varietas una cum Rhamnus vitis idem Burm: fl: indeq. ap.

<sup>ad plas specie spudem et hanc nostra non diversa est</sup>  
an a genere, Phyllanthi diversa, cum stamina 6. stigmata et Cappula  
6 locularis loculis dispermis, habet foliis vero modo floendi et tota  
habitac phyllanthi similes, si a Phyllanthi separare habet licet generis  
Debet conjungi cum Bacynia sa disticha Tostani

6 loculari monospermis

pubescens ♀. foliis ovafo oblongis utique attenuatis pubescentibus vellosulis  
pedunculis brevissimi ramis pubescens. foliis  
glaucis ~~oblongis~~ <sup>oblongis</sup> ramos pubescens. foliis

~~racemosos~~ L.

Habitat in Otaheite

non vidi

glauca Bradlea folii ovati oblongi acuto-glaucis

Habitat in India orientali.

edulis

~~S. foliis angulatis basi cordatis lobatis fructu oblongo echenato~~

~~lascu. Amer. Tab. 163~~

non vidi

ndida B. foliis ovato-lanceolatis glaberrimiis basi rotundatis; pedunculis femineis  
foliatis elongatis ramulis glabratissimis sol.  
*Breynia* <sup>Dufouria</sup> ~~andachnoides~~ Forst.  
Habitat in Otaheite

non vidi

ramiglomerum Glochidion Dost. Schenck gen. 57.

Habitat in Societatis Insulae et Novo Hebrides 5

non vidi

*umbellata*  
~~Umbellata~~ foliis oblongo - obvato ovatis, pedunculis calycibusque glabris  
fructibus setulobatis

Habitat in India orientali  
Arbor <sup>alterna</sup> Ramis <sup>inference</sup> <sup>cuneo glabri</sup> <sup>ramum acum.</sup>  
foliis petiolatis alterna bipinnatis vel parim ultror. ellipticis a optusa  
brevi int. egerima <sup>non pinnatis acutis</sup> glaberrima supra pallide verde atroisque glabri  
ma supra obsoleta rufa pinnatis levigata subtus puberula  
terminata Petiolis brevisimis. Stipulae breves utroque ad basi petoli condita  
ad nata apice petala rigida <sup>minuti</sup>  
Pedunculi oscillari masculi et foecunnei astillares. Pedunculi  
breves 3-5 brevis filiformes uniflori, unicus vel quis longior  
Capitula papere duplo major glabra & lobata.

ovalifolia Bo. foliis ellipticis obtusisimis, pedunculis calycibusque glabris  
fructibus duodecim foliatis

# Brachysa pinnia Gmelin de sem. et. gr. 2. pag. 127

Habitat in Thacia orientali f. capio purpureo-picea  
rami glabri inferne verete superne ang. compusi. folia  
petiolata alterna glabra venosa basi pallide vnde  
triplicana, obtusissima ut emarginata basi panum <sup>sub</sup>acute  
pedunculi petiolis semiuinculari glaber. pedunculi 4-6.  
ut ~~pro~~ pedunculae petiolae uniflori glabri. lacinia calycis ovalis  
marginem panum membranacee. fructus piso duplo major.

# Plathn. tab. 368. fig. 1

ramiflorum 13.

Glochidion ramiflorum Forst.-gen. nov. sp. n.

Habita in forestis insulae et nova Hebrides

oculus

stellata 13. foliis ovatis, corymbis axillaris brevissimi calycibus vellosis  
Habitat in India orientali.

Primum angulata triquetristylata: inferne glabre superne velis brevi-  
bus rionibus disperse. Tolia breviter petiolata alterna remo-  
ta, integerrima, utrinque glabra, simpliciter venosa, glauca  
vnde et trigollicana. Petiolis subhirsutis; pedunculi axillares et  
transversi petioli petolo pubescentes longiores uti calycis et pedunculi  
subhirsuti. Calyx angulatus setiferus; lacinias lineanibus. Corol.  
o. stylis calycis duplo longior apice indentatur

*Phyllanthus stellatus*

petr. obs. fasc. 5

Bractea

- ♂ Calyx perianthium monophyllum per partitum calycinis linearibus, obtusis  
Corolla <sup>vesticula</sup> 0  
Stamina filamenta sparsiparia. Anthers tres oblongo erecta. latere exteriori filamenti  
ad apicem aperta terminata mucrone nigrante filamento transversale  
♀ Calyx Perianthium ut in mare perffuso.

Corolla 0

- Stamina 0  
Style 1. Stigma brevissimum Stigma globosum in partitum <sup>per septens</sup> <sup>truncata</sup>  
Pappula festacea orbicularis utrinque Depressa extus fuscata pro fungo  
loculari, fixa locularis. Semina foliata trigona extus convexus  
intus latenter compressi inter acutis

Fructus repeat officia zeylanicae Gartn: pag. tab. differt vero frui  
Visicatio et relegate ab Aquileia potius enim Phyllanthi species est  
si non genus distinctum.