

Pacalia

glaucæ *Cacalia foliis lanceolatis* ^{sessilibus} integerrimis glabris, corymbis axilla-
ribus terminalibusque simplicibus

Habitat in Carolina Bosc. Ex herbo Parisino

Caulis herboeus, ^{actus} erectus, simplex, teres, striatus, glaber, uti tota
planta. Folia bipollicaria, remota, plerumque alterna, inter-
dum opposita, raro terna, subrotunda, subtrinesaria, vix venosa.

Pedunculi ~~ex axillis supra~~ ^{inferiores} superne, axillares caule altiores, ter gen-
tem breviores, plerumque nudi, interdum foliolo uno alterove ad-
spersi: pedicelli filiformes, superiores semper minores, & quamvis
si vel nudi. Calyx simplex, pentaphyllus, foliolis linearibus. Corolla
lo quinque. Semina obovata, ~~la~~ vix

Sonchus

Lactuca orientalis, dentis Leonis folio, flore luteo C. P. Tournef. Cor. pag.

Habitat in Oriente herb. Martii

Caulis pedalis, striatus, glaber, uti tota planta. Folia inferiora sublyrata, bipol-
licaria, superiora amplexicaulia, cordato-oblonga vel cordata lanceolata,
acuta, omnia glabra, dentata, tenerissima. Pedunculi superiores, rami
pedicelli basi squama instructi. Calycis foliola lanceolata. Pappus sessilis
dentatus. Semina levia

glabrata

Caecalia foliis ^{glabris} radicalibus subcordato - ellipticis inaequaliter crenatis,
superioribus
caulinis cordato-lanceolatis extrorsum dentatis.

Habitat in Peru. Ex herbario Fusseri

Caulis herbaceus, crassitie penna cygnea, erectus, superne divi-pus, stri-
tus, subangulatus; glaber, uti tota planta. Folia radicalia petiolata,
epithamosa, basi subcordata, altero latere saepe interdum breviora, inae-
qualiter acute crenata, obtusa, subtus venosa: caulina subramis flori-
feris attenuata, acuta: suprema integerrima. Pedunculi albescenti,
ramosi; pedicelli uniflori, basi foliolo suffulti. Calyx polyphyllus,
basi squamis nonnullis parvis calyculatus. Foliola linearia

Herba Pentstemon

~~Carduus orientalis cyanifolius Tournaf. Cor~~

~~Habitat in oriente habi. Mus. cono. cers~~

~~Caules herbaceus ramosi, villosi. Folia inferiora petiolata, superiora sessile, frequentius, tripol~~

~~licana vel minor, basi versus attenuata, acuta, villosa, subanastomata, siabra, in cymis~~

~~avenis, cernis. St. terminalis subsessile, magnitudine c. siculo squama ovata, termi~~
~~nata pini brevi, basi utraque spinula 1-2.~~

Lyustriana

Cacalia foliis lanceolatis acuminatis ~~glabris~~ integerrimis glabris,
pedunculis unifloris.

Habitat in Peru. Ex herbario Fursii: 3

Caulis fruticosus: Rami teretes, ~~glabris~~ conferti, saepe terni vel plures ver-
ticillati, tuberculato-cicatricati, glabri. Folia subpetiolata, fre-
quentia, hemipollicaria, basin versus attenuata, subcoriacea, gla-
berrima, avenia, enervia. Pedunculi ex supremis axillis, alterni,
approximati, superne squamis duobus vel tribus subulatis adspers-
i. Calyx polyphus. Loculi plures. Pappus simplex
an huius generis

~~Verniifolium ^{Pentodon} foliis linearibus dentatis scapo superne calyceque subtomen-
toso.~~

~~Habitat ad fretum magellanicum Thoun~~

~~caespitosa
caulis~~

~~Folia angustissima dentibus remotis acutis digitalia Scapus foliis
dimidio longior medio foliolo subulato superne versus florem incurva
calycis foliola subulata. Pappus pileosus~~

Scapigera *Cacalia* foliis lyratis, pedunculis scapiformibus elongatis filiformibus.

Habitat in insula Francia

Caules plures, pollicares. Folia in petiolum pollicaria, in petiolum attenuata, profunde dentata lyratis, lacinia terminali ^{reliquis quadruplo major} oblonga integerrima. ope lentis rare villosa. Pedunculi semipedales, teretes, glabri, superne

semificationibus semel vel bis divisi[†]. Flores rari. Calyx hexaphyllus, basi folioli non nullis calyculatus, sex vel septemflorus.
v. lineari suffult

Scanzera
Bubocaulis

Cacalia foliis lyratis, caulis brevis siccus, pedunculis elongatis

longissimis filiformibus

Habitat in Tava

Raulis semi pollicaris, decumbens, divisus in ramis duobus vel tribus, vix polli-
cibus

Prof. Scanzera
Cacalia

cuneifolia *Cacalia* foliis ^{sub} sessilibus cuneatis integerrimis, pedunculis axilla-
ribus terminalibusque trifloris.

Habitat in India Orientali. Ex herbario Fuscoi.

Caules herbaceus, erectus, six pedalis, ramosus, striatus. Folia infima
subpetiolata, reliqua sessilia, alterna, pollicaria vel parum ultra,
mucronata, crassiuscula, glabra. Pedunculi elongati; ^{cauli} pedicelli terni,
uniflori. Bractea subulata ad basin pedicellorum. Calyx atrophyllus, semi-
pollicaris, glaber, multiflorus. Styli longi, bifidi.

~~Cuneiformis Bidens~~

Habitat in Penn. herb. tussoi

Caulis herbaceus, pilosus, fruticulosus, ramosus, rami diffusi, dichotomi. Folia opposita, semipollicaria, cuneata, petiolata, trifida: lacina serrato-incisa, supraeque margine raris pilosa. Pedunculi a dichotomia et apice, foliiformes, solitarii, uniflori, semipollicares. Calyx simplex: foliola linearia ciliata. Antherae duo divaricatae

buxifolia Calycis: foliis lanceolato-^{subtriangularibus} integerrimis acuminatis,
glabris, floribus axillaribus, solitariis sessilibus.

Pongia buxifolia; glabra, ramis strictis, foliis oblongo-ovatis integerrimis, floribus lateralibus sessilibus La Mart. Dict. bot. 2. p. 92.

Habitat in Peru ex herbario Gussonei &

Rami frequentes, sparsi, post casum foliorum tuberculati; ramuli angulati, glabri. Folia sessilia, frequentia, semipollicaria, subrotunda, aserna, enervia. Calyx simplex, basi squamis paucis calyculatus: foliola linearia. Corollula omnes hermaphrodite

~~Caecilia~~

~~Habitat herb. fr.~~

~~Caules erectus, basi lignosus, ramifus: rami frequentes. Flores Vermenales-~~

~~Solitarii~~

Lunifolia *Cacalia lanato-tomentosa*, foliis linearibus integerrimis, caulis
unifloris pedunculis terminalis, unifloris

Habitat ad fretum magellanicum Commerçon Thoun

Caules suffruticosi, simplices, ascendentes, pa spithamei, uti
folia et pedunculi lanato-tomentosi. Folia sessilia, 8 pass a,
frequentior^{bina}, vix pollicaria, angusta, Pedunculus quia sub
calyce squamulis paucis, subulatis. Calyx multiflorus, polyphylus
basi tomentosus, superne glaber: foliola linear-lanceolata. Corolla
flava.

pauciflora Caecalia herbarica, foliis lanceolatis integris amplexicaulis;
pedunculis caulis alabibus unifloris

Habitat in Ceylona.

Caulis adscendens, inferne flexuosus basi cyropus glaber

Folia inferiora plura, ^{glabris} approximata, superiora ^{angustiora} duo vel tria, remota
semiamplexicaulia, pollicaria, subindegerrima, summa tenera
brevia.

Pedunculi superiores in caule, alterni, solitarii remoti. 3-4.

capillares, bipollicares, uniflori: folium lineare superne
versus florem ecalyculatus multiflorus

Calyx pentaphyllus: folioli lanceolatis glabris

valde affinis Caec. fontefot. an modo varietas

Caulis rigidus, folia haec brevifolia errata in Cupat. no 6

~~altorri folium~~ ^{baichangides} Eupatorium foliis ovato lanceolatis ^{acutis} serratis trinerviis, floribus panicu-
latis.

Habitat in Brasilia 5

~~Frustrum~~ caula ramis nigris striatis, glabris sicut reliqua planta. Folia
^{ret. w. l. d. u. p. n. o. s. s. a. l. t. e. r. n. e.} ovato lanceolata, serrata, trinervia, utrinque acuta, longi pollicaria
Petioli semipollicaria. Panicula terminalis virgata, foliis petalis
ad basi pedunculorum gemmalium et pedicellorum. Calyx imbricatus, hemi-
sphericus multiflorus, minutus, foliis oblongis acutis longitudine flosculorum.
Pappus plumosus
Habitat in Brasilia ad flosculi omnes hermaphrodite

Potui C.

Habitat in Europa C.

ad fontem folium prope me accidit: differt calycibus ovatis, spina inter
media duplo longiore: foliis radicalibus lyratis, minus dentatis, & lobis
luteis Spreng. Bot. Hall. 23.

petufter Cacalia

La Mancha

Habitat h herb. fus.

Caulis herbaceus, sulcatus, erectus, pubescens. Folia petiolata, pedem fere lata, squa-
nosa, venosa, subtus pallidiora, villosula, basi subreniformia, acuminata. Radicum
li supernae, alterni. Corymbi compositi. Calyx simplex, pentaphyllus: foliolo apice
subfusca. Corollula squa-

Cacalia?

Facolia africana lutea, Tanxaci folio, crasso, avenae, enerve

Habitat in Africa herb. Toros n. n. f.

Glabra. Caulis herbaceus, striatus. Folia inferiora petiolata, reniformia
subrotunda, semipollicaria, obtusa et grosse dentata; petiolus pollicaris.
Inferiora ^{minima} subtruncata, laciniis extrema subrotunda, obtuse dentata, lacini
les distantes, oblongo, obtuso, superiora semiamplexicaulia, cordata-lan-
ceolata, pollicaria, dentata, superiora integerrima. Pedunculi superiora
subgemini, elongati, uniflori, squamis nonnullis remotis adpressi.
Flos magnitudine Ciner. pariflorus. Calycis paleola lanceolata. Radix
nullus

-Ineffructuosa C. foliis lineari-integerrimis, pedunculis elonga-
tis unifloris

Habitat in Brasilia.

Vente

Pro ni sparsa, uti tota planta glabri. Folia sparsa frequen-
ter angusta acuta basi attenuata. Peduncululi serminali
aphylli. Foliola Calycis quinque, lanceolata

Linaria Bacata foliis linearibus sparsi integerrimis ad limum porofo
floribus solitariis terminalibus Caron. ii. vol. 3. pag 29. tab. 257.
Habitat in nova Hispania 5

rotundifolia Cactin foliis orbiculato-ovatis repando crenatis, flori-
bus corymbosis caule ramoso Welden. phyt. pag 11. tab. 6. fig. 1.
Senecio indicus atroplicis foliis glabra Monf. hist. 3. p. 106
f. 7. 7. 17. fig. 7.

Habitat in India orientali

Caulis herbaceus erectus, teres glaber. Folia alterna
venosa, utrinque scabra inferora repando-crenata
superiora obsolete crenata. Calyx calyculatus. Pappus
vilosus scaber.

mihi ignota

tenuifolia Achillea foliis filiformibus pinnatis pinnis
cordatis integerrimis, caulebus simplicibus.

Habitat

Caules plures

1 papillaris *Caecalia* caule fruticoso obvallato pinnis petiolariibus truncatis L.S. pl. 1160.
S.V. 733.

Ulcinia caule carnosio, petiolis truncatis obvallato Hort. Cliff. 395.

Caecalienthemum caudice papillari Dill. Clth. 63. t. 55. fig. 63.

Habitat in Ethiopia

non vidi

9 odora *Racalia fruticosa* folia lanceolata plana, floribus umbellatis
C. odora foliis lanceolatis; caule lineis albis striato *Dosk. Desf. p. 146*

Habitat in montibus arabicis felix Yemen. 5

~~Caulis erectus ramosus, foliis deciduis ciliatis, apice folii: ciliis
obtusis foliorum delapsorum albis, protuberantibus; ab hac ciliatura purpur
et deorsum lineis tres nitido elevato decurrunt. Toti folia in apice ramo
rum juniorum sparsa lanceolata, crassa sessilia glabra. Pedunculi termonia
les umbellati 5-7 erecti squamis albis lanceolatis ad basin. Foliole 2 lanceol
atae ad basin calycis. Calyx cylindricus post anthesin glaber Flores 9-
13. *Dosk.*~~

3 Anteuophorium *Palalia* caule fruticoso, foliis ovato oblongis planis, petiolis basi lineari
triplici deductis L. Sp. pl. 1160. t. v. 1793.

Anteuophorium Tod. pempt. Lob. Bauh. p. 307. Monf. h. 3. p. 345. f. 7. t. 37.
fig. 10. Dill. Elth. 63. t. 55. f. 2. 3.

Palinia foliis carnosis planis ovato-oblongis Hort. Cliff. 395.
Habitat in Ethiopia h.

non vid.

4 *Cuneifolia* *Catalia*. Caule fruticoso, foliis cuneiformibus carnosissimis Mant. 110 L. pag. 733.
Habitat ad Cap. b. p. h.
Caulis carnosus, minus C. Ficoides et antecedentibus. Folia parva, avenia, absque
costa, supra plana, subtus non nihil convexa carnosissima. Cum folia non opposita
nequit esse cotyledon. Fructificatio ignota.

non vidi

45 odora Cacalia fruticosa, foliis lanceolatis planis, floribus umbellatis
Cacalia odora, foliis lanceolatis, caule lineis albis fruticosa Forsk
Defr: pag. 146

Caulis erectus, ramosus, cicatricatus: cicatricibus protuberantibus, lineis
tribus ^{albis} elevatis et singulis cicatrice lineis tribus, elevatis, nitidis
sursum et deorsum excurrentibus.

Folia in apice ramorum junctorum ^{reflexa}, parva, lanceolata, crassa,
glabra. ^{ex apice ramulorum}

Pedunculi terminalis, 6. 7., umbellati, erecti, ^{uniflori} basi squamis
nonnullis lanceolatis adpressis.

Calyx cylindricus, pentagynellus, glaber, basi foliis duobus linearibus

Corollula 9 numerari Forsk. 13.

10 *Semperviva* *Cacalia* *carnosa* caule dicto foliis lancicollatis obtusis

C. semperviva foliis carnosis, sessilibus, lanceolatis, subnae confertis ^{100 M}

Defer. p. 146

Habitat in montibus Arab. foliis 2

Caulis vix pedalis. ^{lancicollata} Folia basi ciliatis. Folia sessilia carnosa conferta

glabra, & obtusa bi nitida plana subtus canna longit. carinata bipolli

caria. Peduncululi terminales plerumque bini erecti albo punctati

versus apicem squamati, uniflori, spithamei Calyx polyphyllus multi-

florus. floribus purpureis

4 *Valeina Cacalia* caule fruticoso composito, foliis lanceolatis planis, petiolorum de
catis. *Tabus obsoletis* L. Sp. pl. p. 1168. L. V. 733.

Autor Lavandulo folio Clus. exot. L. I. c. 5
Frutex Indis orientalis folio nerei glauco Bell. Leth. 61. Tab. 54. p. 2.

nec *Cacalia*, nec *Cacalia* frum, an *Tithymaloides* nerei folio frutescens *Valeina* monog.

Valeina foliis lanceolatis planis, caule lavi ventricoso Hort. Cliff. 395.
Habitat in Canariis, forte etiam in India.

Caule foliis deciduis ciliatis, ciliatis rotundis. Folia
nervi lanceolata basi attenuata acuta in tegmina
longum flores in se subumbellati.

aggregata Ononis

28 sempervivens ^{va} Calathea carnosa, caule erecto, foliis lanceolatis obtusis, pedunculis
^{sub geminis terminalibus unisporis}
Calathea sempervivens foliis carnosis, sessilibus, lanceolatis confertis
Forst. descr. pag. 416

Caulis ^{virgata} pedalis, inferne foliosus.

Folia ^{bipollicaria} sessilia, carnosa, lanceolata, conferta, glabra, obtusis, nitida,
plana, subtus carinata.

[Pedunculi terminalis, plerumque binis gemina, erectis, albo
punctati, versus apicem squamis adpressis. Smithsoni,

Calyx polyphyllus, multiflorus.

Corollula purpurea

[Lit. Ita. pag. 93.

5-6.

2 172 10

24 ^{Torsholci} ^{glabra} ^{oblongi} ^{dentata} *Caralia herbacea*, foliis ^{oblongis} ^{dentatis} ^{lyratis} petiolatis, Panícula ^{multiflora}

Caralia foenicifolia Torst. Cat. pl. Arab. n. 485

Habitat in Arabia felici

~~Herbae Caulis ^{pubescenti} striatus ^{superne} divisis. Folia ^{inferiora} ^{scabra} ^{oblonga}, ^{angulato} ^{dentata}, ^{nonnulla} ^{lyrata}, ^{laevius} ^{glaberrima} ^{avenia}, ^{acuta}, ^{interdum} ^{basi} ^{sinuata} ^{lobulo} ^{utrinque} ^{opposito} ^{lan} ^{ciolato} ^{sub} ^{dentato}, ^{sequi} ^{pollicaria}, ^{petiolis} ^{folio} ^{dimid} ^{triplo} ^{breuior} ^{amplex} ^{icaulis} ^{superne} ^{den} ^{delatatus}. ^{Rami} ^{Rami} ^{florif} ^{superne} ^{divisi} ^{foliis} ^{hem} ^{lan} ^{ceis} ^{latum} ^{sub} ^{divisus} ^{ubi} ^{caulis} ^{superne} ^{divisi} ^{tor} ^{pedunculis} ^{pedunculis} ^{dividitur}, ^{stipitem} ^{sub} ^{pedun} ^{culis} ^{pedi} ^{cellisque}. ^{Pedicelli} ^{capitales} ^{foliolo} ^{stipitem} ^{respu} ^{api} ^{cem} ^{calyx} ^{calyx} ^{ulatus} ^{foliolo} ⁵ ^{lineari} ^{lan} ^{ciolato}~~

Paniculo pluro
multiflori fasti
grati

6 *ficoides* *Cacalia*. caule fruticoso foliis compressis carnosis L. Sp. pl. 1160. S. V. 733
Senecio africanus *arborescens*, *ficoidis* foliis et flori. Commel. rar. 40. t. 40. Bradl.
Juss. 5. p. 11. t. 49.
Helinia foliis carnosis lanceolatis compressis, caule tereti Hort. Cliff. 395
Habitat in Ethiopia

non vid.

6-7 *carnosa* C. caule fruticoso, foliis teretiūsculis carnosis incurvis, pedunculis terminatis
libris unifloris nudis Hort. kew. 3. p. 156
Habitat in Cap. b. p. 5

non ad.

7 repens *Cacalia* caule fruticoso foliis depressis carnosis Mant. 110. S. V. p. 733.

Habitat ad C. bona spei

Planta tota statura simillima *C. bicoides*, etiam rore glauco obducta, sed folia supra
depresso concava, nec lateribus compressa. *C. Mant. l. c. Emendatio ignota. Radice*
huic repens non illa

folia angusta sessilia acuta alterna. Pedunculi terminales
interdum gemini pauciflori folia calycis lineari lanceolata

Maciniata *Caecalia* caule fruticoso, foliis carnosissimis pinnatifidis laciniatis L. V. p. 753.

— Jacq. Michx. vol. 3. Gen. rar.

Habitat.

Habitus *Cotyledonis laciniato*.

non vid.

o Suffruticosa Cacalia caule suffruticoso ramoso, foliis linearibus planis sparsis Mant. 109*
L.V. 733.

Cacalia foliis linearibus integerrimis, caulibus unifloris. Ard. spe. 2. p. 39 t. 19
Habitat in Brasilia &.

Caulis suffruticosus, palmatis, vane ramosus scitiformis. Folia sparsa, linearia, integra
gemina, minime carnosus. Pedunculi terniiales, petioli, uni flori, erecti, longior
vane caulis. Calyx simplicissimus pentaphyllus: foliis ovatis obtusis, erectis, loben
bus occultatis membranaceis. Flosculi multi, purpurei. Pappus pileosus. Flos
omnino Paraphylli, sed herba diversa Lin. Mant. L.C.

~~no. 10.~~

910

reticulata
amplexicaulis

Cacalia fruticosa, glabra, foliis cordatis subrotundis amplexi-
caulis denticulatis, floribus paniculatis

Habitat in Insula Bobonis Commerson Dedit Dr Thourin

Caulis fruticosus, ramosus. Rami e nervis foliorum decimen-
tibus angulati, glabri; articulis sub foliis inersatis

Folia fere Bobonis inersata, alterna, ^{obliqua} (cordata), subrotundata,
glabra, venoso-reticulata, denticulata: dentibus, uti
margo ^{inferius} foliorum, ^{reflexo} inersatis. Nervi basi ~~colorati~~ sanguinei

Panicula terminalis, 6 pollicari; pedunculis pedicellis
que glabris: folium parvum ad basin pedicellorum

Calyx simplex, glaber: folioli 12. lineari-lanceolatis, squa-
libus

Pappus simplex, longitudine calycis.

20 pauciflora Cacalia Caule herbaceo, foliis lanceolatis indivisi amplexicaulibus, Pedunculis
lateralibus axillaribusque unifloris
Habitat in Ceylona.

Caulis ascendens inferne flexuosus. ^{in fenore approximato} folia plurima in inferiori parte caulis superioris
^{remota} ~~laxa~~ f. triplex caulica, lanceolata, subintegerrima, pollicaria. Summa linea
non brevis. Pedunculi a superiora parte caulis alterni, solitarii remoti 3-4.
capillares bipollicares uniflori versus florem foliolo petales. Calyx pentaphyllus
laciniis lanceolatis, e calyce alatus multiflorus

obovatifolia *Racalia* fruticosa glabra, foliis petiolatis ovatis triplinerviis, obtusis integerrimis
glaberrimis, thyso terminali, calyce tetraphyllo glabro L. Suppl. p. 351 L. V. p. 733.
Habitat in Mexico &c.

Frutex glaberrimus Viburnum Tinum s. Laurum referens. Folia opposita bre-
ve petiolata, ovata, integerrima, obtusa, pollicaria nervosa: nervis supra
basin excurrentibus; subtus viridia subtus splendens, consistentia
sempervirentium. Panicula terminalis thysoidea, conica. Calyx tetra-
phyllus quatuor floris. Foliola aequalia, oblonga, obtusa, duo et duo opposita
Corolla limbus tubo major, quadri-dentatus. Pappi Congiura. Lin. Suppl. L. V.

Differentia sesquipedalis

non vid.

13 *Cordifolia* *Cacalia* *lutescens* *hirsuta* foliis petiolatis cordato-ovatis nervosis acutis scabris,
Calyce tetraphyllo quadrisloso pubescente L. Suppl. p. 351. S. V. p. 733.
Habitat in America meridionali.

Folia opposita, longe petiolata, late ovata, basi cordata, apice acuminata,
serrata hispida scabra, trinervia et venosa. Petioli tomentosi. Panícula
terminalis, foliis floralibus vestita. Calycis glaberrimi, nudi tetraphylli
foliolo aequalia opposita, lanceolata, glabra, erecta, flores quatuor aequa-
les hermaphroditi. Corolla limbo quadripartito. Stili duo staminibus longiores.
Pappus simplicissimus, calyce longior. Variat foliis integris et serratis L. Ruppel

L.
Q. *laevifolia* et *Cordifolia* flore exactissime conveniunt, et numero constanti
calycis et flosculorum adeo singulares sunt, ut facile proprium genus consti-
tuere possint. L.

non vid.

Astilpeuadea Cacalia fruticosa tomentosa, foliis petiolatis ovatis lanceolatis integerrimis
 supra glaberrimis subtus tomentosis, margini revolutis, paniculis terminalibus
 L. suppl. p. 752. S. V. p. 733.

Habitat in America meridionali

Caules teretes, tomento molli cana, recti. Folia petiolata, opposita, patentia,
 basi ovata et sensim attenuata, ut fere media inter ovatum et lanceolatum.
 Subtus valde tomentosa, supra glaberrima, integerrima, consistentia folii laevi.
 Petioli teretiusculi, tomentosi. Paniculae terminales parvae, congestae. Calyx
 calyculatus squamis majoribus, paucifloris. Herba faciem habet *Astilpeuadis*
 Lin. Suppl. l. c.

16 *appendiculata* *Caecalia fruticosa tomentosa*, foliis petiolatis ovato lanceolatis cordatis
ovatis acutis angulatis subtus tomentosis, petiolis appendiculatis foliopsis
L. Suppl. p. 352. P. V. 733

Habita in Teneniffa locis aquosis

Tribes caule angulato, tomentoso, cano. Folia altiora, longe petiolata, cordata,
subangulata, ovata, acuta, ad petiolum non multum emarginatum, nervosa,
venosa, supra glaberrima viridia, subtus tomentosa-albida. Mox infra folium
in petiolo foliola appendiculata, sive trigone opposita, minima, petiolata, ovata
integerrima, a folio remotiora majora et sensim versus folium minor. Pani-
cula terminalis. Pedicelli bracteati. Flores lutei L. Suppl.
varietas datur sic: hinc caule glaberrimo, foliis cordato oblongis, subdentatis
nec angulatis

non vid.

17 tomentosa Cacalia fruticosa tomentosa, foliis lanceolatis dentatis subtus tomentosis sesquipedibus
L. Suppl. p. 353. LV. pag. 733
Habitat ad Cap. bono spec. h

non vid.

10 *Lanceolata* *Cacalia* *pubescens* ^{glab.} *foliis* *petiolatis* *lanceolatis* *subdentatis* *floribus* *subpaniculatis*

90
hab in amer. merid. *Asiolo*
Rami teretes striati glabri

Folia alterna *petiolata* *lanceolata* *inferiora* *dentibus* *uno* *alterove*
suprema *integerrima*, *glaberrima*, *multinota* *bipollicaria*
pediculis *brevissimis* *linearis*. s. b.

Flores *terminales* *suppaniculatae* *pedicellis* *brevibus* *foliis* *binis* *vel*
3. *linearis* *lanceolatis* *instructis*

Calycis *foliola* *glaberrima* *o* *linearia* *obtus.*

Habitus *prophylli* *et* *pubescens*, *floribus* *non* *solitatis*, *foliis* *lanceolatis*
si *distinguitur*

non eadem cum *ruderali* *Trigu*

36 articulata *Cacalia carnosa* caule decumbente articulato, foliis inferioribus hastatis
superioribus lyratis. Suppl. p. 354. L. V. p. 794.
Habitat ad. B. bono spei

non sicut

19 porophyllum Cacalia foliis ellipticis subcrenatis caule herbaceo et foliis oblongis subcrenatis
Senecio indicus, atriplicis folio glabro Monf. h. 3. p. 106. f. 7. t. 17. fig. 7.

Tagetes foliis integris perforatis Plum. spec. 10.

Chrysanthemum americanum frutescens, balsamico foliis nigris maculis
punctatis Plur. alba. 100. t. 16. fig. 1.

Porophyllum foliis ellipticis Hort. Cliff. 494 *

Cacalia caule herbaceo indiviso, foliis ellipticis subcrenatis L. l. pl. p. 1169
l. v. p. 733.

Totus Globus caulis ~~fr~~ ramosus striatus. Folia opposita alternaque petiolo
ta ^{oblonga} ~~ovata~~ basi ~~truncata~~ angustiora ~~fr~~ apice ~~fr~~ rotundata subre
tusis, mucronata, ~~ovata~~ ^{longitudinibus} ~~ovata~~ ^{sepe folijs} ~~crenatis utrinque~~
binis vel tribus remotis, pollucaria. Petiolus filiformis. Peduncululi termi
nales cum ramus excrevit laterales sunt binii vel ~~fr~~ solitarii, uni
flori ^{ternati} ~~superne~~ ^{et alij} ~~enervati~~ ^{ultrius} ~~bi~~ ^{multiflori} ~~pollucariae~~. Calyx ~~pentaphyllus~~ ^{multiflorus} ~~foliolis~~ 5 lan
ceolato linearibus. semina linearia angusta nigra

14-15 *ruderalis* C. caule herbasco ramoso, foliis lanceolatis integris glaucis Sw. prod. p. 110
Nleeria ruderalis Gay. Amer. 215. t. 127
Habitat in Jamaica C

villoso inferionibus petiolatis
vulgo Cacalia foliis lyratis, superionibus dentatis amplexicaulis dentatis
superioribus lanceolatis dentato incispi

Cacalia villosa Jacq. coll. 2. p. 347.

Habitat ad Cap. b. p. 2

Tota villosa, asperiuscula. Caules plures erecti, pedales, striati. Folia
aeterna, lyrata, squarropula, craspula: infima longa petiolata, spollioni
sequenti dentata, amplexicaulis, obtusa, venosa; summa acuta, sub
lanceolata, inciso-serrata. Pedunculi terminali, ramosi, compositi erecti
florescenti foliola calycis circiter 12 lineari acuta, erecta villosa. Flores cir
citer 35 albedo limbo purpureo. Semina oblensa Jacq.

non vid.

15-16. Sagittata Cacalia herbacea, ^{# amplexicaulis} foliis lanceolatis sagittatis denticulatis.
glabris

Heremium javanicum: foliis lanceolatis amplexicaulis denticulatis, pedunculis unifloris, calycibus glabris Burm.
fl. Ind. pag. 174. Tab. 57. fig. 1.

Habitat in Java.

Caulis herbaceus, teres, striatus, glaber.

Folia amplexicaulia, sesquipolliciana, lanceolata, inaequaliter remote denticulata, glabra, subarenia, acuta.

Pedunculus caulem terminat, bis. tri. pollicans, ^{sub} uniflorus.

sub flore parum crassior

Calyx simplex, glaberrimus, dodecaphyllus: folioli lanceolata.

Corollula longitudine calycis, purpurea.

(Affinis C. sonchifolia, diversa foliis et floris ^{ver} triplo majoribus
Figura Burm & Burm: edita cum aerea planta in omnibus
convenit nisi quod pedunculi quadriflori, quamvis differentia
unifloros habet. ~~Heremium javanicum, Burm. Ind. pag. 174. Tab. 57. fig. 1.~~
~~non est a veris~~

Planta Burmanni non esse speciem Hieracii, ^{indiget quoque calyst}
in iconē ipsius tel. aut. ~~et quod talis apparet~~

forchifolia Cacalia glaberrima ^{amplexicaulibus} foliis ^{ly} inferioribus lyratis amplexicaulibus
dentatis, calycibus polyphyllis
Cacalia forchifolia caule herbaceo, foliis lyratis amplexicaulibus
dentatis Lin. S.V. pag. 733.

21 *Sonchifolia Calceola* caule herbaceo foliis lyratis amplexicaulis Sp. pl. p. 1169. Mart. 463 *

S.V. p. 733. Murray Comment. Gotb. 1772. p. 79. tab. 7.

Tagetina Luzonin. flore purpureo Bot. Gaz. tab. 80 f. 12. Camel. Luz. II. 2.

+ *Senecio maderaspatensis*, Sinapios folio floribus parvis luteis Pluk. Alm. 192. t. 444 fig. 1

+ Macl. Schari Rheid. Malab. 10. p. 135. t. 60.

Sonchus amboinensis Rumph. Amb. 5. 297. tab. 103. fig. 1.

Chondrilla zeylanica minor marina folio sinapios Murr. Zeylan. 61.

Kleinia caule herbaceo, foliis lyratis Fl. Zeyl. 305 *

Herbitaria India

Caulis herbaceus glaber caulis simplex laevissimus striatus. Tota omnia amplexi
caulis remota inferiora lyrata laciniis acutis, ^{impari} *terminalium* impari dentata
acuta interdum obtusa. Superiora cordato lanceolata basi dentata superiore
integerrima. Flori paniculati pauci 3. 5 foliolo bina ^{by longis} *stipitatae* ad primum divisionem
pedunculorum. Pedunculi filiformes nudi. Calycis ^{by longis} *calyculati* foliolis ^{by longis} *oppositis*
cinereis pluribus conglutinatis. Flosculi plures purpurei

22 iniana *Caecalia* caule herbaceo, foliis lanceolatis dentatis L. Sp. pl. 1169. N. p. 734

Habitat in India

Statura sesquipedata. Caulis erectus altus, folia lato-lanceolata vix petiolata, cavaeque serrato-dentata. Pedunculi terminales, rami, longi nudis foliis uno alterove minimo.

non vid.

20-21. Candicans Cacalia herbacea, lanuginosa, foliis radicalibus inferioribus
que ovatis petiolatis crenatis, summis sessilibus ellipticis ceteris
nervis Tab. ~~LXX~~ LXXI

Habitat ad fretum magellanicum Commerçon did. In Thonin
Caulis herbaceus, erectus, pedalis et ultra, simplicissimus, foliatus,
crassitie fere penno Cyneo, uti tota planta, calycibus excep-
tis, lana nivea obducta.

Folia radicalia plura, ^{ovalia} petiolata, oblonga, ^{bipolliciana} ovalia, ^{caulina}
inferiora, uti radicalia, petiolata, ^{tripolliciana} oblonga, omnia crenato-
dentata, obtusa, ^{sequipolliciana} suprema elliptica, basi parum attenuata,
integerrima.

Petiolae radicalium et inferiorum caulis. Tri pollicum crues, plati,
basi latiori, amplexicaules, ^{multifloris} supreme in folium dilatati
Corymbus terminalis, Pedicelli uniflori, foliolis adpressi: folium
lanceolatum, acuminatum, ad basin pedicellorum, uti pedicelli
valde lanatum

Calyx simplex, foliolis lanceolatis, glabri.
Corolla magnitudine nucis congl. Corollelo longitudine calycis

23. 24. Scabra Cacalia herbacea, foliis radicalibus petiolatis oblongis den-
tatis: caulinis lanceolatis sessilibus.

Habitat in Brasilia dedit Dr. Moquin.

Caulis herbaceus, erectus, pedalis, teres, striatus, pilosus, prope stem basi,
scabriusculus, ~~in pennis~~ subnudus, simplicissimus.

Folia radicalia petiolata, bipolliciana, oblonga vel subspathulata,
dentate-subrepanda: dentibus inaequalibus mucronatis. ~~Termina-~~ Termina-
tis; pilosa, supra scabriuscula, venosa, basi parum angustiora
per petiolum parum decurrentia. Caulina parva, tria infima
versus basi caulis approximata; lanceolata, sesqui pollicia
via, sessilia, remote denticulata: dentibus, utrinque quatuor
vel quinque; una tria superiora remota, unguiculata,
lineari-lanceolata, integerrima, erecta.

Petiola radicalium longitudine foliorum, hirsuta, plana.

Floris quinque, pedunculati, alterni, versus apicem caulis

Pedicelli teretes, hirsuti: solum lineari-lanceolatum ad basin.

Singuli pedicelli et aliud in medio

Calyx simplicissimus: foliola 12. lineari oblonga, longitudine flosculorum

Pappus sessilis, ad lentam partem minutissimis denticulatus

24
angulata Cacalia herbaea, glabra, foliis inferioribus oblongis angulatis, ²dentatis
petiolatis, paniculis multifloris

Cacalia sonchifolia Forst. Catal. pl. Arab. nov. 85.

Habitat in Arabia felici

Caules ^{herbaceus,} strati, ^{in petiolum attenuati, sesqui pollicari} superne divisi. Rami floriferi, elongati, apkylli.

Folia ^{petiolum attenuatum} inferiora petiolata, oblonga, angulato-dentata, glaberrima
avenia, acuta, interdum basi sinuata, lobulorumque lanicola

^{respl.} fo. subdentata, superiora sub ramis lanicolata, integerrima
Petioli folio triplo brevior, amplexicaulis, superne dilatatus

Panicula plures, folia ramosi, fastigati. folium setaceum
sub pedunculis et pedicellis. Pedicelli capillares, versus apicem
folio setaceo.

Calyx cylindricus multiflorus, ecalyculatus; folioli quinque,
lineari-lanicolati

Targonia intermedia
conspicua. ~~Pentaurca~~ ^{sub} Calycibus ~~in~~ ^{sub} ~~axillis~~ ^{axillis} squamis ovatis subintegerrimis foliis pinnatis
y inferioribus glabris, ~~superioribus~~ ^{superioribus} ~~pubescentibus~~ ^{pubescentibus} caule tomentoso

~~Patria ignota ex Herbario Tarz. Tozzetti~~

~~Caule tomentoso erecto ^{per} ~~se~~ ^{se} paniculatis. Folia ^{patentia} ~~inferiora~~ ^{inferiora} ~~decur~~ ^{decur} ~~sive~~ ^{sive} pinnata
vel ~~pinnatifida~~ ^{pinnatifida} ~~inferiora~~ ^{inferiora} ~~petiolata~~ ^{petiolata} ~~glabra~~ ^{glabra}, ~~superiora~~ ^{superiora} ~~sessilia~~ ^{sessilia} tomentosa, ~~primis~~ ^{primis} ~~canaliculatis~~ ^{canaliculatis}
~~lanceolatis~~ ^{lanceolatis} ~~obtusis~~ ^{obtusis} ~~integerrimis~~ ^{integerrimis}. Ramulorum ~~integra~~ ^{integra}. ~~Flor~~ ^{Flor} ~~terminalis~~ ^{terminalis} ~~sessilis~~ ^{sessilis}
solitaria. Calyx ~~ovatus~~ ^{ovatus} magnitudine C. leppii ~~ante~~ ^{ante} ~~imbricata~~ ^{imbricata}, squamis glabris
acutis, ~~fenestris~~ ^{fenestris} minutis oculo armato videndis. ~~Flor~~ ^{Flor} ~~purpureus~~ ^{purpureus}?~~

~~an varietas~~ ^{an varietas} ~~si alpina~~ ^{si alpina} ~~convenit~~ ^{convenit} ~~cum~~ ^{cum} ~~figura~~ ^{figura} ~~Pent.~~ ^{Pent.} ~~majoris~~ ^{majoris} ~~lutei~~ ^{lutei} ~~Corn.~~ ^{Corn.} ~~canad.~~ ^{canad.}
~~differt~~ ^{differt} ~~vero~~ ^{vero} ~~ab~~ ^{ab} ~~aliqua~~ ^{aliqua} ~~quod~~ ^{quod} ~~ul~~ ^{ul} ~~foliolum~~ ^{foliolum} ~~impar~~ ^{impar} ~~integerrimum~~ ^{integerrimum} ~~non~~ ^{non} ~~fenestatum~~ ^{fenestatum} ~~integerrimum~~ ^{integerrimum}
~~est~~ ^{est} ~~neq~~ ^{neq} ~~fenestatum~~ ^{fenestatum}, ~~calyces~~ ^{calyces} ~~etiam~~ ^{etiam} ~~trinitis~~ ^{trinitis} ~~forme~~ ^{forme} ~~perulatis~~ ^{perulatis}. ~~habitu~~ ^{habitu} ~~addedit~~ ^{addedit} ~~ad~~ ^{ad} ~~pari~~ ^{pari}
~~culatam~~ ^{culatam} ~~sed~~ ^{sed} ~~foliula~~ ^{foliula} ~~latiora~~ ^{latiora}, ~~et~~ ^{et} ~~calyx~~ ^{calyx} ~~diversus~~ ^{diversus}

Atriplicifolia *Caecalea* caule herbaeae, foliis subcordatis dentatis sinuatis, calycibus
 quinquefloris Gronov. Sig. 110. Sp. Pl. p. 1170 L. V. p. 734.

Caecalea Virginiana glabra, foliis deltoidibus sinuatis subtus glaucis
 Monf. h. 3. p. 94. f. 7. Tab. 15. § 37.

Nardus americana procerior, foliis cosis Pluk. alt. 251. Tab. 107. f. 2.

Porphyrillum foliis deltoidibus angulatis Gronov. Sig. 1. p. 94.

Habitat in Virginia Canada &

magnitudo et ~~structura~~ ^{maximè} ~~antecedenti~~, folia inferiora cordata superiora ^{obovata} ~~obovata~~
 ra quam longa, sinuata, subtus glauca, cœvia. Differt a ~~precedente~~ ^{precedente}
 non modo figura foliorum et glaucis colore, sed et floribus quintuplo
 minoribus, tantum ex quinque floretulis compositis, cum antecedentibus
 ses gaudcant pluribus L. Sp. Pl. l. c.

~~multi spinos.~~

23 *Sarracenia* *Catalpa* cauli herbaeo foliis lanceolatis serratis decurrentibus L. Sp. pl. p. 1169. t. v. 734

Virga aurea f. *Solidago sarracenia* latifolia serrata Bauh. h. 2. p. 1063.

Virga aurea maxima radice repente f. *Doria* major reperit Monf. h. 3. p. 123.

+ *Senecio* perennis *Solidaginis* folio saillant Act. 3. 66.

+ *Conyza montana* foliis longioribus serratis flore & sulphureo albicante Thunb. Act.

Parf. 1705. p. 394.

Solidago foliis lanceolatis serratis decurrentibus caule angulato Monnei Obf.

163. Sauvag. Minipr. 14. Gouan Monf. 444

Habitat in Gallia Australi

Tacite Sen. *Sarracenia*. Caulis angulatus & costa foliorum decurrente
glaber. Folia lato-lanceolata, glabra, & qualiter denticulato-serrata. Calyx
bas. squamis inaequalibus. Calycis foliola apice spatulata. & latifolia omnes
infundibuli formae herb. Ausf.

non vid.

26 Scandens C. caule herbaceo scandente, foliis sagittatis dentatis petiolis simplicibus
26

Hort kew. 3. p. 137.

Habitat in C. b. spec. 4

non red.

16 *suaveolens* *Cacalia* caule herbaiceo, foliis ~~trilobis~~ ^{trilobis} ~~acuminatis serratis~~ hastato sagittatis
denticulatis, petiolis superne dilatatis Hort. Upf. 254 * Sp. pl. p. 1170. ~~Tab. 1. v. p.~~

734.

Habitat in Virginia Canada &

sero glauco Caulis altitudine humano sulcatus, simplicissimus. Folia sagittato triangulari
vel deltato hastata, tripliciana glabra subt. venosa subtus pallidiora ^{non} ~~denta~~
to f. inaequaliter denticulato serrata basi in petiolum producta. Flores panicula
si subfastigati. Peduncululi calycibusque glabri. bracteo ~~stipite~~ ^{basino} ad omnes pedun
di basi pedunculorum & pedicellorum. Squama nonnulla ad basin calycis Calyx
polyphyllus multiflorus, flores albi anthero flavo

Suav. Cac. caule herbaiceo erecto, foliis hastato sagittatis denticulatis,
petiolis superne dilatatis Hort. New. 3. p. 157

hastata *Galium* caule herbaceo, foliis trilobis acuminatis serratis, floribus mutatis S. p. 1170* P. V. p. 734.

Senecio foliis ex dentato trilobis acuminatis serratis Jmel. Sibirica 2. p. 136. Tab. 66.

Habitat in Sibiria 4

Caulis erectus oxyale, sulcatus, interdum rose glaucus. Folia triloba hastato deltoidea basi acuminatis, intermedia majore, inter lobos serrata, basi rotundata, subtus parum villosa, basi saepe intergerima, in petiolum prodeducta. Petiolus ampliusculi. Flores paniculato racemosi. Peduncululus pedicellique villosi. Liliopsis ad basin pedicellorum. Flores mutatis squama una alterave ad basin calycis. foliola Calycina quinque lineari oblonga obtusa. Calycis multi flori 8-10.

20) alpina ^{hirsuta} Cacalia foliis reniformi cordatis acutis denticulatis, calycibus levibus.

Tusflego alpina seu montana Dalechamp. Lugd. p. 1052.

Cacalia glabra folio Clus. Pan. 501 hi

Cacalia pyrenaea allianifolia Tournef. inf. 452.

Cacalia foliis cordatis serratis longissima petiolatis, floribus umbellatis
Hall. h. 137.

C. alpina foliis reniformi cordatis acutis denticulatis L. Sp. pl.

S. V. p. 734. Jacq. fl. Austr. vol. 3. t. 234.

Petiolus amplexicaulis
folio longiori

Caulis teres pubescens simplicissimus. Folio petiolato, cordata acuta
dentata, supra viridia subtus pallida pilis pilissimis venoso reticulata
venis pilis raris. Flores paniculati. foliolis fetaceis ad basin pedicel
lomm. pedicelli ultimi capillares squamis minutis fetaceis. Calyx tetraphy
lus foliolis lanceolato linearibus obtusis bi sive trifloris

Cacalia foliis cutaneis acutioribus et glabris B. Pin. 190

Cattalia Lobel. 4. 592.

Corollula quadrifida Scopoli

Tusflego allianifolia foliis reniformi cordatis serratis utroque glabris
innatis calycibus quinquefloris Encycl.

calycibus glabris

alliana folia

Caulis herbaceus, simplex angulatus, v. illofus. Folia petiolatis cordata, pitha-
mam ^{ata} ~~longa~~, eroso-dentata, basi acuminata, latiora quam longa, superiorum
inferne coarctata, basi auriculo amplexicali dilatata. Stipuli 5que

Catalpa peltata foliis inferioribus amplis cordatis-subrotundis angulatis dentatis, longe
petiolatis, ~~hastatis peltatis pedunculis vespertinis~~ Eni. Caulinis subhastatis basi
annulatis cymbo lato Enéyl.
β. humilior floribus glomeratis lb.

30 tomentosa *Cacalia herbacea tomentosa*, foliis cordatis petiolatis inaequaliter grosse dentatis, Caly-
cibus polyphyllis multifloris ~~se~~ petioli inauriculatis
Habitat in alpibus Aulicis alpinis ?

Affinis maxime *C. alpina* et *albifrons*, sed nec folia basi appendiculata, nec veni-
formia neque calycis 3-5 phylli ^{per} 3-5 flori. totus tomentosus inermis. ^{folia} *folia* corda-
ta, acuta, inaequaliter grosse dentata. Culo altero basi longioribus inferioribus bifur-
catis in pollicaria. Petioli amplius aequalis longitudine folii. Flores corymbosi pedunculati
et pedicellique valde tomentosi basi utique solidis minutis glabris setosis. Calyx
polyphyllus 12-15: oblongis obtusis margine membranaceis latioribus quam in reliquis
floris plures. Calyculus nullus
Sed pro varietate *C. alpina* a *David Mellard* sed revera diversa ~~folia~~ illa si radius ad-
set Senecio vel *diversa* erit.

afend

33 cchinata *Caecalia* herbaria foliis reniformibus cordatis angulato-dentatis subtus tomentosis, foliis calycinis tuberculatis L. Suppl. p. 353. L.V. 1734
Habitat in Teneriffa praecipue maritimis
Simillima *C. alpina*, et differt calycinibus tuberculatis et foliis subtus tomentosis

non vid.

ovata Q. foliis ovatis nervosis utrinque viridibus, calycibus sub
quadrifloris Walt. fl. Car. pag. 196.

Habitat in Carolina

non vidi

1
Mant. 281
bipinnata Cacalia herbaea foliis bipinnatis lineantibus L. Suppl. pag 353. L. V. 734
Cacalia pinnatifida Bergl. foliis duplicato pinnatifido: laciniis linearibus
angustis dichotomis divariatis glabri Berg. Cap. 230.

Habitat in Cap. bono spei

Totus glaber ^{caulis} Caulis erectus sulcato angulatus simplex, folia setis na sessilia, bipinnatifida
laciniis linearibus angustis. Flores in corymbo terminali. foliola linearia integerrima
ad basin omnium pedicellorum. Calyx foliolis quinque linearibus acutis basi
squamosis. Corollula ~~interdum~~ ^{approtinata} quatuor vel quinque, quinquefida.

34

acaulis *Cacalia acaulis* foliis semiteretibus, scapis unifloris Suppl. 353. J. V. 734
Habitat ad Cap. b. pui

non ord:

35 radicans Cacalia herbacea ~~repens, radicans foliis tereti-ovatis L. Suppl. p. 384 f. N. 734~~
~~foliis bipinnatis acutis subtus tomentosis, stipulis oblongis~~
~~rotundatis &c~~
Habitat ad. Cap. bono frei

non vid.

Cataloides Polygamia aequalis 2-flores

Calyx simpliciter 6-phyllus. Receptaculum nudum. Pappus pilosus dentatus

Suffruticosa

Lespley

Stemculis erectus ramulosus villosus. Folia opposita lanceolata inferiora serrata serraturis minutis gracilibus versus apicem 3-5-ellianis floris superioribus integerrima. Flores in summitate caulis plures pedunculati propriis foliis alternis uniflori foliis alternis cuneatis

Calyx tubul. Calyx 6-lobus linearis acuta lobulis brevioribus. Flores 5-6 limbo quinquefidis. Stylus et stamina 6-lobis. Semina longitudine calycis 2-fidita Pappus pilosus pappi pilosus plumeus pilis minutis

semplius undique vincto
si Cataloides distingueret a ut villosa
angustifolia C. caule fruticuloso foliis lanceolata villosa
Cuneatis
pubescentibus

habitat in monte visco Baspley & Thouron

Pae' alia

Habitat herb. Desf.
Caulis crispis fere digitis maximis, glaucus, uti folia. Tota longe petis
lata, sparsa. Pedunculi elongati. Corymbus terminalis subcompositus
Calycis foliola linear-lanceolata
an o laciniata diversa

India Catechu

Habitat in India orientali herba Martii

Caulis ramosus, lanatus, etate glabris. Rami erecti, striati. Folia longe petiolata, pollicaria, cordato-ovata, crassa, venia, nervia, profunde dentata, dentibus dentato-angulatis, Venella lanata, etate nuda. Petioli lanati, folia longiores. Pedunculus terminalis 1-2 floris, adpressis squamis lineantibus, unguiculatis, inaequalibus. Calyx polyphyllus.
Senecionis forte species et potius planta caperpi quam India, a Censurariis
viri folia forte non diversa