

Paltha

1. palustris C. foliis cordatis crenatis
2. paradoxa C. foliis cuneiformibus trifidis

cuneifolia  
paradoxa Caltha foliis cuneiformibus trifidis

Habitat in Terra del Fuego

Planta vix bipollicaris, basi testis squamis lineariibus, unguiculanibus, aridis, flexilibus, et petiolis persistentibus.

Folia radicalia, petiolata, cuneiformia, unguiculata, crassa glabra, trifida: caecina laterales lineares, obtusa, superne vix obfusa: intermedia oblonga, extorsum dilatata, apice subtrifida: lobus <sup>intermedius</sup> ~~intermedius~~ foliorum, structura caecinarum laterali

Petiolus planus, pollicaris

Scapus erectus, uni florus, superne panem craspor.

Corolla petala quinque persistentia

Oederi P. foliis caule caule simpliciter: foliis <sup>caulinis</sup> pennatis pinnis oblongis serratis  
galea obtusa

Pediculari flammea ~~Car. fl. Dan. tab. 30~~

Habitat in alpebus norvegicis

Radix tuberosa longi bipolliciaribus. Caulis erectus digitalis  
vel palmari. Folia plurima radicalia per caulina plerum

que 1. raris 2. 3. pinnis petiolata pennata. penna lata oblon

gata serratis iterum tenuissime subserulatis inter medio

majoribus floralia sessilia pennatisida costa plana lata

ut in hepatica suprema lanceolata, obtusa obscure serrulata.

Calyx sessile tubulosus 5 dentatus dentibus tribus profundioribus

Corolla lutea apice purpurea, glabra

variat caule calycibus hepaticis, vel glabris, floribus totis luteis

vel galea purpurea fere

palustris Caltha foliis cordatis crenatis  
Caltha palustris L. S. V.

802

~~prostratum  
p. 200 - dictamnus calycibus luteis foliis cordatis crenatis  
caulis quibus L. S. V. pag. 53-54  
motu ciliis quibus L. S. V. pag. 200~~

Anathus ~~resinosus~~ <sup>T.</sup> capitulis terminalibus imbricatis, <sup>6</sup> bracteis ovatis lanceolatis  
marginibus ~~subtus punctatis~~ foliis lanceolatis subserratis margine punctatis

Habitat in Regno Neapolitano 5

Plantae ~~fores~~ <sup>vel</sup> ~~digitales~~ <sup>digitales</sup> ~~vel~~ <sup>parva</sup> ~~ultra~~ <sup>ultra</sup> basi ramorum glabrae  
superne villosae rami filiformes teretes inferae glabrae et capsulae  
foliorum mucronati superne villosi subheri. folia sessilia  
unquiculae nigrescentia subtus multistrati dentibus remotis  
nervi serrato basi filicata supra levia, glabra, serie punctis  
nem utroque ad marginem unice. Bractea floribus quatuor  
vel 5 sub pagulo capitulo folia caulium simillima basi  
tantum ovata longitudine capituli. Capitula magnitudi-  
ne papi imbricata bracteis ovatis non coloratis villosis  
nerviis versus apicem raris nervis punctatis dentes  
calycis ciliati

Thymus ~~Indicatum in foliis~~ ~~Indicatum in foliis~~ ~~Indicatum in foliis~~

Paradoxa Q.

Habitat ad fretum magellanicum Commeson dedit Thourin

Planta vix magnitudine digiti minimi. Petioli uti caulis basi squamis ardis, oblongis involucri et tolia pedicelatis, oblongis, unguiculatis, glaberrimis, fusida: laciniis oblongis, intermedia major emarginata, basi ad nervum appendiculae duo lineari oblongo, folio dimidio breviori. Petioli squamis parum longiora. Scapus simplex foliis parum longior uniflorus.

~~Grana~~

Capitolo 3 bralato

bayialensis Palka

Habitat in Siberia herb. Mus.

Tota glabra. Folia petiolata, semipollicaria, cordato-subrotunda, integerrima.

Caules repentes. Flores parvi



1 palustris Caltha L. V. fl. Dan. t. 668.

Populago major Tabern. u. 750.

Caltha palustris flore simplicis Bauh. pin. 276.

B. Populago minor Tabern. u. 750

V. Caltha palustris flore pleno Bauh. pin. 276

Habitat in Humidiusculis Europae &

Caulis decumbentes; Vertices, striati; glabri. Folia plerumque  
radicalia, reniformia petiolata glabra venosa, serrata  
caulium pauca, ~~sepi~~ petiolata cordata acuta crenata  
pedunculul. Terminalis uniflorus. Petala obovata

palustris C. foliis cordatis crenatis

Secund: specimen alexand. V. G. N. B.