

Calypso

velloso P. foliis oblongo-ovatis, subtus pedunculati cymbricis axilla  
ribus trichotomi villosi

Habitat in Andia occidentali & A

Rami dichotomi superne subtetragoni cortice cinereo puncta  
to glabri juniores tomento tenuisimo tecto. Folia petio  
lata, opposita pollicari basi acuta apice parum  
acuminata acumene obtuso supra glabra subtus ut  
vix venosa subtus punctis ve elevati supra junioribus  
subimpresis. ad ultra utroque glabra juniora subtus  
e villi ut perceptibilibus cinerea. Pedunculi axilla  
res oppositi trichotomi foliis longiores. Flores tres sepales  
in ramulis ultimi calycis ubi pedunculi tomento tenui  
tecto

una ramulis  
terminant

ca  
ca. Zeyheri G. Don

androsomoides <sup>myrtus</sup> racemis axillaribus, terminalibusque subcompositis.  
pedicellis trifloris, foliis ovato-oblongis sessilibus

Myrtus androsomoides pedunculis trifido multifloris, foliis

subovatis sessilibus Lin. S. V. p. 462. fl. zeyl. 1814

Habitat in ~~Asia~~ <sup>Reyona</sup> ~~et~~ <sup>heperna</sup> ~~Reyona~~  
Totus glaber. Rami ~~superne obfcuri~~ ~~tetragoni~~ ~~teretifiduli~~

Folia opposita, maxima hujus generis, palmaria et ultra  
venosa, <sup>puberula obfca vena obfca et vena</sup> parum apice attenuata, obtusa, utrinque  
impuñctata

Racemi terminales gemini, ~~interdum quatuor~~: axillares

~~oppositi~~, ~~solitarii~~ ~~compositi~~ ~~brachiati~~; tripolliares. Pedun-

culi tetragoni; ~~trifidi~~ <sup>compres</sup> ~~partiales~~ <sup>ultimi terge</sup> oppositi, patentissimi  
remoti; inferiores trifidi; pedicelli ~~triflori~~ <sup>triflori</sup> brevissimi

Bractea <sup>una flori</sup> subulata, brevissima ad basin singuli <sup>rami</sup> ~~rami~~ <sup>pedicelli</sup> ~~pedicelli~~  
et pedicelli ~~et pedicelli~~, nulla ad basin caly-

Calyx <sup>is</sup> obovatus, magnitudine seminis <sup>obfca</sup> ~~quadrident~~

100 androsomoides Calyx <sup>tatur</sup> ~~trident~~

elliptica Ternstroemia foliis ellipticis acutis integerrimis, pedunculis lateralibus  
Schw. prodr. pag. 81.

Habitat in India occidentali Dr. Joseph

Rami cortice nigro, glabro

Folia superne in ramis parva, petiolata, bipollicaria et ultra, glabra  
nona coriacea, margine parum reflexa, utrinque punctis numerosis

minutis elevatis adpressa, basi angustiora. <sup>petiolis</sup> <sup>supra subarenia nuda</sup>

parum supra  
folia

Pedunculi <sup>pet. foliis breviss</sup> <sup>foliis brevissimi</sup> unguiculatis supra planus subtus convexus

Calycis foliola obtusa, coriacea

elliptica

~~Caryophyllifolia~~ C. foliis lanceolato ellipticis subpetiolatis acuminatis  
notidi, racemi axillaribus <sup>terminalibus</sup> et terminalibusque, pedicelli subpet. flori.

Penn. mar. Arch. Mal. s. pag. 57. Tab. 29

Caryophylli languiscentis vi aromatici Plukh. alm. 100. Tab. 274. fig. 2.

an  
cum foliis petio  
latis et pedunculi  
longioris et laxioris

~~an~~ Eugenia caryophyllifolia foliis ovato-lanceolatis  
acuminatis venis penninerviatis calycis truncato la Mart.  
Encycl. ~~h. 1. 1. 1.~~

Habitat in Andia Orientali

Rami teretes glabri inferne cortice cinereo superiore purpurascante  
covi. Folia inferna breviter petiolata superiora sessilia, tripollia  
na utrinque glabra, subtilesime nervosa venosa, <sup>compuncta</sup> supra  
nitidiuscula subtis parva pallidiora, margine plani integerrima  
pollicem lata acuminata obtusa. Pedunculi axillares oppositi  
terminales saepe 40x folio duplo breviores erecti <sup>2-6</sup> flori com  
pressi fusco purpurascenti parvales oppositi ~~4-6~~ flori  
A. C. in apice singulis brevissime pedicellati umbellati  
differt ab androsamode foliis angustioribus floribus minoribus

Puparia

Caryophyllifolia C. foliis lanceolato ellipticis petiolatis acuminatis, racem  
is paniculatis lateralibus terminalibus

Caryophyllus Conguescente aromaticus Plukn. alm. 80. Tab. 274.

fig. 2.  
Penna Nyasa Rheed. Mal. 5. p. 57. Tab. 29.

~~Caryophyllus~~ Caryophyllifolia La Mark. Care:

Habitat in India orientali &

Rami glabri. Folia opposita pariter integerrima 4-5 pollicum  
pollicem lata. pauci ramuli panicula terminali floribus 6-7.

sessilibus

an ab elliptica diversa differre videtur folus petiolatis  
Racemus longioribus latioribus

*Caryophyllata* M. pedunculis trifida multifloris foliis obovatis Jacq  
obf. 2 p. 1124 fl. Zeyl. 103.

~~*Caryophyllus aromaticus* in id. occidentalis foliis et fructu  
rotundis Plukh. atm. 10 p. 1155. fig. 3.~~

~~Habitat in Zeylona~~

sq. Jacq. et Plukh. ad aërem Sw. pertinet



conyophyllatus  
Conyobos

C. foliis petiolatis ellipticis obovatisque, corymbis terminalibus corym-  
piti

Habitat in India Orientali in Kottler &

Aframiter glabra, cortice griseo pallide fere neri. folia petiolata  
opposita glabra bipollicaria lanicolata elliptica saepe  
pleniusque basi attenuata obtusa plana, subtilesque obscure  
nervis confertis nervis venis, <sup>sapida</sup> <sup>subtus</sup> <sup>superioribus</sup> ~~sa~~ ~~atque~~ ~~elevato~~  
punctata. Corymbi terminalis rami ultimi triflori pluribus se-  
pibus.

quadrat optime cum Myra Rheide Malab. 7.

~~est Myrtus conyophyllata C. f.~~

variat foliis obovatis truncatis et corymbis

Myrtus conyophyllata foliis obovatis Lund. no 103. ex. lufp  
Synonymo Platan

Horatium

1 fagineum D.

*Phytraulica* *C* arborea, pedunculis terminalibus paniculatis trichotomis  
tomentosi, foliis ovatis apice attenuatis lvs. prodr. p. 29

*Phytraulica* arbor, foliis ovatis glabris oppositis, racemis terminalibus  
Brown Jam. 239. Tab. 37. fig. 2.

*Myrtus Phytraulica*, pedunculis dichotomis paniculatis tomentosi  
foliis geminis, subovatis terminalibus lvs. Am. 5. p. 39. *P.*

*Habitat in* Jamaica

Rami vertici oppositi glabri, superne compressi sulci ad geni-  
cula parva lacinosa. Folia petiolata opposita 5-6 polli-  
cari elliptica acuminata acumine obtuso submarginata  
integerrima utrinque glabra obscure oblique nervosa  
ovaria, supra lacin nitidissima. Petiolus b. lami ungui-  
culans fusco purpureo. Pedunculi terminales breves  
geminati uti calycis et germen ferrugineo tomentosi

rigida *C. abouxiensis*, pedunculis solitariis axillaribus subtrifloris, foliis  
ovatis acutis convexis avenis rigidis *Sw. pr. pag. 80*

Habitat in Jamaica;

Habitus Myrti communis ramis teretes glabri oppositi apice super-  
ne compressi virgati cortice unevanescente. Folia brevissime  
petiolata, opposita lanceolato-oblonga attenuata basi acuta  
apice obtusa <sup>an</sup> anevra ~~virgata~~ ~~sub~~ subavenis, utraque  
glabra <sup>impunctata</sup> integerrima ~~supra nitida~~ ~~subtus~~ pallidiora ~~opposita~~  
~~nitida~~ utraque sex pollicaria. pedunculi axillares

9<sup>ta</sup> simili Eug: foetida ut facile unam pro alteram  
superes in hac vero folia impunctata in illa utraque

Zyzygium C. arborens, pedunculis axillaribus trichotomis patulis, foliis  
ovatis obtusis, ramis dichotomis Sw. prodr pag. 29

Zyzygium fruticosum, foliis ovatis nitidis et ramulis ubique  
pergatis Brown. Jam. 240 tab. 7. fig. 2.

Myrtus Zyzygium pedunculis multifloris, foliis geminis subrotatis  
terminalibus, ramis dichotomis Lam. acad. 5. pag. 398 & Lin

S. V. pag

Habitat in Jamaica