

Campsis

adrepens Campfi Lour. fl. Coch. pag 370

Habitat prope Cantonem

Caulis fruticosus, scandens inermis radiculis lateralibus. folia
bipinnata: foliola ovato-lanceolata, pinnata, glabra. flores coerulei
magis rari terminaliter compositi. Cal. 5-neris, 5-fidus lacinis
subaequalibus. Cor. infundibuliformis. Lacinis rotundatis aequalibus
anthera 2. Stylus filiformis. Stigma spatulatum bifidum.
Capsula longa, 4-neris, 2-valvis biloculari polysperma. Semine subrotunda

~~festuosa Justicia frutescens foliis ellipticis, thyrsis terminalibus L.V. p. 62. g. Mart. 172. *~~
~~Justicia paniculata~~
~~Willd.~~

~~Solia nondum habitata in India, Arabiis & articulatius articulis pollicatibus
Petiolus pollicaris ^{ultimo trip} triplex ramis teretibus glabris, purpureis facie Phlogis. Solia petiolata
elliptico lanceolata acuminata integerrima frithamoa glabra.
Racemi ^{terminali} ~~terminali~~ ^{pluribus} ~~pluribus~~ ^{scapo} ~~scapo~~ ^{compositis} ~~compositis~~ ^{racemulis} ~~racemulis~~
bractea opposita pollicaribus axillaribus ~~axillis~~ ^{ex} ~~axillis~~ ^{approximatis} ~~approximatis~~
utroque foliario Pedunculus in pollicaris. Flores umbellati secundi magnitudinis grani
suborbibus oblongis ^{caulis} ~~caulis~~ ^{exterioris} ~~exterioris~~ ^{lanceolatis} ~~lanceolatis~~ ^{acuminatis} ~~acuminatis~~ ^{glabris} ~~glabris~~ ^{interioris} ~~interioris~~
pediculis deciduis ^{pubescentibus} ~~pubescentibus~~.~~

~~Planta arabica et India differt quod duplo minor, foliis subtus venis
racemis axillaribus non vero ^{sepe binis} ~~racemis~~ ^{axillaribus} ~~axillaribus~~ ^{in racemulo elongato} ~~in racemulo elongato~~ ^{articulis ramorum} ~~articulis ramorum~~ ^{tripollicares} ~~tripollicares~~ ^{andiseja planta} ~~andiseja planta~~
ment: altera p. 173.~~

~~lyn. herb. bot. Tom. 9. Verae sunt uti et est facie h. p. arabica~~