

# Capsicum

1. annuum C. caule herbaceo, pedunculis solitariis
2. grossum C. caule suffrutescente, fructibus incrassatis variis
3. sinense C. caule fruticoso, floribus fructibusque pendulis
4. baccatum C. caule fruticoso, locis pedunculis geminis fructibus subglobosis
5. frutescens C. caule fruticoso foliis scabris pedunculis solitariis oblongis  
erectis

Caulis herbaceus,  
Lanceolatum Capsicum (foliis lanceolatis, pedunculis solitariis)

Habitat in India Orientali.

Caulis herbaceus, angulatus, uti tota planta glaber. Rami alterni <sup>semis</sup> ~~sub~~  
si subdichotomi. Folia petiolata, bipollinaria, utrinque attenuata  
Peruvia, Pedunci axillares, solitarii, uniflori. & cores parvi

pubescens Capsicum

Capsicum pubescens; caule suffruticoso, foliis ovatis acutis apice obliquo,  
pedunculis angulatis, fructibus subrotundis rotantibus &c. Penn. 2. pag. 30

Habitat in Peruvia cultis

~~Pa~~ Pubescens. Caule suffruticosa, 4-5 pedalis, erectus, angulatus, nodosus, dichoto-  
mum, lincis dispersis, geniculis atropurpureis, folia e lateribus singulis  
dichotomis, solitaria, ovata, acuta, apice obliquo, integerrima, venis  
prima, pro bullata, nervo superne prominulo. Petioli incurvati, supra  
basili, breves prope basin articulati. Pedunculi e dichotomis, solita-  
rii, rarius gemini, uniflori, petioli longiores, angulati, superne incurvati.  
Corolla violacea; laciniis acuta. Baccis subrotunda, depressiuscula,  
magnitudine mali medoceri, vix di lateribus

1 annum Q. caule herbaceo pedunculis solitariis Len. l. v.

Piper indicum vulgatifimum Bauh. pin. 102.

Sallia-capo-molago Rheid. Mal. 2. p. 7. 15.

Habitat in America meridionali ☉

Caulis erectus pedalis striatus glaber superne dichotome divisus  
interdum glaber interdum pubescens pro parte superne. Folia  
petiolata utrinque acuminata glabra vel venis et margine  
centes pubescentia. Pedunculi in diversis ramorum soli  
solitarii uniflori, fructu superne incrassati. Pedum  
fructus penduli oblongi

2 *grossum* Pappium caule suffrutescente, fructibus in craspatis variis Lin. Mant 2774  
L.V. p. 226.

Pappium fructu longo, ventre tumido; per summum tetragono Tournef.  
G. R. H. 152.

13. Pappium fructu bifido Tournef. G. R. H. 152.

Habitat in India 4-t

Similemum P. annuus, sed caulis perennans, spithamou, subramosus.

Fructus relatione ad plantam maximis, malifere magnitudine,  
varia in variis individuis forma, solitarius; erectus Lin. M. C.

Tolia ovata utraque attenua glabra alterna in ramuli  
gemina vel terna. Pedunculis & ramulis et axillari pedu  
solitarii vel subgemini.

conicum Capf. caule suffruticante, fructibus ovato-conicis pumatis La  
marki P. En 2. p. 26.  
Habitat in Andes 5.  
Folia minora quam in grupo

michi ignotum

ternisiforme Caps. caule suffruticeo, fructibus subglobosis foliariis erectis  
La Mart. P. E. 2. pag. 26  
Piper filiqua parva Bauh. hort. 2. p. 944  
Habitat in Brasilia 5





rugosum Piper foliis ovato-oblongis venosis glabris basi subaequalibus subtus  
reticulatis, ramis teribus, petioli simplicibus.

Habitat in Cayenna. et in Rohr 5.

Rami glabri, <sup>recti,</sup> teres, inferne teretes, superne <sup>recti,</sup> laeves, angulati, articulati,  
ope ~~se~~ lentis quasi pulverulenti: Geniuli nodosi.

Folia petiolata, alterna, quinque-vel sex pollicaria, oblonga, basi ~~ovata~~  
ovata, apice acuta, venoso-reticulata, subrugosa, profertim subtus  
e venis exstantibus, basi utrinque aequalia.

Petioli unguiculari, supra canaliculati, subtus acute carinati.

Spica oppositifolia, pedunculata, solitaria, semipollicari (folia  
<sup>obovata</sup> ~~obovata~~ <sup>abrupte</sup> ~~abrupte~~ <sup>subulata</sup> longiores) teretes, mucrone brevi nudo absque flori-  
bus terminata.

Pedunculi longitudine petioli.

Piper citrifolium fide ~~her~~ La Martre

sinense & caule fruticoso, floribus fructibusque pendulis Lin. l. V. p. 226.  
Gacyn. hort. 3. t. 67.  
Habitat in China

non vidi

4 ~~barbatum~~ *Cappium* caule fruticoso lavi, pedunculis geminis Linn. Mant. 47. x  
L. V. p. 226

~~*Cappium frutescens* L.~~

*Piper filiquosum* magnitudinis baccarum apparat. Bauh. h. 2. p. 944  
*Piper brasiliense* Perita Clus. cur. 55. Pumph. amb. 5. p. 247. t. 80.  
fig. 2.

*Cappium minus* fructu parvo pyramidalis erecto Sloan. h. 1. p. 240  
t. 146. fig. 2.

*Cappium* fructu minimo conico rubro Brown. jam. 176.  
Habitat in Indis

Differt parum a fruticoso: caule venenose, magis frutescente, nec spica  
breviflora. Baccis minimis ovatis, magnitudine Rubrorum; nec magnitudi-  
dine Persici. Ramis divariatis; nec angulo recto ex eundem. Caly-  
cibus dentibus subulatis patulis, nec obsoletis.

Rami tetragoni. Folia ovata opposita interdum terna ovata  
attenuata sesquipollicaria al. <sup>in ramulis tantum</sup> <sup>loco alterna</sup> ~~obtusis~~ vel alterna ~~trisplo~~  
minorum rami ad angulum rectum ~~erecti~~  
fructu subgloboso

Luteum Capf caula frute pienta flexuosa ramuli petiolosque hirsuti, fructi-  
bus ovatis pendulis tuberculatis Ca Mart. T. E. v. p. 26  
Habitat in India orientali  
fructu luteo

5. frutescens C. caule fruticoso scabriusculo, pedunculis solitariis Linn. S. V.

*Capficum brasiliense* Claf. Coed. 340. fig. 2.

*Capficum indicum* Rumph. amb. 5. p. 247. T. 80. fig. 1. 3. 4.

Habitat in India &

fructu ovato elongato