

Carex

helvetica Det. Muhl.

alba Caru

Cypripedium polytachyum minimum specis per maturitatem albiam
debus Tournef. Inst. v. h.

Plukn. Voyb. 91. fig. 2. herb. Mus.

~~refurn. E. p. 404. tab. 232. *Cypripedium* *Caru* *albium* *per* *maturitatem* *albiam* *debus* *Tournef.* *Inst.* *v.* *h.*~~

~~Dentation~~ ~~Crym...~~ ~~foliis ovorari emarginatis, pedunculis capillaribus foli-~~
ramulis ^{ancipitibus} ~~compressis~~ dentatis

Habitat in ~~India~~ ~~Francia~~ ~~et~~ ~~India~~ ~~orientali~~

Rami ~~ramosi~~ ~~paniculati~~, inferne teretes punctis albis adpressi
Ramuli ^{ancipiti} ~~compressi~~, articulati alternatim dentatis apice sive
subulatis. ~~Solia~~ ^{pedicelata} ~~integerrima~~ ~~subar-~~
bitibus glauca unguiculata. Pedunculi axillares solitarii
capillares unisflori sub flore unisulcati

~~Erythroxylon hypericifolium foliis numerosis ovatis, ramis punctatis scabris,
pedunculis longis capillaribus, fructu tuberculato cavan. Desf. Tab 2~~

~~Erythroxylon~~

Lamarck

acuta Q. digna prois filiformibus; foemineis inflorescentibus
nutantibus fructiferis erectis capsulis acutiusculis apice in-
verso.

Carex acuta B. Lea. fl. Swer

Carex gracilis Curt. fl. Lond

Habitat

Radix repens. Folia angusta erecta, apice pendula ulmo bre-
via. Culmiratione soli vario magnitudinis, bipedalis et
ultra, acute triquetra sub inflorescentia saepe cernuis.
Spica mascula duo raro tres ~~idiformem~~ 1-3 pollicares
squamis oblongis obtusis nigris inferiores breviores. Spica
foeminea 3., uti mascula filiformes sub inflorescentia
nutante, maturitate erecta apice saepe mascula. Squamae
oblongae acutae nigrae. Bractea inferior calamo parum longior
Capsula suboctofararia imbricata orata, acutiuscula, com-
presso.

Carex ^{serena} ~~procon~~ la Mantini qua fructa W. & Henning.

acuta & spicis masculis pluribus obtusis, squamis obtusiusculis, caule angu-
lato ~~Curt. Fl. Lond.~~ Hot. Kew. 3. p. 331

rufa a Gramen Cyperoides Bauh. pin. latifolium spica rufa f. caule
triangulo Bauh. pin. 6. Scheuch. agr. 450

Gramen Cyperoides cum paniculis nigris Bauh. hist. 2. p. 494.

Carex spicis masculis ternis, femininis numerosis erectis,
brachiis petiolatis, capsulis bicornibus Hall. hist. 1404

~~Cyperoides latifolium spica rufa~~

Cyperoides foliis canyophyllis, spicis latioribus squamis curtis
Michx. gen. 62. t. 32. fig. 12.

nigra β. Carex

^{acuta} Carex spicis masculis pluribus, femininis subsessilibus, capsu-
lis obtusiusculis Lin. Spic. pl. 1. V.

~~nigra β. Gramen Cyperoides, foliis canyophyllis, vulgato nomine~~
~~hist. 1492.~~

Habitat ubique in Europa pin. agris a in siccioribus.

~~Carex acuta fl. Lond. et V. paludosa Gooden.~~
Var. a ad Car. Britan. Good. forte amandanda
var. β. ad acutam Good. pertinet
vid. Car. ac. Good.

paludosa C. spicis oblongis subobtusis; masculinarum squamis obtusis: foeminae
nearum lanceolatis, capsulis ovato-lanceolatis apice subdentatis
Gooden. in A. J. Lin. 2. p. 202.

Carex acuta fl. Lond.

Habitat in palustribus ad ripas flosanum

Radix repens. Folia semperquam lata, glaucis vix erecta
culmo breviori, basi rubra. Culmi pedalis bipedalis erectus
inaequaliter triente torqueter. Spica plerumque sex angulis
approximata, triquetra angulis obtusis pilis obtusa. Squama
oblonga obtusa. Spica foeminea 3. erecta remote inferior
pedunculata. Squama lanceolata valde attenuata capsula
saepe longioris. Bractea inferiori culmo saepe longior. Capsula
octofloram imbricata ovato lanceolata apice obtuse biden-
tato nervosa. Stylis 3.

crassa R. specis femineis brevissime pedunculatis cylindricis
capsulis ventricosis stratis acuminatis Willden. no 1236
Ech-grum. no 59.
Habitat in Germania.

mihi nota

Ciliata & spicis pedunculatis cylindricis, squamis foeminearum subula
& serrulatis, capsulis superne ciliatis

Habitat in Barbaria

Culmi triquetri, striati, angulis glabri, bipedales. Folia sub
serrata

spicis plerumque octo, 400 masculis, quatuor foemineis
omnes pedunculatae proter duo inter mediae ^{remota serrata}
teretes, 3-4 pollicares acuta, bractea foliacea sub 5 in penon
infimo fulms alborei, pedunculi a vagina eaque lon
giores. Spicis masculis tenuiores squamis cuneato oblongis
obtusis foemineis dorso margineque albidis apice ope latis
serrulatis. Spicis foemineis apice ^{ma} immatura 6 pollicia
res a medio ad apicem masculis tenuiores, natura 10 pollicares
obtusis squamis albidis lanceolatis attenuatis capsula longioribus
apice serrulatis ^{non tamen veris} virgatis dorso linea ^{inter un} ~~viridis~~ capsula

foemineis

squamis multo latioribus obovatis convexo triquetris cellis brevi bifido
maturo superne ferrugineo ope latis ruguloso margine acuto super
ne ciliato Stylis 2.

4
gracilis & specis masculis et foemenais pluribus subfili formibus; floribus
digynis Curtis fl. lond Hort. Kew. 3. pag. 331

~~Habita in Anglia~~

~~Carot acuta variet. 13. Lin. fl. Svec: n. 897~~

acuta Cor. digyna specis filiformibus; foemenais in florentibus nutan-
tibus, fructiferis erectis, capsulis acutiusculis apice induratis Gordon.
in ac. f. Lin. 2. p. 203.

polytricha ♂ spicis foeminei numerosi ternatis pedunculatis
pedunculo intermedio brevior.

Habitat in nova Zelandia

encetonum P. spica mascula oblongo-ovata, foemineis approximatis breviter
ovatis fere semis rotundatis margine membranaceis, fere capsula
subtomentosis Schrank bayesht. flor. 1. p. 207

~~Carex encetonum~~

~~Poll. palat. 006~~

~~huc refertur a Schrank. Scheuchz. Tab. 10 fig. 10~~

~~Habitat in Germania~~

nihil ignote

Hallii
Falta
notus

C. fructus masculis androgynis que pluribus, foemineis
sepius, capsulis ellipticis puberulis Hall. bot. p. no. 1399

Habitat in A. Helvetia

non vidi

Ampullacea C. spicis filiformibus: masculis tenuioribus; foemineis ter-
tibus erectis, capsulis inflatis globosis aristato - rostrato divarica

Hall. 1409 β
Lec. Wildem
Carex inflata

Sci gooden in art. bot. lin. 2. p. 207 x
gramen cyperoides medium angustifolium, spicis terribus
erectis flaviscentibus Monf. Okon. 3. p. 242. J. P. t. 12. fig. 8.

Wied
Carex Hall.
hist. stirp. no
1401. excl. p.

Carex vesiana var. a fl. angl. 413 Poll. 895
Carex rostrata Witte nuy. p. 1059
Habitat in aquis Angliae Germaniae Danicae Norvegiae &
Sueciae

Synon. Monf.
Carex bifurca
Schrank. boyer
sch. flora

radix repens. Elia glauca - viridia, angusta culmo altior
Culmus 1-2 pedalis erectus inf. triquetris inferne angulis obtusi-
superne acuti. Spica masculo 2-3. lineari filiformer subli-

Spica foeminea
interdum natantes

pollinaria squamo obt. lineari oblongo obtusiuscula fusca
vel p. margine pallida vel tota pallida. Spica foeminea

Carex longifolia
Thuiller fl.
Panf.

cylindrico plerumque 2-3. raro 1. bipollinaria erecto
natantes pedunculato. Squama lanceolata acuta capsulis
longioribus bractea ad spicam omnes excepto terminali

inferiores culmo altiores. Capsula 7-8 bifidam imbricatam
globosam acuminatam ore bidentata, nervosa. v. s. t. e

inferiores culmo altiores. Capsula 7-8 bifidam imbricatam
globosam acuminatam ore bidentata, nervosa. v. s. t. e

Cyperoides polystachyon *spicis verticibus erectis* Tournes. *Appl. v. h.* herb mus.
Cyperoides spicis longioribus flavescens Tournes. *4. v. h.* herb Tournes

vesicaria Q. spicis masculis pluribus, foemineis subsessilibus, capsulis obtu-
 sifoliis fl. lan. l. v. p. 8l. dan. 647
 Germen Cyperoides angustifolium, spicis longis erectis Bauh.
 pan. 6.
Cyperoides vesicarium, spicis viridantibus et subsessilibus Scheuch.
 gram. 470.

Carex spicis masculis una pluribusve femininis erectis tereti-
 bus; capsulis ampullatis rostratis Hall. hist. 1401.

~~Carex vesicaria spicis masculis pluribus, foemineis subse-
 silibus, capsulis ampullatis rostratis Hall. hist. 1401.
Cyperoides vesicarium duncubatis, capsulis globosis inflatis acuminatis
 glabrum spica pedicula Hudson. angl. 413~~

Germen Cyperoides major procerum, spicis viridis tereti-
 bus flarescentibus Mon. f. 3. 242. J. O. t. 12. fig. 6.
 530 herb. mus. Habitat in palustribus et a quibus
 et Tournet Carex acutangula Ehrhart

vesicaria Q. spicis masculis circantibus; foemineis oblongis patentibus
 capsulis inflatis oblongis rostrato-acuminatis patentibus Good.
 inflata Hudson. in act. f. lin. 2. pag. 205

rumula C. spicis masculis duabus terminalibus sessilibus, foeminis dua-
bus pedunculatis oblongis erectis Thunb. Fl. jap. 394 L.V. p. 846

Habitat in Japonia arenosis, iuxta margines riarum rarius
Culmus teres finis, subnullus. Folia culmo longiora, attenuata, con-
voluta, glabra. Spica monoica; semines duo, pedunculato, oblongo glabra
glumis turgidis ovatis acutis glabris; masculo duo, lineares: inferior
sessilis, minor, superior duplo longior, terminalis. ~~Thunb.~~ Maxime affinis
C. vesicario ut character vix distinguenda nisi quod fere acutis
Thunb.

non. vidi

mixta C. spica Yerminali mascula, altera androgyna superne
mascula, reliquis foemineis seplibus, capsulis breviter bi
furiatis Hall. hist. fl. no 1390

Cyperoides latifolium, spica spadicea viridi major Scheuchz. agrost. p. 464
Cyperoides palustris angustifolium
Scheuchz. agrost. pag 452

Carex Scheuchzeri Honkeny Synop. 1. pag 391.

Carex spadicea Elwes n. 779

non vidi

ferax

B. spica terminali mascula sessili, reliquis masculis cum
androgynis superne masculis et foemineis pedunculatis erectis
capsulis inflatis acuminatis fl. Tub. p. 200

non vidi

quinata Q. fructis masculis terminalibus androgynaeque mediae inferae
masculae sessilibus, foemineis pedunculatis erectis fl. Tubaz.
p. 209

mibi ignota

palustris C. spicis foemineis sesquibus, capsulis ovatis brevifloris mucronatis Hall. hist. purp. no 1400

Cyperoides foliis caryophyllois, spicis erectis sesquibus, capsulis ovatis brevifloris mucronatis Ray Synopt. p. 410

Habitat in polydopi

Spica multiflora. Capsula compressa apice integro

Carex elata All. fl. Pedem no 2344.

affinis vesiciana

michi ignota

manida Q. specis masculi androgynusque manidi: foemineas
gracillimis supra sessili, reliquis longissime petiolatis
Hall. bot. fr. no 1402.

Cyperoides sylvestris, specis tenuibus a radicibus albis
tribus Schaeuberi p. 454
Habitat in sylvis montis

non vid. prope vesicularum

herba C. spica remotis: masculis pluribus, foemineis subpedunculatis erectis.

Capsulis herba Lin. Spic. pl. p. 1384 ~~Fl. Dan. tab. 479~~ ~~Leers. fl. her. 725.~~ ~~t. 6. 16. f. 3~~

Cyperoides polytachyon Canadense Pluk. alm. 170. t. 34. fig. 6.
monf. her. 3. p. 243. f. 8. t. 12. fig. 10

Habitat in Europa sabulosis ^{repente pilosus}
Radix squamis setosis vestita, repens. Culmi saepe pedales foliosi, tri-
quetri superne pilosius. folia caninata margine canny scabra
vaginis que villosa. folia floralia sub spicis foemineis culmo
altiora. spica remota. masculi terminales gemini distantes,
teretis bractea membranacea spicula brevior, villosi. foeminei
duo vel tres inferne: squamis membranaceis ovatis ~~tri~~ acuminatis
glabris. ^{Capsulis longioribus} Capsula ventricose acuminata hispida subbicornes al-
tero latere planiuscula

herba C. pilosa spica striatis: oblongis; foemineis remoti vaginati
vagini hinc lanato villosi, capsulis herba Good. in A.
soc. her. 2. pag. 200

Synon. flor. Dan. ad filiformem peduncul

ternaria C. fructu summis erectis, masculis duabus; Terminali ses-
sili altera subpedunculata, foeminea pedunculata fl.
Tub. p. 200. 209

non vid.

mutans G. speciei oblongis, masculis sesplebis: foemineis nutantibus
androgynaeque media suberecta, pedunculatis axillari-
bus fl. sub. p. 206

michi ignota

polygama C. fructis mastulis Verminalibus: androgynis superne
mastulis axillaribus: pendulis Pler. tub. p. 206

non vidi

erecta Q. spicis mapulis terminalibus approximatis erectis, andro
gynis superne mapulis pedunculatis axillaribus erectis Flor
sub. p. 209

michi ignota

παροδοχα

Q. fructus duabus supremis masculis oblongis gracilibus
simplicibus, & legum compositis. fructibus inferioribus compositis
• masculis, terminali androgyna superior mascula flor sub.
p. 290

non vidi

Culmus erectus sesquipedalis glaber inferne foliosus vaginatus
inque ter angulatus acutus subscabri venis. folia angusta
marginata ut scabra Culmo alborum ^{hymplex} spicula ^{simplex} superiore androgyna
Cistis 12. Distanter linearis-oblonga usque foliis subscabris
bracteae ovatae acute margine albido ferruginea capsula dimidio
brevioris. capsula ovata attenuata dorso convexo intus plana
acuta quae integra et pedicula.

Carex officinis muricata in Holstria lecta