

Phloris

porquoides - Nardus fructu secunda revoluta angustata, florum duplici serie
La Mart. T. Enc. pag. 152.

non f. p. V. B. f. quat.
Habitat in America

non vidi

ut sequens ad Chlonis spectant

andropogonoides N. spica secunda recta flosculis anfracti basi barbati

Habitat in India orientali.

Culmi plures, simplicissimi erecti digitati vel palmati capillares basi ^{truncati} ~~capiti~~
nati. Folia inferna lanceolata, culmi ultimum culmi subulatum, glabrum
spica terminalis pollicaris. Flores ab altero tantum latere alterni
erecti flosculi basi barbati. Vagula altera f. linearis attenuata altera
congrua apice ~~sub~~ ^{per} rare pilosa latere interno infra apicem anfracta
tota fere longitudine flosculi

Anthoxanthum avenaceum: spica secunda calycibus pilosis; flosculi anfracta longa ~~tota~~ geniculata. Retz. off. 3. pag. 5.

Chloris

repens Chlois

Habitat in Carolina Richard

notes

1. Spaspaloides Cynosuavis fructu digitatis, calycibus bifloris subglobosis. valvula
exterior calycis aristata, lobisulis minutis
Habitat in Cap. b. per Da. Bulow
gramen pedale, Culmo simplicis fimo subaphyllo
Folia radicalia vaginantis, lineari-lanceolata, bipollicaria
acutiuscula, Vagina ~~compressa~~ duplo angustiora. Vagina
compressa, ^{equitantes}
Spicae terminales ^{quasi sessiles}, ~~sa~~ sesquipollicares et ultra, basi
lanata, lineares, angustae, ferrugineae, nitentes.
Flores ^{unicat. males} magnitudine seminis milii, alterni, conferti, obtusissimi
Calycis valvula exterior, corolla brevior, ovata ^{f. paucum} infra
apicem aristata, emarginata, caninata; altera minor
Lobisuli gemini, calycis longiores: valvulis exterioribus carnalis;
canna margine albo ciliata

Chiloni petra Sw.
Agrosti complanata Hort. Ken

Filiformis Cynosuavis Spica ^{solitaria} simplicis disticha, calycibus pubulatis trifloris ^{anastats.} muticis flosculo
sterili mutico.
Habitat in copula Francis

Culmi repentes, fili ramosi, fili formes, inferne compressi, floriferi
erecti, solido unido pubulato. articuli bases pollicares

+ setacea equisanta folia ad nodos aggregata, pubulata sesqui pollicares unduplicata
nodium. Spica terminalis, pollicaris, linearis. Floris distis alternis

Calycis valvula patente, pubulata parvi, valvulis subaequalibus,
longitudine flosculorum calyce longior

Corolla valvula exterior sub apice anastata; anata longitudine
flosculorum flos alter ses fili hermaphroditus, alter breve pedi
cellatus mapulis, tertius sterilis muticus.

radicata Agrostis. Specis subquinis cruciatis basi villosis, valvulis petaloideis anis-
tis. Am. Acad. S. p. 392 & S. V. p. 111.

Gramen dactylon specis gracilibus plerumque quatuor cruciformiter
dispositis Noan. h. 1. p. 111. t. 68. fig. 3.

Gramen cruciatum aspergens, specis puberulis tenuioribus et longioribus
deorsum frugiferis Brown. Jam. 137.

Habitat in India

Simillima A. cruciata sed triplo major. Specis plures quam quatuor,
et intra divituram ubi a pinicem distendunt dehiscunt, villosa.
Flores cum A. cruciata structura conveniunt Cui Am. l. c.

Eklaus Su

Cruciatula. Species quaternis cruciatis basi glabris, valvulis petaloides anthera
to

gromen dactylon bicorne minimum antheris longis armatum Sloan herb.

1. p. 112. t. 69. fig. 1.

Habitat in Jamaica

Chloris Sw

radicata & spicis (subpens) plurimis fasciculatis erectis pilis, flosculis sub
latis anfractis glabris ~~pub.~~ prodr. ^{sp. septibus} ~~pub.~~ monf. bot. 3.
gramen dactylon indicum spicis villosis subrepentibus
p. 105. f. p. t. 3. fig. 15.

~~Agrostis radicata~~
gramen dactylon majus, panicula ~~spica~~ ^{longa}, spicis plurimis
nervi ~~crassi~~ ^{longi} ~~secundum~~ ^{herb.} ~~1. p. 112. t. 69. fig. 2.~~

Andropogon fasciculatus spicis digitatis plurimis erectis
lis adsculatis lobis: flosculis utrinque anfractis her. 1.

V. pag. 904

gramen dactylon spicis gracilibus plerumque quatuor
coniciformiter dispositis Sloan her. 1. p. 111. t. 62. fig. 3.

Agrostis radicata: spicis subquinis imbricatis basi villosis, valva
lis petaloides anfractis her. Am. acad. 5. p. 392 + 111.

culmi inferne
gemulis inflexis

Habitat in India occidentali

Folia supra ^{glauca} pilis ^{raris} ^{acuta} ^{adpressa}. Spicula saepe ad 12. vel
2 inferiores parum remota reliqua umbellate sessile bipollicares
et ultra vulnulo calycino subulato antea longa.
variat foliis acutis vel obtusis vaginis compressis.

~~non ad~~

ciliata C. spicis digitatis subquinis; glumis corollinis margine ciliatis in
prodr. pag. 29

Habitat in Jamaica

gramen bipedale tenue. Folia longa convoluta, glabra margine
op. latis serrulata. Spica bipollicari. glomuli parvi gl. valvula
exterior calycis oblonga compressa carinata subconduplicata infra
apicem angustata basi canna et margine longe ciliata valvula
exterior subulata Corolla valvula altera truncata infra apicem an-
gustata

non vidi

Encicata Ch. spicis subquaternis encicatis, flosculis acuminatis & prodr
pag. 25

Gramen Dactylon bicorne minimum aristis longis armatum Poon
hid. p. 112. t. 69. fig. 1

Agrifolus Encicata spicis quaternis encicatis ^{basi} glabris, valvulis pete-
lodes aristatis Lin. S. V. pag. 111

Habitat in Jamaica

Culmi plures filiformes ^{inopleta} semipedales ^{ramosi} glabri inferne geniculati ^{tenuis} uno ^{glabro}
inferne uno altero infra. folia setacea. Spica 3-4 ^{semp} peripollinari flores
secundi aristis longo

— filis setaceis

polydactylon A. specis fasciculatis, petalis exterioribus aristatis; flosculi infimo
no ciliato-barbati Len. Am acad. 5. p. 412 x
Gramen dactylon elatius, specis plurimis tomentosiss. Sloan her. 1. p.
111. 4. 65. fig. 2.

Habitat in Jamaica

and synogon barbatum Len Am 5. pag. 412 x

Specis plurimis ^{5 me} aggregata terminalis, secunda, sessilis longitudine digi-
ti. Flores sessiles alterni. Calyx bivalvis, inaequali acuminatus, glaber
brevior biflorus. Flosculus sessilis prior. Glumina petaloidea bi-
valvis: Valvula exterior ciliata lana alba longitudine flosculi
apice terminata arista recta, longitudine flosculi. Flosculus
pedicellatus alter minor glumina corollae bivalvis. Valvula
exteriori imbricari aristata. Len

Folia convoluta
margine ore
centis ciliata

polydactyla Philoni specis plurimis fasciculatis laxis, glumina corollae margini
ciliato villosi Sw. prodr. pag. 26

~~novae~~

Philoni specis Swartzii

~~novae~~

sublot. diversa

Synonym Sloanii ad spec. Pappali a Swartzio amanda x

retroa @ spicis quaternis strictis erectis, spiculis confertis glabris
muticis, culmo compresso Sw. prodr. p. 25.
Habitat in Jamaica

Cynosuavis

~~Cynosuavis~~ conferenda cum Cynosuavis paspaloides contendit
quidem Swartzii eandem esse sed mihi videtur differ

pilosa ^{villosi} C. specii geminatis, foliis subulatis
Aegypti pilosa Petrii ?
Habitat in India orientali