

Picuta

1. *virosa* C. umbellis oppositifolius, petiolis marginatis, seminibus ovatis cista
tis multatis
2. *maculata* C. foliis serratis mucronatis, petiolis membrana ^{transversis} opus
bilobis, seminibus mucronatis angulatis
3. *bulbifera* C. ramis bulbiferis

Angelica lucida quid vide Off. no 132

Picuta nova Aubl. est Acrostichum filigosum Linn. est vero Aekoff. fil. Linn. vel Pteris
vel propriè genens

Lucida C.

Herb. mus.

Cauli crispè digiti, sulcatus, glaber. Folia bipinnata, pinnis ovatis, acu-
to serratis. Serraturis mucronatis, venosa, pellucida vel mucosa, rugosa
sepe biloba vel bipartita, glabra. Umbella latiuscula et vermicularis

perennans. P. foliis biternatis: folioli obtusi obliqui serratis fl. luteis d. p.
fructibus Walt. fl. Car. p. 116
Habitat in Carolina.

non vidi

bulbifera Cicuta ramis bulbifera her. S. V. 286.

umbellifera aquatica, foliis in minutissima et plane capellaria forma
ad idem Praei Suppl. 260.

Omni foliorum laciniis sp. capellanibus, caule angulato Gron. Virg.

^{31.}
Habitat in Virginia, Canada herb. 4. us.

Caule angulato. Foliorum laciniis capellariis.

Folia bipinnata: pinnulae lineari-lanceolatae, remote serratae, inferae
laciniis 3-4 terminalibus tripartitis. Ramuli axillares pollicares, oppositi
sub foliis, bulbiferi, oppositi vel alterni, vel superne alterni bipolli-
cares longiores. Bulbi alterni cum foliolo lineari, aequo summo. Um-
bello saepe radiato. Involucrum triphyllum: radii breves. Flores parvi

maculata ^Q foliorum serratis mucronatis, petioli membranacei apice bilobis
Lin. f. v.

Myrica Mitch. gen. 10.

Angelica virginiana, foliis acutioribus, semine striato minore, cumini
sapore et odore Monf. hist. 3. p. 201.

Angelica Canadensis, oleo trifolio, flore albo, seminibus luteis
striatis, cumini odore et sapore Plukh. alm 31. 7. 76. fig. 1.

Habitat in Virginia aquosa

caulis altus fusco purpurascens inferne uti et petioli
caulis altus fusco purpurascens inferne uti et petioli

Angelica canadensis
Genus folia ovata
Deli radice Tournef.
Inst. r. h.
herb Vaill.

maculata serratis glaber. Olea subtripennata
foliola opposita lanceolata ^{cuti emulata} inaequaliter serrata, glabra
foliola endardum opposita. Vagina unguicularis. Pedun-
culi axillares et terminalis involucrium parvum eva-
nescenti. semina ^{fibris} striata ^{oblongum} striata bisostriata
petioli basi utraque in membranam desinent

caulis
seminibus mucronatis bisostriatis

brevispinis

virga & umbellis oppositifolius, petiolis marginatis obtusis Lin. S. V. Mant. 355.
Fl. Dan. T. 200.

Sum enico folio Bauh. pin. 154.

Habitat in Europa paludibus sterilibus &
Caulis teres fistulosus glaber striatus. Folia ampla bipinna-
ta glabra, pennis lanceolatis utrinque acutis, serratis
nisi laticoribus quam in maculata pubes pallida
radii umbellae basi purpureo. vagina sesqui pollicia
res margine membranacea. Eruentis compressis
truncis

seminibus ovatis costatis muticis