

Census

Specimen ex Am. boreali det. Ruff.

Cissus

Cissus mucronata foliis ternatis oblongis serratis glabris membranaceis
Vitis Infolia, una corymbosa, acinis mucronibus rotundis Plum. Cat. pag. 18.

Icones Tab. 259. fig. 4.

Habitat in India occidentali non Rahrz

Rami angulati, glabri.

Folia petiolata, ternata. Foliola lateralia sesipha, bipollicaria, lanceolata,
elliptica, obtusifida, mucrone brevi terminata: intermedium petio
latum, subtripollicare, ellipticum, interdum utrinque attenuatum
lateralibus duplo latius. Omnia serrata: serraturae remotae d. frontibus

Cissus oppositifolia
liis

Petiolus sesquipollicaris, utrinque membrana marginatus.

Cyma pedunculata, axillaris, quadriradiata: radii inaequales, unguem
laras, divaricati, pilis raris adpressis, bis bifidi. Pedicelli ultimi ~~trifidi~~

basi foliola plura
munita, subulata

undecim vel tridecim, breves.
Flores minuti, dichotomia pedicellati.

Litr. C. pag. 17.

Calyx marginis brevis, subquadrilobus.

Petalae quatuor, oblongae.

Stamina quatuor, longitudine corollae, neutra infesta

bas. margine nectonifero cinctum.

Germen ovatum, glabrum, Stylus longitudine corollae

Bacca parva, oblonga obtusa

(Rami quidem angulati, ad non e membrana decurrente ut in Risso trifoliato,
nec folia bifida subdentata

① glandulosus Cuspis foliis ovatis serrato-dentatis, pedicellis calycibusque hispida glandu-
loso
Cuspis glandulosus; foliis serrato-dentatis; pedicellis et calycis basi glandu-
loso Boiss. descr. p. 34.

Habitat in Arabia

. Caules teretes. Folia opposita, sequipollicaria, ovata acuta, serrato-
dentata, carnea. Stipula deflexo carnofo lanicolata, margi-
ne membranacea. Curvis interfoliaceis. Pedunculi cymosi Boiss.

non vidi

dissectus Cypus foliis quinatis oblongis obtuse serratis subtus tomentosis

Habitat in Guinea Prod

Herbaceus. Ramis dichotomis, dissectis, pubescentibus, pedibus cinereis, verticillatis.
Folia petiolata, alterna: foliola petiolata, intermedium bipollicare, lateralia
minora, basi attenuata, apice obtuse serrata, serraturis mucronatis
^{minutis} terminatis. Cirsium oppositifolium. Pedunculi axillares, foliis multo
longiores, tomentosi, primum trifidi, demum bifidi, reliqui racemosi; pedicelli
2-3. ex eodem puncto. Calyces parvi unani, parvi

3. latifolia Cuscuta foliis cordatis acuminatis fetaceo serratis

Cuscuta latifolia Cas Marti.

Schunambra - valli Rheod. T. 7. tab. 11

Caulis albido farinosus, articulis nodosis. Folia glabra. Peduncululi dichotomi
umbellulis ut in congenitis

asicyoides ~~st.~~ - sicut ut foliis ex parte cordatis, acumine acuto

a tomentosa nostra non diversa

1 vitiginea Cuspi foliis cordatis sublobatis repandis tomentosis.

Arbutula barre fera circum plicabilis, vitiginea folio subtus lanata, fructu
raemglo feravidio odore Plukh. mant. 27. t. 337. f. 2.

Cuspi flora Rayl. 60* Am. Acad 1. p. 390*

Cuspi foliis cordatis subquinquelobis tomentosis L. S. V. pag. 158.

Habitat in India Orientali ^{et} ~~per uos~~ ^{particulat}, ~~consp~~

Ramuli teretes, tomentosi, et de caespitibus uti in ~~aliqua planta~~.

Folia cordata, coriacea, nervosa, repanda, et sublobata, obtusissima, latera fere
quam longa. Pedunculis ~~et~~ stipulis ovales membranaceae, acuto oppo-
sita semiunguiculares deciduae. Pedunculi oppositi folii, dichotomi
fructus pyriformis

+ Viti Madagascariensis, fructu aereis, folio subrotundo et anguloso Pet. v.
mus. n. 696.

mappa Cuspi

Habitat

herba Martii

Bolea petiolata, opposita, pollicaria, oblonga, glabra, corymbosa, subaequali
minata, obtusa. Stipula lanceolata. Cyma trifida, dichotoma.

Dipolov Cusur

Habitat in Andra Orientali herb. Fus.

Raulis angulatus, ramosus. Folia petalata, Tripolliciana, ovato-oblonga, vel subcordato-oblonga, attenuata, ^{in pediculis} glabra, tenera, remote setacea serrata, vix venosa, subsque nervia, subtus rufescentia, obtusa. Corolla oppositifolia. Sepala ovata, obtusa. Pedunculi oppositifolia, semipollicaria, pediculi oppositi. Pedicelli umbellati, plures, glabri

micranthus Cusis

Habitat in Domingo herb. Desfont. 5

Rami glabri, inferne teretes, punctato scabri, superne angulati, laces. Soli
petiolata, semi polliciana, ovata, 3-5 loba, angulata. Lobus intermedius lon-
gior, serratus 1-2, glabra. Petioli longitudine folii. Pedunculi axillares, solita-
rii, petiolo brevioris. Umbella composita, radii 3. parthales 5-6. Flores
parvi

palmata Cuspi

Habitat in Madagascar Commerc. herb. Cuspi.
Rami teretes, glabri. ^{Sp. pollicibus} Dolca petiolata: inferna s. fida: lobi oblongi, glabri, venosi,
remote setaceo ferrati, ferraturis acutis mucronati: suprema cordata, in-
divisa, densius ferrata. Pinnis oppositis: foliis. Stipula oblonga, obtusa.

Cuscut.

Habitat in Borana Commes. herb. fus.

Rami glabri, angulati. Folia petiolata, distantia, 5nato vel 7nato digitata,
foliula simplicia, bipoll. et una, ex venis minoribus, canaliculata, ex trospum
remota serrata, serraturis subulatis, utraque attenuata, acuta, glaberrima.
Reritur oppositifolius. Flores non vidi.

brasiliensis Cuspus

Habitat in Brasilia herb. Tuss.

Rami teretes, articulati, villosi. Folia petiolata, bipollicaria, oblonga, attenuata, remote setoso serrata, binerviata, nervoso-venosa, basi versus intermedia, villosa, supra minus. Umbellae oppositifoliae, compositae, quinque radiatae secundariae 7 radiatae. Pedicelli villosi.

Cyclamenifolia Cofus folio cordatis remotissime seduceo subserratis, umbellis compositis

sitis folio cyclamini uva corymbosa, acinis nigricantibus Plum.

Habitat in America herb. Suriani

Tota glabra. Caulis flexuosus, superne angulatus. Folia petiolata, tripollicaria, acuminata, serraturis circiter 8 utrinque, minutis, subvenosis, glaberrima membranacea. Petoli folio dimidio breviores. Pedunculi axillares, sesqui polliares, radii umbello univesales 5-6, femi polliares; apice interdum bifidi; pedicelli plerumque sex, femi unguiculares. Flores parvi.

larescens Cusur

Habitat.

hab. Thauri

Rami tomentosi, incani, uti folia subtus. Folia sesqui pollicana, subcordato-ovata
obtusiuscula, remote serrata, supra oculo armato adpressa vilius minutius. Pedun-
culi oppositi folii. Umbella decomposita, incano. Circhi simpliciter

biflora Passiflora

Habitat

Herb. Koenigii

Cuscuta strata foliis digitatis; foliolis oblongo lanceolatis serratis, caule strato
flexuoso, cymis coarctatis fl. Penn. 1. pag. 64. tab. 100. fig. b.

Habitat in fylis Chili

Caulis fruticosus, scandens, ramosissimus; stratus. Rami pubescentes.
Folia aeterna, petiolata: foliola 3-4. saepius 5. glabra, superne serrata
per ramosi acuminatis, ^{lateralibus} ~~terminalibus~~ minoribus. Petioli pubescentes. Folia
breviora. Stipula ovata, acuta, femina plerumque obliqua. Cymis
oppositi folii bifurcati. Cyma oppositi folia et terminalis, dichotoma
multiflora, filorum longitudine. Pedunculi et pedicelli pubescentes.
bractea ovata minuta, acuta, concava. Petala lutea fl. Penn.

90

Charact. Cusi emendat. ind. loco titato vix a Linnæi diversis nisi
quod Bractea biloculari, bi-4 sperma. ^{filis, acuminatis} semina umbilicata rugosa

Cistus obliqua foliis binatis, ternatisque serratis, lateralibus cordatis obliquis
intermedio ovali, floribus umbellatis corymbose fl. Peruv. 1. pag. 69. Tab.
101. fig. 6.

Habitat in praeceptis Huancu et Tarma

Caulis scandens, ramosus, tetragonus, striatus. Rami pubescentes. Folia
aeterna, longe petiolata, binata vel ternata, acute serrata, venosa,
supra pilosa, lateralia cordata, basi obliqua altero latere angustiora
Stipulae oppositae, ovatae. Pedunculi oppositifolii, longitudine fere
foliorum. Cyma 3-4 radiata, involucriata. Involucrium 3-4 phyllum, ovatum
Pedunculi dichotomi, inaequales. Corolla lutea fl. Peruv.

Cuscuta compressicaulis foliis cordatis acute serratis, floribus umbellatis
pedunculis dichotomis fl. Peruv. t. pag. 64. Tab. 100. fig. a.

Habitat in ruderalis Provincia Chanay.

Pubescens. Cauli fraxidens, ramosus, tetragonus, compressus, geniculatus
repens procumbens radicans. Folia alterna, petiolata, obtusa vel acuta
venosisima, supra nitidiuscula, nervella subtuloba. Petiol. teretes,
lenitoli. Stipulae oppositae triangulari, marcescentes. Sepala simplicia
ad medium squamae duo opposita, lineares. Pedunculi petiolo longi
res oppositi folii. Cyma 4-7 radiata. Pedunculi dichotomi, bracteis 2.
serratis. Flores 2 dichotomia, solitarii. Petala lutea. Bacca subrotunda
diperma, purpureo-nigra fl. Peruv.

Cissus granulosa caule angulato granuloso, foliis quinatis, foliolis obovatis
dentatis, cymis patentibus fl. Peruv. 1. pag. 64. Tab 101. fig. a.

Habitat in praeceptis Huasbahuasi

Globa. Caulis scandens, ramosus ^{longis} cymis, tenax Vitruginosus. Folii alterna,
petiolata: foliola extus ^{longis} dentata, lateralia minora. Petioli foliis
duplo ^{breves} ~~longiores~~, granulosi. Costae bifurcatae, striatae. Cymae oppositifoliae
terminaliaesque, dichotomae. Corolla lutea. Baccae nigrae, magnitudine
cucurbitae fl. Peruv.

canescens Cuspid. folius cordati oblongi serratis subtus ramisque tomentosis

Habitat in umbrosis Peru. Dombey 4

Rami, et folia subtus raris petioli et pedunculis nudi. Folia bipollicaria
sa, remote serrata, serraturis acutis, eglandulosis, crassiuscula venosa, state sub
tus glabra. Stipula oblonga, parva. Cinchi oppositi folii, pedunculi oppositi
folii: Umbella subdecomposita: nonnisi quinque radiata: secundaria 2 radia
ta, cum pedicello uniflora in centro

non compressis in aulis fl. Peru

Cannabina Puffo.

Habitat in monte vides herb. Tusf.

Rami angulati, punctati, glabri. Folia peti. ovata, 5-6 digitata: foliola
lespica, lanceolata, tripollicaria, remote serrata, basi integerrima, serr
turi seta mucronatis, obtusa, mucronata, venosa, glabra.

Latifolia Cusper.

Habitat herb-*ha* Martini.

Rami glabri. Folia palmata et ultra, cordata, attenuata, serrata, serraturis mucronatis, venosa, vena subtus bilobata, supra oculo armato adpressa vellis obsoletis. Coeternum glabra.

obtusifolia Cusps.

Habitat herb. la Mantini

Folia ternata: foliola semipollicaria obovata, v. illopa, serrata. Cyma trifida

Cuscuta

Habitat Capeno herb. la Mantia &

Caulei tetragonus, fruticofus, glaber. Folia simplicia, opposita.

pelosa Cuscuta foliis Vernatis: foliulis obovatis dentatis glabris, caulis
et cymis que glandulosis pilosis
Habitat in India orientali Rottler
folia ad spem Angli rarissima

~~foliata rugosa foliis subseptem, foliis oblongis utraque
ae uncinatis serratis; terminali spili: fructibus ellip-
ticis quadrisulcatis Wild. Ber. Baumg. pag. 154~~

Punctilobus *Cuscuta* *varius* *puncticulosis*, foliis ^{lato-} cordatis ob
tusis, aluminè abrupto obtuso, semiremotis lobis
taceis. Act. Soc. h. N. Par. 1. pag 106
Habitat in Am. merid.

michi nota

ovata Cuspis foliis ovatis sensim et obtuse acuminatis gla
bri. Ceviter petalis serratis Act. f. h. N. Paris. 1. pag. 106
Habitat in Amer. merid.

La Mart.

erosa Rufus glaberrima Infoliata petiolo submarginato; folioli
oblongo ovatis basi subacutis obtusius erososubdentatis mem-
branaceis cyma longe pedunculata, subpaniculata Aot.
foc: par: 1. pag. 106
Habitat in Amer. Merid.

10 acida *Cissus* foliis ternatis obovatis glabris carnosis incis^{is} L. S. V. pag. 158. ^{Swartz obs. pag. 497}
Bryonia ^{renata} *trifoliatum* indicum, foliis succulentis crassis et ~~crassis~~ ^{renatis} Pluk. alm. 76. tab. 152.
fig. 2.
Bryonia alba triflylla geniculata, foliis crassis acris Sloan. h. j. 1. p. 233. tab. 142 f. 6.
Vitis trifolia minor corymbosa, acinis nigronibus turbidatis Plum. fr. tab. 299 f. 5.
Colum cauponis Rumph. Amb. 5. p. 458. t. 166. fig.
Triplo triflylla scandens et claviculata, foliis crassis serratis Brown. jam. 1. p. 147.
Habitat in Jamaica, insula St. Lucie et India orientali
Caulis lignosus alte scandens. Folia opposita, ~~alt~~ floralia alterna, petiolata, ternata
foliata ~~obovata~~ obovata, glabra dentata, incis^a, carnos^a, intermedium petiolatur.
Peduncul^{is} ⁵ ~~radicata~~ ^{radicata} oppositi folii, tripollicares. Flores bis umbellati ~~racematis~~ ^{racematis} ovatis acutis
ad basin pedunculorum partialium et pedicellorum. umbellule interdum iterum diviso.
radix univesalium et partialium quinque

Roigii Cuscuta foliis ovatis subcordatis attenuatis setaceo serratis glabris, caule
tereti umbellis triadatis

Habitat in nemorosis India Orientali

Facies C. siyodese. Caulis articulatus glaber. Folia petiolata opposita

basifere transverse glaucos vix simpliciter nervosa, acumina
obtusis remote setaceo serrata. Pedunculi ^{petiolo dimidio breviores} oppositi foliis radii
3 longitudine pedunculi pedicelli 5-8 umbellati

Cuscuta

Columella pedata foliis pedatis, racemis planis dichotomis

Leucis fl. Cochlea pag. 86.

Habitat in Cochlearia 5

fructus pendens & res, terrore folia petiolata. ovata,
crenata tomentosa & racemi laterales Calyx sub
truncatus Petala 4. nectarium 4-filiatum. Stamina masculi
ne calycis infundibulo. Stigma simplex bacca subrotun
da, 2 loculis appensa

non vid.

Cissus

Cissus rhombifolia, pubescens, foliis ternatis serrulatis: intermediis rhombeo
ovato, lateralibus semiorbati

Habitat in insula Trinitatis von Pohl.

Rami angulati, uti tota planta villosi.

Folia ternata: folia ^{lateralia} sessilia, bipollicaria, latere exteriori triplo
latiora: intermedium petiolatum, tripollicare, ovato
rhombicum, utrinque ^{attenuatior} ~~acutius~~ ^{acutum}, omnia ret venosa, tenuissi-
me nervosa.

pubescens B. foliis conjugatis cinctis: foliis cordato-ovatis subtus pu-
bescentibus Lin LV Mill. Dict. 14.

Habitat in Campeche &
Coastal

Specimen Beynonian misit Dr. Von Rohr et St. Martha 1021.

vellosa

quo huius proxima non eadem certe statuere nequeo.
Rami teres grisei, superne compressi, paleis tomento lanugineo terti.
Folia petiolata opposita conjugata foliolo petiolato ovata
membranacea venosa supra fere glabra nisi in et circa costam
villis minutis subtus & villis griseis mollibus, integerrima, byssis
caulis petioli communes pollicares, partiales femurquic alares
nervi acuminati
uti communes grisei, pubescentes terti. Cirrhii in apice petioli
acuti
filiformis grisei. Calyx campanulatus glaber truncatus.
Corolla magna laxe 5-pollicari nupuracea glabra
nervi lutei

non vidi

umbellata Cuspi foliis ovatis integerrimis, floribus umbellatis Lewis. fl.

Certh. pag. 84

Habitat Perhincina 5

~~Caudis quadrangulans perenni longis ramis glaber virens,
articulatus scandens, crispis, sulcatis. 5~~

Caulis fruticosus, volubilis longis, ramis. folia opposita
glabra. umbellae terminales. Corolla alba campanulata
intus lobata. Baccis subrotunda

collis intus lanatis

orientalis Cusper folius subbipinnatis foliolis ovatis serratis, caula frute
siente La Mark T. En. pag 332. tab. 84. fig 2.

Habitat in oriente 7

Fraxus vitis arborescens at major folius minus compositis foliolis latiori
bus

connuens - Cus. folius subbipennatis, foliolis ovatis obtusiusculis sub
dentatis, petalis connuentibus La Mart. T. Enc. pag 332.
Habitat in Madagascar.

Mappa Cuspi folus bipinnatis lobus folioli ovatis integerrimis
La Mart. T. En. pag 332.
Habitat in sapula francis

non vid.

obtusifolia Cuspi: folus ternatis; foliolis obovatis obtusi dentatis
pubescentibus La Mart. T. Eni: pag. 332.
Habitat in Andra orientali

non vid.

Cinerea C. S. f. foliis ternatis folioli pubescentibus dentatis lateralibus
subcordatis, petioli teretibus. Lamour. T. Enc. pag. 332.
Habitat in Ind. orientali.

non vid.

Canepiensis Ruspis foliis ovato-oblongis, obliquis denticulatis subtomentosis
Canepiensibus La Mart. F. K. pag 331
Habitat in Peru

non vid.

ovata Cuspus foliis ovatis acumineatis venter dentatis utrinque lobatis
Lam. Mart. T. E. pag. 331.

Habitat in Guadeloupe

Cuspus Brown. T. 4. pag. 1. 2.

angulata Cuf. foliis ~~sub~~ subpentagonis angulato lobatis venulatis pubes-
tomentosis La Martk. T. Enc. pag 331.
Habitu in Andra orientali

tomentosa Cus. f. foliis subpentagonis, obtuse dentatis subtus tomentoso ferrugineis
nisi La Mart. T. E. pag. 330
Habitat in insula Bourbonica
Folia basi subtruncata

Sw. obs. pag. 407

5 Sicyoides *Cuscuta* ^{sub} *foliis cordatis* ^{ovatis} *nudis, setaceo serratis ramulis teretibus* L.S.V. pag. 15. *Tripl. Ar. pl. 16*
Byronia alba genicula lato, *violae foliis*, *baecis e viridi purpurascantibus* Sloan h. t. pag
 + 223. tab. 144. fig. 1.
Fosida scandens, *foliis oblongo ovatis ad margines denticulis setaceis* Brown jam. p.
 147. tab. 4. fig. 1. 2.

Cuscuta foliis simplicibus nitidis Jacq. amer. 22. tab. 15.
 + *Vitis foliis dentatis* Plum. fr. 259. fig. 2.

Habitat in Jamaica Cambai
 Caulis lignosi altissime scandentes. *Folia cordata* *petiolata*, *alterna*, *cordata*
vel ovata, *acuminata*, *obtusiuscula*, *glabra*, *exteriorum* ^{vamata} *setaceo serrata*,
sesquipollicana et ultra. *Umbello* *oppositifoliae* *longitudine petioli* ^{3 vel} *quon-*
que radiati, *parthali bifido quinqueflori.*

varietas *foliis* *latioribus* *fructis* *cord. folia* *at foliis* *setaceo serratis*
nei in Tegernsee

umbellis sive radiatis

13 *Digitatus* *Cisus* ~~fol.~~ foliis quinatis ternatisque oblongis serratis, ramis ^{glabris} verticillis
~~foliis~~ *Palanthi* *Digitatus*; foliis inferioribus palmatis quinatis, superioribus
ternatis; omnibus petiolo communi sustentatis *Boiss. & Ehrh. pag.*
35. t. tab. 3.

Habitat in Arabia

Folia inferiora quinata, folioli petiolati, oblongi serrati glabri
uti totus fructus globosus verrucosus

17 trifoliata Cesp. foliis ternatis subrotundis hispitis subdentatis ramis membranaceis angulatis
L. V. pag. 159. Jacq. Amer. p. 23. tab. 182. fig. 10. pl. sel. pag. 20 * Swartz obs. pag. 50

Bryonia alba triphylla maxima Leon. Jam. 1. p. 233. tab. 144. fig. 2.

Gofiola triphylla scandens; foliis ovatis subdentatis petiolo communis marginatis,
calycibus majoribus Brobn. Jam. 147.

~~Calyx~~ Habitat in Jamaica, Cathayno.

Caules lignosi, cernosi, scandentes; rami petiolique membranacei angulati, pilosi.

Folia alterna, ternata: foliola acuta, inaequaliter serrata, hispita,
venosa nervosa, lateralia semicordata, medium ovatum. Umbello ut in
Cesp. fig. 2. at paulo majore. Jacq. l. c.

9 ^{lespilifolia}
~~Lespilifolia~~ ^{lespilifolia} Cuscuta foliis ternatis ~~(foliis ovato-cordatis serratis)~~
Solanum ternatis, foliis ternatis, complicatis, serrato-dentatis. Boiss. Desf.
pag. 35.

Habitat in arabia

Caules scandentes, crassitie pennae cygnae, teretes, albido-pulicatis, glabris, articu-
latis. Folia ternata, sessilia, ovata, acuta, basi cordata, grosse serrata.
Stipula gemina patenti recurva, ovata, acuta, semipollicaris glabra.
Culmi oppositifolii, bipedi. Boiss.

7 rotundifolius Buple foliis ser cordato subrotundis serratis glabris Synt. bot
Solanthus rotundifolius; foliis cordato-orbiculatis, serratis Boiss. Desf. pag
35. Gen. Tab. 4.

Habitat in Arabia

Caulis teretes articulati, geniculis basi incrassatis. Folia subrotunda, cuneo
obtusiuscula, glaberrima, subavenia, remote serrulata. Pediculi dichotomi

umbellula ~~radiata~~ racemosa, opposita multiflora 40 circiter
in singulis nudunculis ~~comuni~~ pedunculis ~~compositis~~ ^{collegatis} ~~foliolis~~
~~cordifolia~~ differt foliis ~~ser orbiculatis~~ cordatis acutis integerrimis

6 quadrangulans Cuscu foliis cordatis carnosis serrato-dentatis caule tetragono humidum
Sculo Lin. Mant. 39 * L. V. pag. 150

Planta base. fera scandens epidendros Planch. phyt. 310. f. 6.

Tunis quadrangulans Rumph. amb. 5. p. 83. v. 44. f. 2.

Plantae quadrangulans; caule articulato, dichotomo, tetragono; foliis
carnosis cordatis, serratis semicomplacatis G. Hoffm. de stir. pag. 33. t. c. tab. 2.

Habitat in India, Arabia.

Caules dichotomi, tetragoni, ^{corticati} in rascati, carnosus, geniculis contractis, laevibus.

Folia alterna, ^{sub} carnosus, cordata, subhastata, utrinque glabra, angustis
† Petioli angulati, ^{se} remoteque serrata. Peduncululi uti cuncti oppositi foliis bi vel ter
umbellati Lin. G. Hoffm. t. c.

2. cordifolia Cespis foliis cordatis integerrimis L. S. V. pag. 158.
Sicut foliis subrotundo, utra corymbosa conulea Plum. 4c. 259. f. 3.
Habitat in America.

10 pedata Cyperus foliis quercata pedatis: foliolis lanceolatis serratis subtus tomento
sic symb. bot.

Belutta - Tjori - valli Rhed. Mal. 7. Tab. pag. 19. Tab. 10

4 Cyperus pedata La Mark. Encycl.

Habitat in India orientali

Lambucus cana Rami tomentosi incani fr. vel 4 pollicaria
densus Burman. scolia pedolata, pedata, s. 6-7: foliola petiolata, lanceolata, incana
And. pag. 75 qualia, lanceolata, attenuata inaequaliter serrata, acumine
propter Synon. indegerrimo, supra vellosiuscula, subtus vel villosa - tomentosa
Rhedii incana, seriosa.

capitulum tomentosum Petiole tomentosi incani - petioli bifidi: partiales laterales ramosi
hub. La Mark. intermediis simpl. indivisus folia monophyllis

Cyma dichotoma s. sexies divisa, incana. Flores parvi

Calyx obscure quadridentatus, minutus

Corolla petala quatuor

Nectarium ad marginem germen cingens

Stigma acutum

Obovata Cuscus foliis ternatis ~~obovatis~~ foliolis obovatis integerrimis glabris
In insula St. Crucis legit Dr. Rector ^{Wet,} qui cum multis aliis plantis ame-
ricanis ~~benevole~~ ab ipso in America detectis benevole com-
municavit

Caulis scandens crotchi, glaber. Orthus oppositi foliis, longitudinis folii, bifidus
Folia petiolata, alterna, ternata: foliola obovata, mucronata,
glaberrima, membranacea, integerrima, latera ^{laci} minoria, sessilia
intermedium petiolatum triplo majus.

Pedunculi axillares, solitarii, foliis longiores, glabri, trichotomi, ramu-
li triflori: floribus pedicellatis.

gustus acerrimo Ruck.

relegiosa P. foliis lan petalisque lanceolato ellipticis

utroque acutis

Nürvala Rheed. Mal. 3. p. 49. Tab. 42.

Praterea relegiosa inermis, foliis subaequalibus

Koepf. prodr. pag. 35. pl. excul. pag. 45. & prodr. n. 203

Habitat in India orientali et Societatis Anglorum

Prami. Venter cortice olivaceo, punctis elevatis
albidis adspersus. Folia petiolata parva ad

apicem ramulorum. foliola terna petiolata

attenuata integerrima glabra, tenues nervos
Spithamou. lateralibus parum minus margine

altero angustiore. Cyma Pedunculi ad apicem
alterni tripollicares, axillares, interdum cy-

mosi, uniflori sexies. petala sequipolliciana

calyx ungue petalo quadruplo brevior. Perianthium

un quadrupartitum laciniis lanceolatis

ungue utantibus. Germen oblongum

7 crenata

Populus foliis ternatis; foliolo subrotundis crenatis

Folium Causonis Rumph. 5. pag. 450. tab. 166. fig. 2. et folia nimis
ovata et acuta

Vitis trifolia: foliis ternatis, foliolo subrotundis ternatis Lin l. V. pag. 244
Rami villosi uti petioli folia juniora et pedunculi.

Solida Folia petiolata ternata: foliolo petiolata, pollicana Cameraria
parva minima, altero latere angustiora, crenata crenis remotis
microratis

Stipula parva oblonga obtusa.

Pedunculi folia longiora Cyma dichotoma. Pedunculi folio longio
ri.

Calyx minutis quadridentatus

Petala quatuor fornicata

Nectarium utraque in congenentibus inter stamina et germen.

Filamenta quatuor corolla breviora

3 Carnosa Cuscuta Cuscuta foliis ternatis ovatis ^{obtusis} seratis glabri, ramis teretibus
prostratisque teretibus

non valli Rhed. Malab. 7. tab. 9.
uni crepitans Rumph. amb. 5. pag. tab. 165

Cuscuta carnosa La Mark. Enc.

Habitat in India orientali

Rami, sicuti tota planta, glabri, striati, teretes.

Folia petiolata, ternata: Foliola petiolata, ovata, carnosa,
obtusis, serrata, lateralibus minoribus, pollicis aequa.

Petiolus communis longitudine foliorum.

Fl. pedunculata folio longior, ramosa, fl. umbellata.

Humbella radii
terni, ramis
dichotomis
Pedunculis universali
folio longior

Differt a C. trifoliata, glabritate vellosetate, folioli lateralibus
obtusis, ramis petiolisque, mem. bracteolatis
a C. acida dignoscitur foliis serratis nec unguis obovatis unguis

4. *Palaefolia* Buddlejia foliis angusto lanceolatis ~~caetero~~ ^{caetero} ~~superne~~ serratis, paniculis racemosis.

Habitat in Bonaria 7

Rami ~~offici~~ ^{rami} teretes, ~~teretibus~~ ^{teretibus} tomentosis ~~iniani~~. ^{foliis opposit} folia ~~opposita~~ ^{opposita}, angusta lanceolata, elongata, ^{triplo die aerea ex utroque latere} a medio ~~exteriorum~~ ^{exteriorum} ~~versus~~ ^{versus} obtuse serrata, inferne ~~integerrima~~ ^{integerrima}, margine ~~paucum~~ ^{paucum} reflexa, supra glabra, subtus ~~tomentosa~~ ^{tomentosa}. ~~venis~~ ^{venis} ~~glabris~~ ^{glabris} ~~racemis~~ ^{racemis} ~~terminalis~~ ^{terminalis}, ~~spithameis~~ ^{spithameis}, ~~foliosis~~ ^{foliosis}. ~~Paniculi~~ ^{Paniculi} ~~oppositi~~ ^{oppositi} ~~florae~~ ^{florae} ~~paniculati~~ ^{paniculati} ~~pedunculi~~ ^{pedunculi} ~~oppositi~~ ^{oppositi} ~~pedicelli~~ ^{pedicelli} ~~ramosi~~ ^{ramosi} ~~paniculati~~ ^{paniculati}. ~~foliola~~ ^{foliola} ~~florales~~ ^{florales} ~~foliolum~~ ^{foliolum} ~~stipitacum~~ ^{stipitacum} ~~ad basin~~ ^{ad basin} ~~pedicellorum~~ ^{pedicellorum}. ~~Floris~~ ^{Floris} ~~carneis~~ ^{carneis}. ~~Calyx~~ ^{Calyx} ~~et~~ ^{et} ~~corolla~~ ^{corolla} ~~exsert~~ ^{exsert} ~~tomentosa~~ ^{tomentosa}. ~~laciniis~~ ^{laciniis} ~~calycis~~ ^{calycis} ~~ovato~~ ^{ovato} ~~obtusis~~ ^{obtusis}. ~~limbus~~ ^{limbus} ~~quam~~ ^{quam} ~~nifidus~~ ^{nifidus}; ~~laciniis~~ ^{laciniis} ~~rotundatis~~ ^{rotundatis} ~~et~~ ^{et} ~~glabris~~ ^{glabris}. ~~Stylus~~ ^{Stylus} ~~longitudinem~~ ^{longitudinem} ~~corollae~~ ^{corollae} ~~stigma~~ ^{stigma} ~~obcordatum~~ ^{obcordatum}, ~~tunicatum~~ ^{tunicatum}.

Pisus

virginica Pisus foliis cordato-subrotundis lobatis subinde usque villosis dentatis

Pisus Flor. Len. Flor. reyl. 60.

Pisus virginica: foliis cordatis subquinquelobis tomentosis Len. Spec. pl.
pag. 170. exclusis syn: proter Plukenetii

Rami villosi, subtetragonis superne incani, subtetragonis.

Folia petiolata, alterna, bipollicaria, integra, ^{caeterum versus apicem} superne f. interdum
subangulata, tri ^{loba} vel subquinqueloba, neriosa, pubes villosa

crenata obtusis dentata: lobus intermedius longior major. pinna
petioli angustissimi. Stipula parva, ovata, membranacea, decidua

Umbella pedunculata, oppositifolia, triradiata; radix dichotome ramosi
divaricati. Floris incani

Fructus pyriformis, magnitudinis pisii, glauco coeruleo semiter

vitiginea Cusis foliis cordato-subrotundis lobatis subintegrisque villosis dentatis

~~Cusis foliis cordato-subrotundis lobatis subintegrisque villosis dentatis~~

~~Cusis foliis cordato-subrotundis lobatis subintegrisque villosis dentatis~~

1-2 Latifolia Cuscut foliis cordato-ovatis villosi acuminatis seta longioribus, ramis tetra-

gonis

Schinambu-valli Rhed. mal. 7. pag. 21. tab. 11.

Tunis crepitans major Rumph. amb. 5. pag. 446. tab. 164. fig. 1.

Cuscuta latifolia La Mart. Encyclop.

Habitat in India orientali &

Rami articulati, lignosi, uti ~~folia~~ ^{petioli et} pedunculi, involucri ferrugineo

tomentosi; ~~stata~~ ^{stata} propositum superne, inferne stata glabra

Folia petiolata, alterna bis. tri-pollicaria, ^{superne} cordata, ^{plenumque} fere basi ~~subtus~~

^{basibus} basi oblique ^{sub} ~~versis~~ ^{versis} absque sinu, adalteria supra glabra, subtus, uti juniora villi

ferruginei proferunt in nervis, ~~tomentosa~~ ^{sub} subtomentosa ^{ad} cum veno

sa, nervosa: nervis oppositis, acuminata, serrata: serraturis seta

^{cei} Petioli semi-pollicari. Stipula oblonga, laterales

umbellae composita, pedunculata, ^{foliis brevioribus} quinquemurata: partiales

Planis ⁱⁿ caudice octo provolucrium nullum nisi villi ferruginei ad

Basice Trifolium ⁱⁿ pinniformis, glaber

12 repanda Cuscuta foliis cordatis integris sublobatisve utrinque glabris ^{repandis}

Habitat in India orientali

Rami teretes, flexuosi articulati, tomentosi, stulte glabri

folia pedicelata, ~~cordata~~ bi-^{stri}trilobata, latiora quam longa, cordata ^{obtusolobata lobata}
firmiora quam in congenitibus, integra vel ^{obtusolobata lobata} ~~obtusolobata lobata~~,
nervosa, adultiora utrinque glabra, juniora villosa, repanda; crenis
mucrona minuto terminatis, obtusissima, interdum acumine brevi
obtusis terminata

Stipulae ovales, membranaceae, acutae, oppositae, semialternatae

Pediculus oppositifolius; radii umbellati tri; dichotome ramosi.
pedicelli ultimi umbellati. Squamae villosae, brevissime ad basin
sui pedicellorum

Stylus ovario Bacca pyriformis, magnitudine pisii, stylo persistente
mucronata

- Cuspi obliqua
1. Cuspi obtusifolia, folia ternata villosa, foliolis ovalibus, obtusis, crenatis
Ind. Onent. Sonnerat.
 2. — trilobata foliis ternatis tripartitis (vel ternatis) ovatis serratis ^{diffinitio mea} Pheed.
malab. T. 7. pag. 85. tab. 45. Viareta. - Tjoni - valli
 3. — repens foliis cordatis acutis pedunculis pedicellisque umbellatis
Si Henam - pulli Hort. Mal. T. 7. p. 91. tab. 40
 4. — Schembra - valli Hort. Mal. p. 11. Tom. 7. T. tab. 6. folia cordata flores sabrai mosi
 5. — Wallia - pura - pitica ib. pag. 13. tab. 7. folia triloba dentata
 6. — Wallia - Tjoni - valli ib. p. 15. tab. 8. folia, quaterna, ternata, simplicia
lanicollata, remote serrata
 7. — Balutty, Tjoni - valli ib. p. 19. t. 10. Cuspi pedata Ca. marthii flamma
quatuor

} La Martia
 } Encycl.