

Cladonidium

1. infortunatum C. foliis cordatis attenuatis integris subtis villosi
elentico
2. fortunatum C. foliis lanceolatis integerrimis
3. phelonioides C. foliis ovatis integris dentatis angulatis que, pedunculis
axillaribus subtis flori
4. calamitiforme C. foliis ovalibus subdentatis nudis, panicula bra-
chiata.
5. paniculatum C. foliis quinquelobis denticulatis glabris, panicula
brachicata
6. squamatum C. foliis cordatis obscure angulatis, panicula ramis
dichotomis
7. trichotomum C. foliis lobatis indivisis que ovatis, panicula
ramis dichotomis glabris
8. diversifolium C. foliis integris trilobis que ovatis, ramis dichotomis
villosis, pedicellis ultimis rademosis

Cleodendron feriatum Less. 17 anthe
pyramidata
deuminata

capitatum Alerodendron foliis ovato-oblongis integerrimis, floribus ter-
minalibus capitatis

Wolhamera capitata; foliis ovatis integerrimis scabris, floribus
terminalibus capitatis, calycis foliaceo. Wald. Spec. III. 1. p. 384

Habitat in Guinea Isert 5.

Rami inferne teretes, cortice cinereo, superne villis aequot adspa-
si. Folia pedicelata, opposita vel alterna, quadri-collicaria, basi
interdum ^{sub} cordata, acuminata, subtus venosa, apulis minutissimis
scaberrima, integerrima vel extrossum dentibus utroque 2-
3 remotis grossis serrata. Pedunculus terminalis, brevis.

flores saepe 14. Calyx basi coarctatus, ^{interdum 4-angulatus} campanulatus, angulatus,
laciniis ovatis, acutis, pallidis, raris pilosis, venosis, ciliatis

Corolla fere tri-pollicaris; tubus angustus villosus, infundibul-
formis: limbus squaeformis; laciniis subrotundis, patentibus.

Clerodendron

Habitat in Philippinis herb. Gui

Rami tetragonis, glabri. Folia petiolata, palmata, cordata, obscure remote denticulata, supra glabra, squamata. Panicula ramis dichotomis an squamatum meum.

nitidam Cleodendron

Habitat ad Surram Leonam herb. La Martini

Globum. Bolla petiolata, ses qui pollicaria, ovato-oblonga, venata, integerrima, nitida, subvenosa, acuminata, subtus punctata. Peduncululus terminalis, tetragonus, primum 4fidus, dein duobus divisus. Corolla purpurea, pollicares. Calycis laciniis lineari-conicatis

Plendendron

Tava Commerfon herb. Fusf.

folia petiolata, basi cordata, quinqueloba, spithamea: lobi ovati, acuti, integerrimi, inferne duo major, supra ^{minuti} ~~minuti~~ nervis, subtus punctati. Racemus terminalis ^{pedalis} rami oppositi, bis dichotomi, bipollicares, patentes. Flores pedicellati, oppositi: foliolum oblongum, petiolatum, utraque ad basin sinuuli rami pediculi ope lentis villosi. Bracteae subulatae, parvae. Pedicelli oppositi. Corolla semipollicaris: lacinae ~~ob~~ longa, glabra.

Denticulatus Cleodendron.

Habitat in insula francis Commafor herb. Lus.

Rami glabri. Folia petiolata, 3-4 pollicaria, cordata, lobo altero altero incumbente, ut si-
nus non apertus nisi basi ad insertionem petioli, acuta, remote dentia alata, venosa, supra
pilis minuti raris albis. Petioli longitudine foliorum, spatium utraque inter petiolos basi
utraque peloso, incanus. Pedunculi et calycis colorati, glabri. Calycis lacinis acuta
crata. Corolla semipollicaris.

infortunatum. foliis cordatis tomentosi fl. Zeyl. 232 &

Petalites agris Rumph. amb. 4. p. 100. t. 49.

Peraya Rheed. mat. 2. p. 41. t. 25.

β. Clerodendrum foliis lato acuminato Burm Zeyl. 66. t. 29.

Habitat in India & ^{corolla extus}

Rami petioli folia subtus et panicula pubescentia.

Folia petiolata palmata supra glabra, nervosa, venosa.

panicula terminalis bracteata. folium parvum ovatum

reticulatum ad ramificationes. Pediculi parvissimi tubo

tomi. Pedicellus in singula dichotomia calycis lacinis

lanicollata. Corolla pollicaris. Stamina corolla longum

Rami et folia subtus flavo villosi subtomentosi

atque
attenuati subtus villosi

Fortunatum C. foliis lanceolatis integerrimis Len Am. acad 4. p. 320 &
of Beck. t. 220. t. 11.

Habitat in India &

Rami teretibus pubescentes. folia lato lanceolata venosa
tripolliciana utrinque glabra venosa. Pedunculis axil-
laribus foliis brevioribus pubescentibus bidentatis pedicelli ultra
tres vel quatuor racemosi. calycis lacinis ovatis

phlomidis Pterodendron ^{Pterodendron} foliis ovatis integris dentatis angulatisque, pedunculis
axillaribus subtrifloris
Volumenta multiflora, ramis inermibus, foliis oppositis cordatis
subdentatis acuminatis, pedunculis axillaribus multifloris
Burm. ind. pag. 134. Tab. 45. fig. 1.

Pterodendron phlomidis foliis ovatis serratis Canis Lin Suppl.

Habitat in India pag. 292 x I. V

Folia utrinque acuta, pollicaria, alia integerrima, dentata
vel dentato-subangulata in eodem ramo, in cano. v. resp. India

Petioluli filiformes, inermes, laevi

Pedunculi plerumque triflori. Bractea duae in singulo pedicello

Calis + campanulatus, quinque fidiis: laciniis lanceolatis acumi

Embellis trifloris pollicariis pubescentibus

Rami subtetragonis cortice griseo. ramuli villosi inermes.

^{articulata}
Cephylla Tamane ramis ^{articulatis, apicibus turbinatis mucronatis} ~~articulatis~~ ^{foliis turbinatis vaginantibus basi mucronatis}
^{racemosis}
Thuja ^{aphylla}: Probilis quadri valvibus, foliis turbinatis vaginantibus
 line mucronatis, frondibus umbriatis Lin. Am. ser. 4. p. 295 *
 P. S. pag. 861.

Habetatur India Orientali

^{filiformes}
~~Sunt ex virgatus ramis ramulosis basi articulatis umbri-~~
~~cales squas foliis crassis vaginis foliaceis~~ ^{terribilibus} ^{quercifoliae} ^{ragini} ^{persisten-}
^{rosi} ^{persistentibus} ^{quor-} ^{dibus} ^{ramulosis} ^{virgati} ^{ramulis} ^{filiformibus} ^{gracili-}
^{alternis} ^{articulatis}, ^{articulis} ^{turbinatis} ^{ad} ^{ab} ^{altero} ^{latere} ^{mucro-}
^{ramulorum} ^{virgato} ^{natis} ^a ^{mucronibus} ^{alternis} ^{umbriatis}. ^{Spica} ^{ad} ^{apicem}
^{ramulorum} ^{virgato} ^{pollicares} ^{umbriata} ^{teretes}. ^{Bractea} ^{ovata} ^{acuta}
^{patentissima} ^{persistente} ^{concava}. ^{Pappi} ³ ^{phyllis} ^{foveolis} ^{setis}
^{longis} ^{alternis} ^{basi} ⁱⁿ ^{vaginam} ^{linearia}. ^{Corolla} ^{magnitudine} ^{Tamar} ^{gallica} ^{famena} ⁵ ^{petala}
^{connata} ^{linearia}. ^{Pappi} ⁴ ^{rayona} ^{acuminata} ^{semina} ^{ut} ⁱⁿ ^{reliquis}

Calamitofum *C. foliis ovalibus subdentatis nudis* Kun. Mant. 90 + P. V. pag

578

Volhamenia alternifolia

Burm ind. 1737. Tab. 48. fig. 1.

Habitat in Java &

Solia petiolata & *pyroptera* *f. ovalis*, serrato-subdentata
utroque nuda. Panícula brachiata. Flores congestius
paulo minores Kun

non vidi

paniculatum Clerodendrum foliis quinquelobis denticulatis glabris, pani-
cula brachiata, axillis lanatis.

Clerodendrum paniculatum: foliis lobatis serratis, panicula
amplissima ^{in magna p. 904} Lin. S. V. pag. 578. ex cl. p. Synon: Burman. fl. Ind.

Habitat in India

Rami tetragoni, glabri, purpurei, quadrifurcati lateribus sulco
exarati.

Folia petiolata, opposita, palmata, inferiora spithamea, quin-
queloba, basi, cordata, utrinque glabra, remote denticulata.

Lobi lanceolati acuti.

Petiolis teretes, sulcati, crassitie, penno columbino, basi
~~axilla~~ ^{axilla} ~~petio~~ ^{petio} longis crispis albidis caulem ambientibus
sive ac ad basin ramorum panicula.

Panicula terminalis, semipedalis, brachiata. Pedunculi
partialis horizontales, oppositi, glabri, multoties dichoto-
me divisi: ramuli ^{ultimi} patenti fini. Pedicelli capilla-
res

Foliola petiolata, cordato lanceolata ad basin peduncu-
lorum partialium: subulatum ad basin ramulorum

Calyx glaber: laciniis lanceolatis

verte

Corolla pollicaris. Tubus feli formis. Lacinio lambi oblongo

Squamatum
5-6 cordifolium Clerodendrum foliis cordatis obscure angulatis; panicula ramis

dichotomis glabris

Habitat in India Orientali La Mart.

Rami tetragoni, glabri, sulcis quatuor exarati.

Solca petiolata, bipollicaria et ultra, cordata, basi sine profundo,

panem acuminata, tri- vel quinque nervia: nervis adspen-

dentibus, tenera, glabra, versus basin obscure angulata,

versus denticulata, interdum pelis minutissimis ad marginem

ciliata, subtus tecto squamis orbiculatis minutissimis

Petiolus unguicularis subuloser

Panicula terminalis, palmatis, patens glabra. Pedunculi

et ter ~~minime~~ ^{mediis} ~~minime~~ ^{minime} ~~incompositi~~ ^{incompositi}: pedicelli ^{v. r.} ~~an~~

floris in singulo dichotomia.

Soleola approposita, petiolata, cordata, subtus villosiuscula

subprimis ramificationibus, fusum versus per se angu-

sticoria: sub ramulis subulata, sessilia

Calycis soleola ovata, ^{acuta} ~~rotata~~, glabra.

Corolla tubus unguicularis, glaber

Hamina volla duplo longior

Differt a Clerod. infortunato, foliis basi sinu profundo ^{et ipsi} mixioni
bus, glabris, Panicula ^{calyce et corolla} glabra. Ramis tetragonis linea exaratis

a Clerod. trichotomum, ut ex specimine ab amicisimo Thunb.
missum, foliis basi sinu profundo, foliis omnibus integris
nec aliis lobatis aliis integris. Convenit sive calyce

Anichotomum Q. foliis lobatis indivisisque lato-ovatis integri panicula
Anichotoma Thunb. fl. Jap. p. 2564 L.V. pag. 578.

Habitat. in Japonia ^{per omnes}
Rami ~~glabri~~ ^{utri usque} partes superiores obtuse tetragoni
Folia longi petiolata lato palmam lata s' ad 6 pollicaria
attenuata venosa, subtus pallidiora, ad costam et nervi-
miponibus villosa, inferiora subtuloba superiora indivisa.
Panicula terminalis ^{per} magna primum *Trichotoma*
dica ^{usque ad apicem} *Trichotoma*. Pedunculus solitarius in singulis ^{usque} sursum
pedicelli uniflori lacina calycis ovato

5-6 diversifolium Clerodendrum foliis integris trilobis que ovatis, panicula ramis dichotomis
villofis
Habitat in India orientali dom. La Martia

Rami tetragoni, quadrifidati, superne villosi

Folia petiolata, ovata, integra vel triloba, lanera, trinervis, basi
acuta, subtinnervata, glaberrima; sequi lobus intermedius ac-
minatus, laterales breves.

Petiolus folio triplo vel quadruplo brevior

Panicula terminalis, semipedalis, ramis ramulisque villosis

Pedunculi parciales oppositi, patenti, bis vel ter dichotomi;
pedicelli demum racemosi.

Pedicella ad basin pedunculorum parvialium petiolata, oblonga, utrin-
que acuta, integerrima; elegia subulata.

Flores pedicellati, aeterni, distantes

Calyx pubescens; laciniis oblongis acutis

Corolla tubus pubescens, prolixius: labium superius bifidum; la-
ciniis erectis lineari-bus: inferius in lobum patens sinum, lobis
oblongis lateralibus brevioribus.

(Cl. trichotomo differt: foliis angustioribus, ramis superne villosis
Panicula ^{villoso} dichotoma tandem racemoso, laciniis calycis oblongis
lobi corolla vix aequales

Cobata Plerodendron
Volthameria foliis quinquelobis denticulatis, ^{glabris} panicula brachiatis, axi-
li lanatis
Habitat in India orientali Honig

Rami tetragoni, glabri, purpurei, quadrifurcati. ^{quinquelobo.}
Folia petiolata, opposita, palmata et est pithamso, utrinque basi cordata
utrinque eglabra; quinque nervia: ^{remote denticulata;} lobi lanceolati, acuti
Petiolus ^{tertius} ~~fulcatus~~ ^{emissis} pennis columbino: axillis basi pilis
longis crispis albidis, densis caulem amplexibus ut et ad basin ramorum
panicula

Panicula terminali magna semipedalis, brachiata, pedunculis par-
tibus horizontalibus ^{oppositis} dichotomi glabri multoties divisi, ramulis patentibus
Flores ^{capellae} foliolarum petiolarum ^{terminales} lanceolatas ad basin pedunculi ^{partia}
cum, seta subulata ad basin ramulorum.

Calyx glaber lanceolatus Cociniis
Corolla pollicaris: tubo filiformi: laciniis limbi oblongo.
est est Plerod. paniculatum