

Plethra

1. *alnifolia* C. foliis obovato-lanceolatis, racemis simplicibus, calycibus acutis
2. *paniculata* C. foliis lanceolatis, racemo composito.
3. *arborea* C. foliis oblongis coriaceis, racemo simplici, calycibus obtusis
4. *trifolia* C. arborea, foliis oblongo-lanceolatis integerrimis subtus incanis: racemis paniculatis filiciformibus tomentosis

tenifolia

V. folia C. arborea, foliis oblongis lanceolatis integerrimis subtus in
canis: racemis paniculatis spiciformibus tomentosis Sw. prodr
pag. 74

Loas. herb. 2. Pl. 8. 190. fig. 2.

Volhamenia Brown. Jam. 24. Tab. 21. fig. 1.

Texas occidentalis Lin.

Habitat in India occidentali

Rami teretes superne tomentosi folia petiolata alterne bis. tri
peltata supra glabra arena hic usadem nervi lineati
subtus venos nervi purpurei, extropum interdum subserotum

boti ramosi

lobi rotundata apice obtusiuscula. Racemi terminales
pedicelli velis pedicelli bracteati calycibusque ferrugineo tomen
topi, pedicelli sparsi breves reflexi. calycis foliola ovata
obtusae

nonodi

acuminata Clathra

Habitat in America septentr. Richard.

habeo

Herba Plethra

Habitat in Brasilia herb. Lusit.

Rami crassitie perna cygneae, superne ferrugineo villosi-tomentosi. Folia opposita
apice ramorum, petiolata, approximata, tri-pollicaria, oblonga basi versus an-
tice sensim angustata, rigida, supra oculo armato punctis villosis stellatis scabris
siccata, integerrima, ~~acuminata~~ emarginata, subtus venoso-reticulata, tomento
laevis, nervi venaeque ferrugineo villosi-tomentosi. Racemi terminaliter
plures, simpliciter vel subcompositi, ferrugineo villosi-tomentosi, quadri-pollicaria
res, multiflori. Flores parvi, pediculi angulati.

2

paniculata *Clethra* foliis lanceolatis utraque ^{nuda} ~~globis~~, racemis composito.

Clethra paniculata fruticosa, foliis lanceolatis utraque nudis, flori-
bus paniculatis. Hort. kew. pars 1. 2. p. 73.

Habitat in America septentrionali

Pedunculis et Calycis ^{lenatis} ~~stomatofis~~, foliisque basi angustatis ut in ^{lenatis} ~~procedent~~
differt vero foliis angustioribus, brevioribus, pallidioribus ~~obtusis~~
~~marginibus~~ ~~ac~~ apice saepe magis rotundioribus, subtus nudis utraque
nuda, praecipue vero racemis composito ~~sed~~ ^{semper} ~~ali~~ ~~com~~
~~posito~~ ~~pro~~ composito ex racemis palmaribus simplicibus, sciden-
tibus. Bractea subulata pedicello brevior. Flores minores foliorum
calycina acutiuscula

1. alni folia Clethra foliis obovato lanceolatis, racemis ^{simplicibus} pinniformibus, calycibus ^{acutis}, bracteis ^{persistentibus} Hort. kew. pag. 73.

Alnifolia americana serrata, floribus pentapetalis albis in pinnam dispositis
Pluk alm. 10. t. 115. fig. 1.

Clethra alnifolia Gron. Voy. 47. Du Ham. 1. p. 176. t. 71. Lin. L. V.

Clethra alnifolia foliis obovato lanceolatis, racemis pinniformibus,
calycibus acutis, bracteis persistentibus Hort. kew. t. 2. p. 73.

Denudata ~~o~~ foliis utrinque nudis
pubescentibus β . foliis subtus tomentosis bracteis persistentibus

γ . foliis subtus tomentosis bracteis deciduis
Habit in Carolina, Virginia, Pennsylvania &

fructu ^{rami} et ^{teretes} tomentoso cinerei. folia petiolata, alterna bipollicaria
^{pub. ult. no} subtus tomentosa, incana, acuta, a medio ad apicem serrata ^{dentatis}
dentibus mucronatis, supra vincta adpersa pilis minutis raris, secundum
venas pubescentes basi attenuata. racemus terminalis palmatis
pedunculis incanis adpressis pilis ferrugineis uti bracteis pedicelli ^{conspic}
breviores ^{bracteis} sparsi, bracteis lanceolato pedicello longioribus Calycibus
tomentosis

varietas γ . sub nomine glauca ex hort. Tramon differt absentia pilorum ferrug.
nearum in pedunculo, bracteis deceduis cotenim similibus. Stamina et stylus
corolla duplo longiora

3 Arborea *Plethra* foliis oblongis coriaceis, ^{ferratis ~~prope~~ ~~apice~~} racemis simplicibus, floribus calycibus obtusis
Plethra arborea, arborea foliis oblongo lanceolatis utrinque glabris, racemis simplicibus, calycibus obtusis Hort. kew. 2. p. 76 73

Habitata in Madera

Arborea. Folia oblonga, ~~se~~ subobtusata vel oblongo lanceolata, petiolata, glaberrima mucronata consistentia foliorum thicis, superiora crenato-denticulata, inferiora a medio ad apicem remote serrata, serraturis spinulosis pungentibus, supra laevissima lucida sesqui pollicaria. Pedunculus communis striatus pubescenti pilosus. Ped. palmari. Pedicelli tomentosi incani. E locis annui triplo majores quam in prociacibus. Calycis foliolis ^{petalis} tomentosa incana, ovata obtusa. Corolla calyce duplo longior. ^{petalis} Hamina longitudine corollae. Germen pilosissimum. Stylus longitudine staminum