

Collinsonia

Genis. plant. Schreb. n 51

~~Cor. inequalis : labio inferiore multifido, capillari. Sem. 1. perfectum~~

~~1. canaliculatis foliis ovatis, caulibusque glabris~~

~~2. scabriusculis & foliis ovatis subcordatis pilosiusculis, caule pilosiusculo scabido~~

57

~~78~~

60. Collinsonia. Cal. 2-labiatus : labio superiore tridentato. Cor. inequalis : labio inferiore multifido, capillari. Sem. 4. tria abortientia.

canadensis 1. Collinsonia foliis ovatis caulibusque glabris Hort. Kew. 1. p. 47.

Collinsonia canadensis Lin. Hort. Cliff. 14. t. 5

Habitat in Virginia, Canada Sylvis 4

Caulis erectus, bipedalis, 4-5 angustus. Folia ^{palmaria, petiolata} oblongata; basi ovata,
~~grassa serrata, serrata mucronata~~. Racemi terminales, infera
^{vel quatuor}
saepe ramis duobus oppositis. Pedicelli oppositi. Bractea lanceolata

vapeas *Sarmienta*
fl. *Penus. sp. D. tab. 7. fig. 6.*

Utricularia foliis carnosis scandentibus. *Sewill. Penus. 3. p. 69. tab. 43.*

Habitata in nemoribus Chile copiose.

*Caulis ramosissimus, sarmentosus, Vires: rami dependentes. Folia opposita, breviter
petiolata, ovata, carnosae, venia, pubes albicantia, Anagae punctata. Pedun-
culi terminales, longi, filiformes, uni vel biflori. Bracteae duae, oblongae, crispae
sub pediculo Calycis. Calycis hirsuti. Corolla punicea, extus pubescens*

lacini 4 aequales s. emarginatae lobis, imbricatis

*Corolla tubus obovatus ventricosus, basi angustissimus, fauce corollata
limbo 3-quefido laciniis aequalibus. Stamina plura recept. carnosum*

Scabruscula & Collinsonia foliis subcordatis pilosis acutis, caule pilosiusculo
et scabro Hort. Kew 1. p. 47

Habitat in Florida orientali &

Folia ovata, subcordata, ~~quasi dimidata~~ ~~semiterminaliter~~

Pedunculi villosi. Racemi inferne paucis duobus vel quatuor.

1. *Canadensis* Collinsonia L.V. p. 72. 1. Hort. Cliff. 14. t. 5. ~~Walm 47. p. 317. Spec. pl.~~

~~p. 29.~~
Habitat in Virginia, Canada sylvis

Canadensis foliis ovatis caulibusque glabris Hort. kew. D. p. 47

Caulis erectus bipedalis glaber folia petiolata opposita
ovata attenuata serrata nervosa venosa, palmata
serraturis mucronatis Racemi terminales inferne
ramis duobus ^{oppositis} vel quatuor ~~op~~ ~~ramis~~ ~~compositis~~.
br. pedicellis oppositis bracteis lanceolatis

tuberosa
Serdina

3 Collinsonia foliis oblongo-ovatis utrinque acutis glabris.

Collinsonia Serdina: panicula Vermenali ramosissima Walt. fl.

Carol. #149.

Habitat in Carolina Bosc 4

Radix Tuberosa. Caulis infera teretis, scelus, superne obtus tetragonus,
interdum pedalis, ramosus, opre lentis rare villosus. Folia desqu pollicis
ni, nervoso venosa, subtus pallidiora, opre serata, remota
mutuati racemi ex apice caulis et ramulorum axillarum
elongati: pedunculus pedicelli et calycis, prosertim ore filioi. Pactea
narrow, subulata. Pedicelli oppositi, remoti. Calycis et flores multo
minoros quam in canadensi.

Collinsonia Tuberosa; radice tuberosa; foliis subrhombica ovalebus,
utrinque angustato-acutis: calycis dentibus petalacis, tubo longis
ribus. Michaux fl. bor. Amer. 1. p. 17.

Serotina ~~C. panicula terminali ramis~~ ~~lima~~ Walt. fl. Car. p.

49

Habitat in Carolina

non vidi

procox 4. Collinsonia foliis infinis petiolatis, floralibus pluribus ovatis sessilibus,
floribus axillaribus lateralibus & terminalibus. Walt. Carol. 65

Habitat in Carolina

Cum differentia Waltheri, aliam plantam a Col. scabrescens indicare
videtur, potius uti speciem distinctam tradere jureat usque dum melius

discernitur

Habitus Comp. spec. Caulis herbaceus, tetragonus, grosse serrata; serraturis mucronatis,
folia opposita ^{reticulata} pubes ad circulum armatum punctata.

~~Tabernaemontana~~ ~~Quilleyana~~

~~Habitat in America boreali herb. Richard.~~

~~Caulis pedalis interdum, ramosus, folia sesquipedaliana, oblongo
ovata~~

herb