

Collyris

Cardata. Cor. globosa. ~~fulva~~ Corpus truncatum quinqueangulum, fulvum
lineare erectum, bifidum laciniis reflexis clavatis ad siniculum angulatum
Corporis truncati

Collyri
minor Mateola foliis oblongo-ovatis.

nummularia lactea minor Rumph. H. Amb. 6. p. t-176

Fig. 1.

Habitat in India Orientali Vioniz

Rami filiformes, teretes, glauci, et folia radianter, flexuosi. So

quadriplo
minoris quam
in precedente ves
lea sessilia, opposita, vel unguiculata, integerrima, utrinque al-
ta, carnosa, avenia, enervia, subtus glauca, et ~~et~~ cores utille

Ar adino multis effigie prociubens ex cauliculis
geniculi radiceo, tuba capsularis ex insula Cheusan.
Platan. a Malth. p. 4. v. 354. f. 2. huc pertinet?

Ungitana l. foliis cordatis ^{oblongis} crispis calycibus spumosi sulcatis strigosis bracteis
concavis mucronatis 'Etl. - l. pag. 34. n. 30.

Salaria Ungitana foetidissima hispida flore albo Tournef. J. b. 4. p. 174
Habitat in agris boreali &

Habitus l. spumosi. Caules basi frutescentes ramosi erecti, villis
longis tomentibus adpersi intermixtis brevioribus glutinosi in calycibus
rigidioribus albis ^{et pilulis} pellucidis. Folia rugosa h. p. tripolliciana
ventricelli ^{circiter} octoflori bracteis cordatis ciliatis longi-
tudine calycis. Calyx amplus. Corolla alba

Salaria foetida foliis cordatis inaequaliter dentatis rugosis h. p. bracteis
cordato-acutis ciliatis longitudine calycum La Mart. Tab. Enc. pag. 69

proter Synon: Tournef. Commes. de Specimen herb. Fussiae attulit
Salaria orientalis v. *bucifolia*, flore partim albo, partim flavescente
Tournef.

Collini
Majus *Motela folis orbiculati*

Habitat in sylvis aridis et collinis prope Malaccam et Solangor
hony.

Caules scandentes, truncis arborum afflexi truncis arborum
affixi mediante squamis oppositis, reniformibus, albidis, latera in
seniori fibrillata radiantes, proterea Vesiculae plures, sessiles ovato
oblongae, alt. bi pollicari vel ultra, obtusissimo, extus glaberrimae
cuticibus magnis notatae, intus parum inflatae, purpureae, fibra
longitudinalis libera, ramosissima basi vesiculae inserta, saepe aqua
dulci plena. Rami teretes, flexuosi, radiantes, inferne a tubis
culis pluribus nodosi, superne squales. ^{oppositae} Folia remotissima,
petiolata, vix pollicaria, subrotunda, acuta, interdum parum
oblonga, carnosae. Clorox axillares.
Calyx parvus, quinquefidus. Corolla globosa, ore quinquefido: lacini

nis erectiuscula, parva. Corpus truncatum apice convexum, quinqueangula-
re, ad singulum angulum fulcrum erectum, apice bifidum: lacinis re-
curvatis, longitudine fere corporis truncati, apicem versus intraspiratis
clavatis magis linearis, quam in affinibus; angulis fereis, marginibus
parum callosis, ~~ad~~ ad singulum angulum fulcrum erectum, apice
bifidum: lacinis recurvatis, clavatis, longitudine corporis truncati.
Hypomochlii parva, ut in affinibus inter squamulas: Anthera
oblongo-clavata, plana, callosa, flavo. Stigmata duo, in centro corpo-
ris truncati eoque parum longiora, ante coronata, eoque parum longiora.
Sollivuli lineari-oblongi, rostrati, latere superiori planiusculi,
apicibus incurvatis. Semina comosa.

Acrida
9

~~300~~

Cor. niger: labes inferiori reflexo bifido. P. quinquepartitus. Cap. bi
loculari 4 valvi

i. magellanica

