

Columella

Flor. Peru. prod. p. 3. t. 1.

color: ~~rubra~~

Cal. 5quepartitis. Cor. rotata. Anth. plicata. Caps. Didyma ~~valvulis~~
duplicatis ~~polygamis~~

62

~~62~~ Columella. Cal. 5partitis. Cor. rotata^{super}. Anth. plicata. Caps. 2-val
vis, 2-locularis: valvulis duplicatis.

~~quatuor inflexum~~

obovata? Columella
foliis obovatis serratis integerrimisque, ^{lesibus} pedunculis subuni flori +
Columella obovata, foliis obovatis serrato-dentatis integerrimisque, pedunculis uni
flori Fl. Per. 1. pag. 20. Tab. 12. fig. 6.

Habitat in collibus aridis Tamao & Penuria;

Frutex octopiedalis. Caulis erectus, teres. Rami equoqi brachia, oppositi. Folia
opposita, conferta, sessilia supra concava, nitida, glauca, subtus incana, mi-
noa integerrima. Pedunculi terminales, breves, uni flori; foliolam utrinque
ad basin pedicellorum, ad duo sub calyce. ~~Caulis luteus.~~ Fl. Per. 1. c. +

Habitus. Caulis arborosus: Rami oppositi. Folia opposita, indivisa,
serrata. Pedunculi ramosi. Flores lutei.

4 terminales: foliola

Duo sub pedum

culis, linearia,

decidua.

oblonga 1. Columella foliis oblongis extorsum serratis petiolatis, floribus corym-
bosis +

Columella oblonga; foliis oblongis extorsum serratis floribus corymbosis

Bl. Per.: 1. p. 20. t. 8. f. a

Habitat in Peruvia Sylvis +

Arbor quadrangularis, ramosissima. Rami veretes; cortice cinereo.

Folia approximata, supra glabra nitida, subtus pubescentia, albi-
cantia. Corymbi terminales. Pedunculi trifidi: pedicelli uniflori

foliis laevis duo opposita in singulo. Corolla lobata. La cinis con-
cava. Bl. per. l. c.

~~nervosum ^{foliis pinnatis} ^{foliis} Tasmense. foliis ovatis trinerviis Lour. fl. Cochinch. pag. 20.~~

~~Habitat in Cochinchina~~

~~Truxes scandens, herborum ramis. Folia impari pinnata foliolo
longipolliciana, integerrima acuminata, nitida. peduncululi
terminales, multiflori. Corolla lacus 5-0~~

non vidi