

Conium

1. maculatum C. seminebus striatis, pinnis oblongis obtusis obtuse serratis in apice.
2. latifolium C. foliis radicalibus bipinnatis, pinnis oppositis cuneatis trifidis
seminebus mucronatis
3. niger C. seminebus submucronatis, pedunculis falcatis folioli canaliculati
obtusis
4. africanum C. seminebus mucronatis, petiolis pedunculisque corvibus
5. tenuifolium C. foliis radicalibus caulibusque simplicibus lineantibus (Linn.
fili folium foliis filiformibus involucri elongati Thunb. prodr. p. 50)

Debotomum Conuini

Desfont.

Trochileum leptaneum ciliata folio semine striato Tournef. Inst. v. h.

Habitat in Leptanea Schousboe

Conium

4 tenuifolium Conium foliis radicalibus caulinisque simplicibus linearibus

Habitat in Capite bono spci. Dedit Dr Burmannus.

Caulis herbaceus, erectus, pedalis, vix ramosus, tenuis, teres, glaber, leviter striatus.

Folia omnia simplicia, angustissima, ~~inferiora quadripollicaria basi~~
parum latiora, suboperculata, glaberrima; ~~inferiora quadripollicaria~~
~~et parum ultra: caulina remota.~~

Pedunculi laterales et terminales, pauci, remoti.

~~Umbella~~ parva, quinquedactyla. ~~Involucrum~~ pericha umbellula
circiter 8. vel 10 flori. ~~Involucrum pentaphyllum: folioli subulati.~~

~~Involucrum~~ pentaphyllum: folioli subulati, rigidiusculi, umbella
brevioribus. Involucella longitudine umbellula.

Semina oblonga, striata, parva, glabra. Styli persistentes; divaricati

maculatum C. seminibus striatis Lin. t. V. Haecq. fl. austr. 2. t. 156.

Cicutaria major vulgari Clef. hist. 2. p. 200

Cicutaria major Bauh. pin. 160

Habitat in Europa cultis, agnis, rud eratis &

Caule erectus oxygale ramosus fistulosus teres striato pallidus
maculis purpurascentibus adpressis. folia magna tripartita
& - petioli oblongis obtusis, obtuse serrato incisi. involucri
foliola parva.

Suffruticosum & seminebus angulato-
stratis, caule suffruticoso
Merg. fl. cop.

non vidi

latifolium C. ^{radicalibus} foliis bipinnatis pinnis oppositis cuneatis trifidis seminibus
municatis

Habitat in umbrosis laevis

Caulis erectus rimosus glaber striatus. ^{pinnis foliorum} Soliola radicalium.

Latitudo magna, bipollicata basi attenuata estrosum dilatata
trifida glabra laciniis lanceolatis intermedio superne et
utroque dente ~~latitudo~~ ^{metata} superiora pinnata hae foliolis
lanceola mucronatis summa simplicia. Umbella conoide
involucra 12 phylla parva semina oblonga

Sede All. huiusmodi

negosum Concuⁿ Semenbuⁿ negosi Thunb. prodr fl. Cap. pag. 50
Habitat in Cap. b. spai

non vidi

rigens
agens

♀ seminitibus submunitatis, pedunculis sulcatis, foliolis canaliculatis
obtusis ~~Hort. Clus. Mant. 352 & 512 & 515. v. p. 270. Spec. pl. pag. 350.~~

~~Caucalis africana, folio minore nuda Boeck. hyst. 1. p. 63. & 63~~

~~Habitat in Africa~~

Habitat ad Cap. b. Speitz

~~Boeck. loco~~

Planta spontanea saepe vix digitali rigens. Rami longiores paten-
tissimi. Folia bipinnata pinnis corduplicatis obtusis crenulatis

*Pedunculis
elongatis* in culta ^{sup} pinnatis pedis duri colore tuta. Umbello oppositi folia

Involucro pentaphyllo foliolo uno vel altero in strutura fere
foliorum caulis. Involucelli foliola pauca & phylla foliolis
crenatis margine membranacei. Semina mucicata

Culta laxior evadit ramifera vagina foliorum brevi margine
lato membranaceo.

africanum \odot Seminibus mucronatis, petiolo pedunculo lobatis Lin. Spec. pl.
p. 350 + Mant. 352. S. V. pag. 270. Gaëq. hort. 2. 4. 194.

Caucalis africana, folio minore nudo Boerh. Lugd. 1. p. 63. 4. 63.

Habitat in Africa \odot

Planta glauca ut Seseli, folus fumariae et angustioribus tripennatis
pennis multo sedis. Involucrum foliola 6. bene lanceolata. Pedun-
culi oppositi folii. Semina oblonga mucronata, majori gibba quam
lata, stylis persistentibus

C. graminifol. B.
Daundum Pal.
suffruticosum Berg