

Connarus

nemorosi. C. foliis ternatis quaternisve ovatis acuminatis acutis
utrinque venis ex partibus
Habitat in nemoribus. And. orientalis. Virens.
Rami teretes fusi et punctis numerosi elevatis ferrugineis
siccis. Paniculae. Sola pedicelata ternata vel pinnata
to bipinnata ~~cum~~ abque impari. foliola pedicelata ovata fere
pollicaria glabra membranacea utrinque venis venoso
reticulata glabra. paniculae ex apice pedunculata
calycibusque breviter tomentosis. Petala linearia unguiculata
absque setis ad basin. fructus pollicariis oblongis glabris

Pentagynus Connarus Cavon.

Habitat in herb. Aust.

Rami Verdes, glabri, cortice purpurasciente. Folia petiolata, alterna,
ovata, parum acuminata, integerrima, pollicaria obsolete trinervia,
venosa, subtus venis erectantibus. Panícula Verminalis, patens.
(racemo decomposito) Calycis et pedunculi orulo armato vomentis
an monocarpo-lin.

Connarus

Habitat in Puerto Rico Herb. Deffend.

Folia ternata et pinnata: foliola internati opposita, in pinnatis alterna
^{binucleata}
vena utrinque exstantes, oblonga, supra nitida Racemi axillares; Terni,
compositi. Calycis laciniis ovatis. Calycis laciniis ovatis.
forte non nisi varietas Caranensis

Connanis

Connanis La Mare

Habitat in Capenna herb. Desfont.

Solucola 6. alterna, unciam terminalata, 5 pollicaria, interiora minoru, ellipti-
ca, acuminata, utrinque glabra et venis exstantibus, integerrima. Petioli fis-
sili, pariter breves, nigri, Racemi compositi

monocarpus Connarus Lin fl. reyl. 2404

Phaseolus arboreus reylanicus monocarpus Raj. suppl.
430.

Phaseolus reylanicus trifolius, phaseolifolius, floribus
prosp. spicati Burm. reyl. 199. t. 89.

Habitat in India t. Specimen misit Thunberg

Com. pennatus foliis ternatis p

folia ternata vel pennata, foliola ovata oblonga
acuminata, obtusa submarginata utraque glabra, uterque
rima supra lora obtuse venosa, subtus tenuis p. veno
so reticulata venis exstantibus. floris paniculati villosi
subtomentosi

— foliis ternatis folioli ovatis acuminatis obtusi emarginati
subtus venis exstantibus floribus tomentosis
~~an lei et pennatus La Mad. Casuarina plant~~

pentagynus \odot foliis ternatis Vernatis Vinensis, flentibus pentagynis
La Mart. Emph. Car. des pag 376 tab. 223.

Connarus pentagynus foliis omnibus 3 foliatis: foliulis ovatis acutis
subtus venosis, flentibus pentagynis La Mart. Car.

Habitatio Madagascar.

Rami teretes glabri purpurascens folia petiolata alterna.

Foliola ^{petiolata} ternata ovata acuminata acuta Vinensis supra diversa
loria subtus tenuissime venosa ^{et emedum 2. pollicia}
et lateralia rariore axillares ^{folia parva breviora} et terminales compo-
si. Calycis tomentosi petala glabra

foliis ternatis foliulis ovatis acuminatis acutis Vinensis
venis immersis, folia parva rariore axillares terminali-
busque compositis

Connarus

africanus

Connarus foliis ternatis lanceolatis - ellipticis, venis subtus protuberantibus.

Connarus africanus: foliis ternatis ovatis utraque acutis, subtus nervosis, floribus paniculatis monogyris. La Mart. Enigil.

Top. 2. p. 95

Connarus africanus foliis trifoliatis, petalis ebarbatis, floribus paniculatis Caran. Suppl. 7ma tab. 221.

Habitat in montibus Terra Leonis dedit dn L Hertert

Rami teretes, glabri

folia petiolata, alterna: foliola petiolata, palmata et ultra acumine Congo ternata, glaberrima, supra ^{loris} nervis et venis

transversis coarctis subsimplicibus immixtis, subtus exstantibus, pinnate gerrima intermediis nygosis, intermediis longior

Petiolis partialis infra foliolo inermis nygosis long

Panicula ternata, plata

Burmannii Drosera scapis radiatis calycibusque glabris, foliis spatulatis sessilibus
Ror. foliis zeylanica, foliis rotundis in orbem dispositis expansis velatis
Her m. mus. zeyl. 10.

Ror. foliis foliis circa radicem in orbem dispositis Burm zeyl. 207. t. 94.
fig. 2.

~~Ror. foliis rotundifolia L. fl. zeyl. 120~~

Habitat in Zeylona

a Dr. rotundifolia distingui meretur foliis sessilibus rotundis basi attenuatis
feminaeque ciliatis ciliis tenuioribus aequalibus: a cuneifolia
distinguitur glabritate scapis et pedicellorum et calycibus, et quod triplis
minor

monocarpus Connarus. L. Fl. Zey. 112. * Sp. pl. 944. L. V. 612.

Phaseolus arborefcens Zeylanicus monocarpus Raii Supl. 430.

Rhizis Zeylanicus trifoliatus, phaseoli facie, floribus copiosis spicatis Burm. Zeyl. 199
L. 89.

Habitat in nemorosis Indiae h.

Rami teretes cortice fusco ferrugines punctatis ^{conacica} supremi ~~rami~~ in punctatis.

Foliae petiolata ~~alterna~~ ^{pedunculata} ternata foliola petiolata ovata, intercostima venosa
nervosa pollicaria. ^{cap. nigrosi} floris paniculati versus summum ramum

thyrsoides ^{vel contracta} axillares terminalesque floris copiosi, calyx tomentosus deciduus fructus
pedicellatus glaber

Decumbens P. foliis ternatis, ^{cordatis} hirsutis, folioli ~~obovatis~~, caule decumbente
herbarico Thunb. in Nom. Ant. 2. bot. 1 N. pag. 1. tab. 7.

Hermannia triphylla Lin. Am. C. p. 91. Spec. plant. p. 942.

Habitat in Cap. b. Spei

Caulis decumbens pilosus, ramosis. folia pilis sparsis hirsuta.
foliola sessilia unguis gularum. Stipula ovata. Flores ternales
subcapitati 3-5. Thunb.

Diversa ab Hermannia triph. Cavan.
non vidi.

Subus P foliis ternatis foliolo oblongo utrinque venosis basi
attenuatis

Habitat. in Ind. Orientali. China 5

Rami teretes quibus puncta. Solia alterna petiolata.

Foliola oblonga utrinque attenuata, prope basin basi, cetera
gemina obtusa utrinque venisissime venosa cetera medi-
um tri-pollicaria. ^{ut supra minus} Petiolis supra planis subtus connatis.
Paniculo axillari et terminali. Stamina 5que
connata.

an hujus generis

pinnatus Connarus folius Vernati ^{quina} pinnatus que, natali basi leptoph
Caron Desf. p. 375 tab 222
Connarus
quinatus, folioli oblongo ovalibus floribus & gynoio la
Mark Desf. encycl.
Habitat in India orientali &c.
Tota glabra Calce

monocarpus
non forte eadem ac ~~monocarpus~~

Santaloides Connans foliis pennatis: foliolo ovatis acuminatis, pedunculis axillaribus aggregatis, floribus racemosis paniculatis
Santaloides Desf. Fl. Reyl. n. 408.

Habitat in India orientali &

Arbor ramis teretibus, glabris, cortice fuso-purpureo-florescente.

Folia petiolata, parva, remota, pennata cum impari, bi-
quadrijuga. ^{subopposita,} foliola ovata, petiolata, ^{superiora} ~~superiora~~ ^{intermedium} ~~intermedium~~

integrum may

aequilongia, aequalia, ~~acuminata~~ ^{integrum}, acumine obtuso,
glaberrima, utrinque venoso-reticulata; venis parvis
protuberantibus, supra nitida; ~~integrum~~

Petoli pariales, vix semiunguiculares, supra canaliculati

†
Peduncululi axillares, 4-6, foliis dimidio breviores, glabris, par-
tiales parvis, versus apicem flores 3-4, pedicellati.

Calyx ^{ovatus} quinquefidus ^{partibus} ~~intermedium~~ ⁶ ~~fidus~~: laciniis lanceolatis, acutis,
in eodem sensu campanulatis quinquefidis ~~intermedium~~ ⁶

Petala quinque, calyce duplo longiora.

Fructus unguicularis glaber

Stamina si filamenta 10, inaequalia, longitudine corollae,

stylis ^{basis} ~~unicis~~ ^{conata}

capsula unguicularis, glabra

Brachycarpa ~~Cecome~~
Floribus ~~Arabianis~~ foliis omnibus ternatis obovatis, piliquis
oblongis scabris, fementibus glabris
Gleome ornithopodioides Desf.

Habitat in Arabia

Tota planta ad lentem ~~caulis~~ ~~teretis~~ ~~ramosus~~ ~~teris~~ ~~gle~~ ~~scabriusculis~~ ~~suffruticulis~~ ~~semiter-~~
adpressa ~~ramis~~ ~~subrosis~~ ~~ad~~ ~~lentem~~ ~~caulis~~ ~~per~~ ~~ramis~~ ~~elevatis~~
elevatis ~~ramis~~ ~~non~~ ~~scabris~~
solis petiolata ~~goleola~~ ~~foliis~~ ~~obovatis~~ ~~parvis~~ ~~sex~~ ~~femur~~
quiculana, ~~hatis~~ ~~scabris~~ ~~petiolata~~. ad lentem punctata

Petiolus viscosus.

Pedunculi axillares uniflori glabri vel subviscosi capella
subcapellares - unguiculari.

~~Aliguo~~ ~~in~~ ~~calyce~~ ~~viscoso~~ ~~vis~~ ~~unquiculari~~ ~~obtusiusculo~~
viscoso ~~scabro~~ ~~stilo~~ ~~terminato~~. Semina numerosa, fusca,
glabra, ~~laxa~~
minus viscosa ~~et~~ quam arabica

memo fides Conarus foliis pennatis subdecemjugis: foliolis ovatis oblongis emarginatis,
racemis axillaribus.

Habitat in Insulis Nicobanicis ad Id. Front 5.

Rami teretes, superne villosi.

Folia versis apices ramorum, petiolata, alterna, approxumata, pin-
nata cum impari, 9-11juga. Foliola brevissime petiolata, opposita
vel alterna, unguiculata, interiora minor, glabra, ^{hirsuta} tenuissime
~~vellosa~~ venosa, obtusa, profunde emarginata, subtus pallidiora.
Petioli leviter villosi.

Peduncululi ~~laeviter villosi~~

Peduncululi axillares, plerumque terni, longitudine foliorum, sub
canescentes. Pedicelli parvi siliiformes, femiunguiculares,
infimi apice saepe triflori. Bractea minuta ad basin singuli pedi-
celli. Flores parvi

Calyx Corolla Stamina, Pistillum utraque procedente