

Conocarpus

1. erecta C. foetida, foliis lato lanceolatis cylindricis
2. procumbens C. procumbens, foliis obovatis lanceolatisve basi biglandulosis
- + 4. racemosa C. foliis ovalibus oppositis mucronatis, fructibus segregatis
- + 3. articulatus C. foliis ovatis oppositis acuminatis, capitulis axillaribus globosis solitariis

4 racemosa C. foliis lanceolatis-ovatis obtusiusculis, fructibus segregatis Lin.

Conocarpus floribus completis remotis Jacq. am. 100. t. 53.

Mangle pilifera, foliis ellipticis ex adverso nascentibus Koun-hist. 2. p. 66. t. 107. fig. 1.

Habitat in Jamaica maritimis

Ramifera, glabra, purpurascens. Cilia petiolata opposita ovata
coracea integerrima ^{utrinque glaberrima} ~~anthera~~ ^{et} ~~ob~~ ^{subtus} ~~obscure~~ ^{obscure} venosa, ver-
sum marginem in anostomosis puncta nonnulla parva, bi/tri-
policaria, mucronata interdum retusa. Flores distincti spicati
speciosi axillares et terminales ~~et~~ ^{interdum} terni pedunculati
terminali inferne ramosi, canis. bractea ovata decidua.

pedunculati compressi floris decidui pentapetali
fructus oblongus basi parva ad emarginatam angustam glaberrimam unicus mucronis et
superiore alveola dilatata apice calyce tryncato mucronato

racemosa C. foliis ovalibus, oppositis mucronatis fructibus segregatis

Stamina decem proprii generis est Richardus sphenocarpa

an hujus generis

procumbens *Conocarpus procumbens* foliis obovatis Haëgu. Amer. 79. 8. 52. Fig. 2.
-P. V. p. 217.

Budecthia fujina, foliis subrotundis Cunn. herb. 501.

Habitat in Cuba.

An varietas *C. erecta*?

Rami cuneatis glabri: Ramuli compressi angulati: Folia
alterna petiolata acuminata, coracina avenia basi
in petiolum deprehensa, bi- & trijollucana. Capitula
racemosa, subglobosa. Pedunculus flexuosus.

procumbens *C. foliis lanceolatis obovatis lanceolatisve basi longioribus*
obtusis

ab erecto specie non differet. Richard interdem humilis rami
decumbentibus et prostratis glandula minute in axillis nervorum
subtus
an modo varietas

erecta & erecta foliis lanceolatis l. v.

Gayu. amer. 70. v. 52. fig. 1.

1. Conocarpus manghana arbor curasavica, foliis foliis Catech. cur. 2. p. 33
t. 33.

Alnus mantima myrtifolia conarionum Plukh. alm. 10. v. 240. fig. 3.

Alni fructu laurifolia arbor mantima Sloan hist. 2. p. 10. v. 161. fig. 2.

Innomerata Plum. ic. 135. v. 144. fig. 2.

Habitat in Jamaica, Bermudensium, Brasiliae mantimis

Rami ^{glabri} inferne teretes, superne angulato compressi cum angulo acuto
utroque decurrente. Folia petiolata aeterna, triplicia
nervi, subcoriacea obscure venosa attenuata utroque attenuata
integerrima glabra. Pedunculi axillares, solitarii, terni
interdum bini pedunculati, pedunculo compresso glabri apice
diphylli: pedicelli tomentosi, et terminales plerumque rari
capitula oblonga tomentosa. Flores pentandri pentapetali

erecta & erecta foliis lato lanceolatis eglandulosis.

glandulas huius etiam adfert Sw. in obs. an eius erecta
Gayu. procumbens?

~~Articulatus~~
3 articulatus

C. foliis ovatis opproptis, capitulis axillaribus globosis solitariis

Habitat in India orientali

a ~~stipite~~ utrinque

Rami ~~obscure~~ verticillati, glabri articulati, ^{sub articulis} dilatati
compressi. Folia petiolata, ovata, ^{subcoriacea}, integerrima
oblique alternata ^{versuta}, ^{secundum nervos} ^{venosa}, ^{lineata} glaberrima ^{acuminata} ^{longa}, fere
palmata. Petiolus brevis. Stipula ovata ^{parum} attenuata
capitula pedunculata magnitudine ^{pediculi} ^{magis} ^{veluti} ^{conyli}. Calyx
superus 5fidus pubescens: laciniis linearibus lanceolatis.
Corolla 5petala: petalis oblongis fere longitudine calycis
filamentis nulla anthera oblonga s. longitudine corollae
stylus longitudine filamentum superne crassior

sub nomine Cenchona missa a Monzyo