

Etopis

~~Polypodiaceae~~ <sup>Cato</sup>  
5 ~~Coryzoides~~ *Eriogon* aborea foliis lanceolatis serratis supra scabris, subtus reticulato rugo  
amygdalena *Coryzoides* floribus corymbosis

habitat in Ampela Borussia  
aboreis Pauli Rami tenete villosi

folia, petiolata alterna, lanceolata, serrata, obtusa cum mucro  
ne supra scabris prope in retrosum pilis vix congruuis puncto minutis  
fimo infidentes, subtus villosa vix in die mollesca, rugo reticulato rugosa  
ut in sa lvia, bi pollicaria Pedunculis

Corymbi terminales multiflori villosi scabris

Calycis dors. villosi margine membranacea

Corollula radii tubulosa

Pappus ut in reliquis

differt a molli serratis a digitibus minoribus apice cartilagineis  
foliis subtus reticulatis minime fereis

*Coryza amygdalena* foliis petiolatis ovato-lanceolatis  
serratis subtomentosis, floribus corymbosis La Mant  
line

forte potius *Coryza* cum Corollula radii tubulosa

elliptica Lam. foliis ellipticis <sup>integerrimis</sup> subtus villosi subtomentosis  
corymbis terminalibus compohtis patentissimis

Habitat in Senegal Desfontaine 5

Frax. Baucher. arborea. Rami teretes ~~vel~~ subtomentosi  
sternuosis ~~vel~~ folia petiolata supra glabra ovata  
subtus venoso reticulatis magna. Pedunculi est aqua  
ramorum parvae: sparsi <sup>tenues</sup> patentissimi tomentosi ramis  
patentissimis bi- trifloris floribus pedicellatis. Calyx  
hemisphaerici imbricati villosi squama exteriori  
ovata acuta. Pappus flavescens

populifolia Conyza incana, foliis ovatis integerrimis petiolatis  
corymbis sem pluribus.

Conyza populifolia foliis cordatis petiolatis tomentosis  
integerrimis, calycibus hermaphroditiis La Mart. Cony-  
zop. pag.

Habitat in tupa la Franciso

Rami uti reliqua partes ~~incana~~ tomentoso incana.  
Tolia sparsa fu bipollicaria supra nervosa subtus albi-  
dura venosa. Petioli subtus convexi supra cana-  
liculati. Corymbis terminalis cum ramulis floriferis  
inferiore sa corymbo rami altiore florifero ~~pedunculato~~  
adsumi squamis parvis adpressis linearibus acutis  
Calyx <sup>hermaphroditicus</sup> imbricatus squamis lanceolatis tomentosis  
petulis. Pappus setis interioribus Conyzei unctis setis  
sublanicolatis brevioribus exterioribus multo  
longioribus omnibus hermaphroditis an proprie  
genens

visida

~~glauca~~

Corymb foliis ellipticis glaberrimis Venerariis integerrimis  
acuminatis.

Habitat in Brasilia  
<sup>arborescens</sup>

Rami angulati glutinosi glabri folia petiolata albena  
basi angulata <sup>ramis oppositis</sup> ~~acuta~~ obsolete venosa nervi tres Venae  
proter marginales plerumque integerrima interdum uni  
cum versus apicem marginis albenis dentibus duobus  
remotissimis, acumine brevi ~~et mucronato~~ <sup>et mucronato</sup> ter minute  
bis. 3 pollinaria pedunculata ex alis fissuris et  
apice fastigiati erecti. Flores <sup>parvi</sup> in corymbo decorymbo  
fitti. pedunculis, pedicellisque glabris angulatis, squa  
mula lineari & basi suffultis. Squama calycina  
lineari lanceolata glabra. Pappus rufoescens.

27 ~~Carinata~~ <sup>longe</sup> ~~longe~~ <sup>oblongis</sup> ~~oblongis~~ <sup>resplendens</sup> ~~resplendens~~  
punctatam ~~longe~~ <sup>foliis</sup> ~~foliis~~ <sup>ellipticis</sup> ~~ellipticis~~ <sup>punctatis</sup> ~~punctatis~~ <sup>longe</sup> ~~longe~~ <sup>integerrimis</sup> ~~integerrimis~~ <sup>terribus</sup> ~~terribus~~, calycibus <sup>disi-</sup> ~~disi-~~  
superantibus <sup>sub</sup> ~~sub~~ <sup>pano</sup> ~~pano  
Habitat in madagascia~~

Rami lignosi <sup>fronati</sup> ~~fronati~~ glaberrimi uti tota planta. <sup>foliis</sup> ~~foliis~~ <sup>elliptica</sup> ~~elliptica,  
integerrimo, <sup>simplici</sup> ~~simplici~~ <sup>ter</sup> ~~ter <sup>venosa</sup> ~~venosa, utrinque punctata <sup>superior</sup> ~~superior~~ <sup>sequi</sup> ~~sequi  
polliciana <sup>supremo</sup> ~~supremo~~ <sup>acuta</sup> ~~acuta <sup>inferior</sup> ~~inferior~~ <sup>obovata</sup> ~~obovata <sup>subretusa</sup> ~~subretusa. <sup>mem</sup> ~~mem  
in axillis <sup>rigidis</sup> ~~rigidis~~. Corymbi <sup>terminalis</sup> ~~terminalis <sup>pauciflori</sup> ~~pauciflori <sup>folioli</sup> ~~folioli <sup>floralibus</sup> ~~floralibus ad  
basin pedicellorum linearis lanceolatis. Calyces <sup>subglobosi</sup> ~~subglobosi~~; folioli  
subaequalibus <sup>se</sup> ~~se <sup>anosis</sup> ~~anosis <sup>nitentibus</sup> ~~nitentibus, acutis, <sup>longis</sup> ~~longis~~ <sup>disi-</sup> ~~disi-  
superantibus.  
Corollula <sup>radii</sup> ~~radii~~ linearis. Pappus <sup>purpurascens</sup> ~~purpurascens <sup>pelosus</sup> ~~pelosus~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~

~~adhaec forte inter Penya~~

10 <sup>291</sup> oppositifolia <sup>E. Gray</sup> ~~Eryron~~ foliis ovatis acuminatis integerrimis trinerviis, corymbis Verminalibus  
Habitat in Draplica monte vide Thonin.

Caetera ramis teretibus ~~pubes~~ glabris striatis. Folia opposita, petiolata, ovata  
acuminata, trinerviis, nervi versus apicem deliquescentes; utrinque glabra  
ferri pollicaria. Ramuli floriferi ex axillis supremis longitudine foliorum  
foliis minoribus. Pedunculi oppositi multiflori, laxis angulati, floris Vermis  
flos hemisphaericus, Verminalis primum florens. Calyces glabri squamis  
oblongis obtusis fuscis margine pallidis. Pappus ferrugineus calyce duplo longior  
vilosis. Flosculi plurimi nudi setacei.

Genus dubium ob foliis oppositis et corollis radii capillaribus idcirco forte  
Eryron

oppositifolium L. f. g. foliis ovatis



*pyrifolia*

*Coryza* foliis <sup>ovatis</sup> linear-lanceolatis, acuminatis subdentatis  
hi glabri cymbo paniculis  $\frac{1}{2}$  laxo. Ca. Mart. Cuv.

May

Habitat in Hava  $\frac{1}{2}$

Stami glabri, flores albi, calycis glabri. Pappus ferrugi-  
neus

non vidi



inerva *P. solus oratis subnodatis pulso v. fidi dentatis basi auricu-*  
*latis, receptaculi d. sic fawolo Hort. Keiv. 3. p. 104*  
*Habitat in Cap. b. nei 7*

non vid.

Scorpioides Conyza

Conyza scorpioides foliis ovato-lanceolatis serratis <sup>et petio-</sup>  
latis integris lobatis, spicis secundis revolutis nudis La-  
marck Encycl.

Habitat in Brasilia &

Descript quere inter amara.

<sup>foliorum</sup>  
Sessilis *Coryza* foliis ellipticis utraque acutis glabris speciosis axillaribus  
ternis vel quatuor secundis foliis secundis  
Habitat in Montserrat Pagan

Descriptum inter Americis inventum

nervosa

Com. foliis ovatis nervosis glabris. spicis ramis secundis  
foliis

Habitat in Anfula N. Carolinae West

Rami suavi tomentosi nervi obliqui subtus inani  
Flores papilionacei

inter americanas Descript.

nyda Con: folio petiolati, ovato integris p. latis subtus venosis  
p. flexuosi, flonibus secundis geminatis ~~fl. latis~~  
s. p. 104 Sw. p. 104 pag. 113

Habitat in ~~Inf. St. Helena~~ Tomaria 5 subrotundo  
Caulis et rami teretes incani tomentosi. folia ovata  
glab. et integerrima venoso pollicaria <sup>p. latis</sup> ~~v. latis~~  
laterales pedunculis flexuosis calycis gemini sepales  
tubinati glab. valde imbricati squamis ex venis  
ovatis interonibus lanceolatis

ovatis integerrimis

уездн. Конста



arborescens *Coryza* foliis <sup>ovatis</sup> lanceolatis integerrimis subtus tomentosis, racemis axillari  
secundis, bracteis reflexis. Lin. S. V. pag. Sw. obj. pres

*Coryza arborescens*, floribus coeruleis Plum. spec. 10. 4. tab

foliis elliptico lanceolata sub  
130. fig. 2.

*Coryza fruticosa*, flore pallido purpureo, capitulis e late  
nibus ramulorum spicatum exsertibus Roux hist. jam.  
pallida non tomentosa  
in spec. herb. 4. usq. 124. herb. 1. p. 257.

Habitat in America meridionali

Tridax ramis depressis, scaber. Rami divaricati divergentes  
Tolice interdum herpetice nigricante villosi. Folia petiolata lato-lanceo  
subtus glabra herb. la  
la masoi  
hirsuta. Racemi terminales et axillares erecti floribus  
subsessilibus alternis squame calycinis pubescentes  
Swartz

Obj. Folia cordato-ovata habet Plumieris figura sic haud  
exacte respondet ad ~~differentiam~~. ad Desimp  
Prami in mea angulata Swartzii. cum descriptione ha Manthii cum specimen  
*Coryza* a Prohrio foliis nigrosi muso nec abluat a Swartz.  
nisi fig. foliorum et quad rami tomento in uno. nec nigricant.  
+ etc.

*Indicofa Conyza* foliis ovatis integerrimis obtusis, floribus sessilibus alternis,  
ramulis flexuosis L. n. 1. V

~~*Eupatorium americanum*~~

Habitat in America meridionali

mibi haud nota

altissima Achyranthus

Tortuosa Con. foliis ovato-oblongis integerrimis, caule tortuoso  
fruticoso, racemis reflexis Hort. Cliff. 405 & Lin Syst. V.  
Conyza madagascariensis fruticosa tortuosa caule, corni plis  
Vaillant. Act. parif. 1719. pag. 396.

Habitat in Madagascaria et sera Guineæ  
rami subherulati, simpliciter. Folia in ramis inferioribus obtuse  
ovata, superioribus ovato-lanceolata, integerrima venosa, duntaxat  
scutellari, pollicari. Racemi terminaliter. Verticillati. In  
alio supremis simpliciter. Flores alterni secundari. Squamæ  
calycis ovatae Lin

mihi ignota

bracteata *Coryza foliis lanceolatis lobis puberulis* sp. n.  
axillaribus terminalibusque secundis foliis  
Habitat in Tripula St. Crucis Ror. west. 5  
Rami albo tomentosi: flores inferiores gemini & levis  
solitarii

reliqua inveniuntur in d. sp.  
americanas

flectuosa C. foliis canicolas nigosis <sup>puberulis</sup> subtus ~~allopis~~ ~~subtamen~~ ~~topis~~  
specis axillaris terminalibusque secundis  
Habitat in Insula St. Crucis  
- forte non haesi varietas C. bracteata

13) ~~Platanus~~ <sup>argentea</sup> Conyza <sup>tormentosa</sup> ~~nitida~~ sericeo  
Eriogonum glandulosa foliis oblongis amplexicaulibus, ~~nitida~~ sericeo  
subidentibus floribus terminalibus sessilibus congestis

Habitat in Insula Bourbonia.  
Caulis fruticosus Rami uti tota caeteri pilis densis sericeo nitentibus

Folia oblonga amplexicaulia, imbricata, semipollucaria, integerrima, obscure lanuginosa nervi apicem non attingentes.  
Flores aliquot subsessiles in apice magnitudine Eng: acris

Calycis squamae carceolatae.  
Flores radii nudi

Pappus ferrugineus calyce longior  
Planta eleyana.

Conyza argentea foliis ovatis tomentosis sericis semianplexicaulis, floribus sessilibus terminalibus congestis  
de Martii Encycl. pag.

9 elongata <sup>Conyza</sup> Engelm suffruticosa foliis oblongis serratis <sup>ligabris</sup> Pedunculis elongatis  
simplicissimis unifloris terminalibus solitariis unifloris  
hab. in Isla Bourbon

caulis rami tenues teretes villosi

folia alterna, sessilia oblonga basi angustata, dent serrata serris  
tunc utrinque 3-4 remotis apice dente incurvato nudo,  
villosa scabra unguiculata villosa scabriusculis

Pedunculus terminalis solitarius uniflorus quadripollicaris  
nisi foliis 3bus & remotissimis, subovatis super dentatis supremo  
integerrimo

flor longioris atri

capillares

Corollula radii tubulosa stylis duplo brevioribus numerosis

Pappus albidus

Trochlea Calycina linearia



1  
Laurifolia  
nerifolium  
Coryza  
Euphorbia arborea foliis integris lanceolatis <sup>extremum latioribus</sup> integerrimis serratisque  
floribus corymbosis

habitat in Anful: Bobonia 5

Caulis arboris tamen teretes laves fusci.

Folia alterna, lanceolata figura nervi oleandri basi in petiolum  
approximate attenuati inferna supra serrata, superiora integerrima, glabra  
venosa tripollicaria

Petioliis ample basi dilatatis

Corymbi terminales multiflori, Pedunculi suffulti bractea  
linear

Calycis hemisphaerici. Squamis oblongis <sup>glabris</sup> margine aridis glabris

Corollula radii lineares numerosi

Pappus ~~ferrugineus~~ purpurascens longitudine calycis

Coryza laurifolia foliis oblongis lanceolatis basi in  
petiolum angustatis subintegerrimis corymbis terminali  
composito La Marti. Encycl.

B. folius extremum latioribus

ovata *Coryza* foliis ovatis serratis subtomentosis acutis, caule corymboso  
caulis subglobosis Lin. S. V. undulato

*Coryza arborescens* purpurea, folio verbosio dentato  
Plum. ic. 97.

12 *Eupatoria coryzoides* madagascariensis, foliis glabris  
flores purpureas Pluken. alm. 141. T. 177. fig. 2  
Habitat in America meridionali

*Coryza major* odorat. f. Bauh. pin. floribus purpureis nudis  
Sloan hist. jam. 1. pag. 258. tab. 152. fig. 16

13. foliis dentatis magis virentibus. L. Mantk. Em.  
*Coryza arborescens* purpurea, folio verbosio dentato Plum.  
Spec. 9. Busch. 4c. 97. ovato

Rami tomentosi foliis ovato-lanceolatis, denticulatis  
3-4 pollicibus subtus leviter tomentosis. Calycis  
villosi squame oblongo obtuso

Pluken. ex herbis sancti

vesta

Tridax 3-6 pedalis. Calyx imbricatus squamis profundis linearibus. Corolla flosculi  
foeminei, numerosissimi, & paucissimi in centro. & infundibuliformis, squedentatis, dentibus  
patentibus sculis, acutis. Stamina filamenta 5. limbo breviora. Tubus antherarum corolla  
longior, apice 5que dentatus. Dextriculis hiantibus. Germen O. Stylus capillaris, basi tumi-  
ditus. Stigma simplex, teres, stylo vix crassior, apice pubescentum, glandulis cor. & persum. Pappus  
pilosus denticulatus, corollula parum brevior  
♀ Corollula utraque, ad gradum, tubo capillari, superae vix latior, & dentata  
dentibus recurvis. Stamina O. germen oblongum, villosum. Stylus corolla longior  
Stigmata 2. fetacea, erecta, villosa. Pappus utraque. Pterostachyum medio  
depressum Richard

appendiculata <sup>lato</sup> Coryza foliis lanceolatis serratis subtus tomentosis  
petiolis dentatis

Coryza appendiculata foliis lanceolatis serratis subtus  
tomentosis basi appendiculatis La Martk Enigyl  
neg.

Habitat in insula Bourbonica

Rami teretes e caespitibus foliorum subtulati superne  
tomentoso incani. Folia <sup>basi attenuata</sup> ~~fracta~~ frequentia tri vel  
quadripolliciana acuta grosse serrata. Petiolus  
~~bas~~ semipollicaris utrinque terminis 3 bus foliaceis  
lineantibus parvis appendiculatis. Corymbus ternarius  
laxus, erectus, compositus. Pedunculi elongati utriusque calycis  
tomentoso incani. <sup>ultimi</sup> Pedicellum lineare ad basin pedun-  
culorum. Pedicelli 3. ~~asse~~ <sup>asse</sup> unguibus adpressi foliis sita  
caes intus glabri

Balsamifera  
~~lanicollata~~

C. foliis lanceolatis serratis subtus reticulato  
rugosis allopis, petiolis acutis

Habitat in Java

Rami teretes superne tenuiter tomentosi. Folia petio-  
lata, <sup>superne</sup> ~~superne~~ <sup>subnatis</sup> ~~superne~~ lanceolata <sup>superne</sup> 4-5 polli-  
cari <sup>superne</sup> ~~superne~~ angustiora. Pedunculi et axillis <sup>superne</sup>  
buis frequentis solitarii folio longioris. Pedunculi <sup>superne</sup>  
approximati subumbellati uni bi, 3 flori. Pappus ser-  
reus

Congra odorata Rumph. amb. 6. Tab. 24. fig. 1.

Congra arbor zeylanensis subtorrido foliis maxime  
tomentoso Pluken. alm. 64

Congra balsamifera foliis lanceolis subtus tomento  
reticulisque dentatis serratis Lin. S. V.

subcordata P. inermis foliis cordato-subrotundis, cunctibus peltis sub  
elevatis s gonio, angulis apice mucatis Sw. prod r. pag 60  
Habitat in India occidentali 5

~~Calyptro~~  
~~dentata~~

~~Coryza folio ~~laciniato~~ <sup>oblongo. cuneiformibus sublinearibus inaequaliter</sup> ~~subtus tomentoso~~ <sup>petiolis</sup> ~~dentato~~ <sup>dentato</sup> ~~lanceato~~  
~~hinc s. v. petioli dentatis~~~~

~~Coryza odorata Rumph. Amb. 6 tab 24 fig 1~~

~~Coryza odorata Zeyla densis subrotunda folio maxime  
tomentoso folio Plukh alm. 64.~~

~~Habitat in India orientali et Arabia~~

~~Rami teretes villosissimi folia frequentius, alterum~~

~~oblonga basi attenuata supra fere nuda subtus villosa~~

~~longis densis tomentoso subspicua, inaequaliter serrata~~

~~petioli a folio decurrenate alati ala denticulata~~

~~Corymbi terminales compositi~~

~~verecor quod Synonymi Rumph. petius ad appendiculata  
la Mantia petio~~

4 ~~glutinosa~~ ~~tenax~~ ~~Engusa~~ <sup>Coryza</sup> Caule fruticoso superne glutinoso foliis lanceolatis <sup>ovatis</sup> petiolatis  
ferratis floribus corymbosis  
~~Patriam ubi ignota est horti Parisiensi~~ Habitat in Europa fronsis

Caulis fruticosus quadrifidus <sup>petalis</sup> <sup>venosa</sup>  
Folia petiolata alterna, glabra <sup>venosa</sup> <sup>venosa</sup>, serrata lanceolata  
pabentia bipollicaria — Petioli breves  
Corymbi terminales multiflori. Pedunculis squama una alterave lin  
cari glutinosi

lanceolatis obtusis  
Calyces subglobosi parvi, folioli membranaceis nitidis, squamis calycinis  
Corollis flavis

Pappus albus.  
Corolla radii nudi.

Coryza glutinosa foliis lanceolatis serratis utroque induribus  
junioribus crispis La Martie Encycl. pag.



Juaveolens C. foliis oblongis subferratis <sup>asperatis</sup> glabris corymbis Verminalibus  
decompositis

Coryza odorata

Coryza

Habitat in Arabia

Russii Yanicae superae pulverulenta foliis acuta basi non  
angustata calycibus glabris

Francis Conroy

Carolinensis. P. foliis lanceolato-oblongis integerrimis, corymbis axilla-  
ribus, pedunculis multifloris

*Coryza Carolinensis* Jacq. coll. 2. p. 271.

Habitata Carolina

Arbutifera sp. pedalis. Caulis erectus, glaber, diametro pollicari. Rami ter-  
tes pures. Tomentosi rari. Folia alterna, lanceolata-oblonga cum  
nerviis, integerrima, petiolata, utraque tomentosa mollia, v. sicidula 4-  
6 pollicaria. Ramuli floentes breves, oppositifolii, pedunculati in ramulis  
axillares multiflori flores corymbosi. Calyx tomentosus cylindricus  
imbriatus foliolis lanceolatis, flosculi feminei limbo trifido hermaphrod. Jacq.

non vidi

Ponyza

est in herb. specim. gnyza haud perfectum a Töpfl.

Boichani foetida in herb. h. Ponyza foetida la

ad Pongrom

Ponyza purpurea la March. Stobelina speciosa h. gna