

12 Corispermum

13

Cal. 0. Petala 2. Sem. 1. ovale, nudum

Digynia

12. Corispermum Cal. 2-phyllus. Cor. 0. Sem. 1. ovale, convexo-planum,
nudum.

orientale Conispermum floribus axillaribus solitariis pedunculatis, foliis linearibus
energis inermibus

Conispermum orientale; foliis linearibus angustis, summitatibus flori-
feris supaniculatis pubescentibus La Mart. illustr. t. p. 15.

Habitat in Oriente

Caule erecto, ~~herbaceo, ramis~~, cortice viridi canescente, superne
albido lineatus, oculo armato villosus. Folia bipollicaria, basi
sensu sensim angustata, glabra, obtusiuscula, crassa. Flores ~~versus~~
apices ramorum, alterni, pedunculati, folio florali duplo pediculi
folio florali linearis duplo brevioris.

Clavata *S. Intra* articuli emarginato bifido, spica Vermineali
bus
Habitat in India orientali 5 ^{somali}
Caulis teres Rami oppositi articuli femineunguicula
res superne latiore. Spica pollicaris

hyssopifolium? Conispermum species terminalibus, floribus distantibus, foliis ^{enerviatis}
~~et~~ praeterea inermibus

Conispermum hyssopifolium Fuss. in Act. par. 1712. p. 244.

Conispermum glabrum, majus, longis dimidiatis primis spec. Am man.
rath. ~~primis spec.~~ p. 22 s. prima spec. p. 150.

Conispermum floribus lateralibus Flin. Hort. Ups. 2.

Conispermum hyssopifolium Pall. fl. ross. 2. p. 112. tab. 90. fig. E.

Habitat ad Wolgam Tartaria: Gyllau Russia, Monspelie, Loricaria
novis O.

Caules semipedales, teretes, saepe flexuosi, pallide lineati, glabri, ste-
ta saepe purpurascens, ~~rami~~ Rami alterni, patentem, ramulose. Colla
sesquialtera, bipollicaria vel parum ultra, acutiusculo, minime mucrona-
ta, crassiuscula. Bractea lanceolata, margine membranacea. Semina
orbiculata, membrana cincta, bractea breviori

Alisum vidi ~~fructu~~ bracteis ovatis acuminatis minoribus, f. de tantu
modo variatis ^{planta}

Quadrac.

Hall: stamina Flin. Culta vidi. Pallas 29 in Spontanea. ^{Quo seltno}

Hort. Ups. Hort. Ups. 2. in Culta Tab. 4 Hall. ed. h.

erecta B. caule erecto glabro, floribus diandris Lin. b.V. p. 524
 Boerhaavia floribus paniculatis, seminibus pentagonis truncatis Lin
 Mant. 315. Jacq. hort. 1. tab. 5. 6.
 Boerhaavia diandra glabra & effusa, floribus diandris Lin. Princ.
 p. 44 ~~1. 2. 3. 4. 5. 6. 7. 8. 9. 10. 11. 12.~~
 Habitat in ~~India~~ Vera-Cruce Americae 4

erecta B. caule tetragono glabro internodiis viscosi; floribus
 paniculatis fructibus tubinato clavatis muticis
 Boerhaavia erecta foliis ovatis acutis subundulatis glabris
 caule erecto, fructibus tubinato clavatis muticis Sw.

fructu truncato:
 Specie forte non
 distinguenda differt
 ab erecta semina
 erant laevia nec
 constricta puncta

off. pag 10
 Caulis 1-2 pedalis erectus ramis ad petiolum immixtis
 folia basi ovata acuminata subangulata margine
 undulata purpurascencia subtus alba petioli pube
 sentes. Pedunculi in axillis foliorum ter minalem
 natiuales 3 flori. Filamenta duo. Semen pentagonum
 globosum

Squarrosus ³ Corispermum species axillaribus floribus confertis imbricatis, foliis
~~lanceolatis~~ ^{enerviis} bracteis que mucronatis subpungentibus.

~~Rhagotis~~ Rhagotis semine Pastinaca Buxbaumia 3. p. 30. T. 55.

Corispermum Amman. ruth. no 225. spec. 2da p. 159.

Corispermum Squarrosum species squarrosis Lin. Hort. Ups. 3.

Corispermum hyssoifolium Pall. fl. Ross. 12. p. 412. T. 98. f. A. B. D.

Habitat ad ripas fluviorum Russia et Sibiria ☉

Caulis ramosus, terre, albedo lineatus, superne villosus. Folia angustiora
quam in hyssoifolio, at longiora. Spica angustilobis, primum minus

res ^{sub} resiliter, dein pedunculato, villosa. Flores in finibus parum remoti,
reliqui conferti. Bractea infimo linearilanceolata, superiore ovata

lanceolata, apice potentes, marginem membranaceo, ubi folia mu-
crona brevi flavescente terminata.

Stamina quinque numeravit Pallas.

Campestri Collectio Hort. Kew. 1. p. 5. *

Habitat ad P. l. p. i. ^{inferne}

Caules teretes, ~~stomente raro~~ pubescentes, ramosi. Folia
verticillata, ^{versus unum} ~~versus unum~~ ^{latus} ~~latus~~ ^{vergentia}, lineari-lanceola
ta acuta ⁱⁿ ~~in~~ ^{termina} ~~termina~~ ^{glabra} ~~glabra~~ ^{vsq. uncialia}.

^{in mucron} ~~termina~~ ^{plurima} ~~plurima~~ ^{membranacea} ~~membranacea~~ ^{lanicolata} ~~lanicolata~~ ^{perpetua} ~~perpetua~~ ⁱⁿ ~~in~~ ^{capitulum} ~~capitulum~~ ^{axillas} ~~axillas~~ ^{trium} ~~trium~~
+ ^{vel 4.} ~~vel 4.~~ ^{foliorum} ~~foliorum~~ ^{occupari} ~~occupari~~ H. Kew.

purgens ⁴ Conispermum floribus axillaribus aggregatis; foliis linearibus lanceolatis
nervosis pungentibus, bracteis subulatis pungentibus
Aegrotis foliis arundinaceis Buxb. cent. 3. p. 30. v. 56.

Conispermum squarrosum; spicis axillaribus squarrosis, foliis lanceolatis
acuminatis pungentibus nervosis Pall. fl. ross. 2. p. 113. t. 99.

Habitat in arenosis aridissimis deserti Romani et Cappici.

Caulis erectus, teres, striatus: striis albidus, villosus, proscithum superius
apice puberulescens, ramosus: Rami alterni, caule simillimi. folia
sessilia, distantia, bipollicaria, patentissima, linearibus ^{lanceolata} subfalcata,
glabra, nervosa, striata, in spinam pungentem flavescentem atti-
nenda. Spica axillaris, brevissima glomerata in axillis folio-
rum aggregata, inter distincti bracteis subulatis, rigidis, pun-
gentibus. Reh.

~~Boiss.~~ Stylis mucro bifidus: Lemnae haud marginatae Pall. l. c.

Habitus Caulis herbaceus, ramosus: Rami alterni. folia sessilia, linearia,
integerrima.

Lemnae haud marginatae Pall. l. c.

niger. Coni per mum.

Coni per mum squarrosium.