

Cosignia

Triphylla Cospignia

Habitat in Insula Franca Combeson herb. Guss. 3

Rami superne compressi, pulverulento-tomentosi, uti petioli, pedunculati; Calycis et capsulae. Sola petiolata, ternata: foliola petiolata, 3-4 pollicaria subincurva, basi acuta, obtusissima, integerrima, serosa, supra nitida, subtus tomentosa, uncinata. Racemi compositi. Fructus semipollicari

pentaphylla *Cosynia* Tus: gen. pag

Habitat in Insula *Sancti* Tus: 5

Rami teretes flavescenti tomentosi, foliis deciduis tuberculatis.

Folia petiolata alterna versus apicem ramorum approximata
pinnata cum imparibus foliis opposita, duo interiora

minora subsessile, tria ^{terminalia} in sequentia sessile terminalia

reliqua majora petiolata ^{et} omnia lanceolata basi acuta, apice
obtusius emarginata, integerrima, ^{et} glabra, venosa subtus uniana

Petiolus pubescens. Thyrsus terminalis ^{et} hemisphaericus

Pediculi inferiores conquis axillares duplicato racemosi
racemi superiores simplices. Bractea parva linearis ad basin
pedicellorum. Calyx tomentosus

Affertur summa cum secundo