

Crasula

umbella Crasfula

Crasfula umbella Jacq. Coll. 4 p. 172.

Habitat ad Cap. b. p. 2

Caulis unus vel alter teris carnosus, basi crassitie calami, spithamou erectus  
subpellucidus nitidus simplicissimus. Folia radicalia nulla, caulina  
duo etiam unicum subrotunda, convexula, repandula areola nervia  
carnosa, fragilia supra vicia radius pallidi ad centem punctata  
subtus purpureo rubentia diametro <sup>unq.</sup> triunciali perforata. viciis  
longis, gemini oppositi & yollucar. Pedicelli inferiores oppositi uniflori  
sub ea pedicelli propriis umbellati summi verticillati. Caulis simili  
verticillo terminatus. Flores parvi. Calyx punctatus Petala lanceo  
lata intus albida, extus carnea. Squama necdiferis fusis vanat una  
padem fructificationis interdum addit Jacq.

capitata Craspula foliis globosis-ovatis quadrifariam imbricatis, cyma  
Inflor. floribus sessilibus Cav. ii. et Desf. 1. pag. 50. Tab. C. fig. 2.  
Habitat monti. C

crenulata Crasf. foliis connatis lanceolatis punctato crenulatis, caule foliis  
corymbis decomposito Lin. suppl. 189. l. V. p. 307

Habitat in C. bono speciei

Paulis erectus, simplex & teres. Folia sesqui pollicaria lanceolata attenuata  
margine cartilagineo squamato, ora vaginatum denticulato ciliata et  
latera inter folia adpressa squamis micantibus. Superne divisi. Peduncu-  
li dichotomi ad basin diffusum. Folia binaria subulata, opposita distenta  
ad basin dimorum corymbis. Calicula calycina glabra subulata  
cannata. Petala linearia extorsum parum latiora

alba <sup>connatis vaginantibus</sup> Crasula foliis lanceolatis ciliis cartilagineo ciliatis impunctatis caule simplicissimo.

Crasula alba foliis lanceolatis, oppositis, vaginantibus. Forsk. descript. pag. 60.

Habitat in montosis Arabia

Cauli Crasulae penno arvensis, <sup>fruticosa</sup> ~~palmaris~~ erectus, vaginis foliorum <sup>superne dichotome divisus +</sup> ~~textus~~. Folia <sup>conferta</sup> ~~conferta~~ sesqui pollicaria obtusiuscula plana. <sup>perennalis</sup> ~~perennalis~~ <sup>bracteis</sup> ~~bracteis~~ <sup>subquasi</sup> ~~subquasi~~ <sup>dichotomia,</sup> ~~dichotomia, <sup>pedunculis</sup> ~~pedunculis~~ <sup>squamis</sup> ~~squamis~~ <sup>si</sup> ~~si~~ <sup>pellucidis</sup> ~~pellucidis~~ <sup>retorsum</sup> ~~retorsum~~ <sup>fiat</sup> ~~fiat~~~~

Seduno flambus corymbosus Forsk. fl. arab. pag. no 273 hinc perthe

Scabra *Crasula foliis lanceolatis<sup>5</sup> pubulatis canicollatis connatis scabri castelagines  
ciliatis caule retrosum scabro*

*Crasula mesembryanthemifoliae, foliis longioribus appensis Dill. Elth. 117.*

Tab. 99. fig. 117

*Cotyledon africana frutescens, foliis appensis angustis acuminatis, flore vire  
scente Mart. Pent. 24. t. 24.*

*Crasula scabra foliis oppositis petentibus connatis scabri ciliatis, caule  
trosum scabro Lin. S. V. pag. 304*

Habitat in Cap. b. p. 5

rami et teretes aequae ac folia adpressis squamis albis retrosum apice paten  
tibus folia unguiculata supra connata sub vaginantis supra  
plana puberula convexa patentia, margine squamoso crenulato  
(pubulata) prima terminalis calycis foliola pubulata squamis  
adpressa. Petala longitudine calycis

subt. n. goni connatis papuloso - scabri caule retrosum scabro  
corymbis composito

an mea dicitur a hunc

prunofa Crasula

perforata Craspula foliis connatis perfoliatis ovatis ciliatis remotis, caule erecto  
paniculato Thunb. prod. pag. 58

Habitata cap. b. sp. a cap. foliis annulata <sup>approximat</sup>  
Caulis fruticosus folia oblonga glabra, ~~unquiculata~~ ~~Corimbis~~  
~~trichotomis~~

foliis opposito perfoliatis ovatis ac caule simpliciter floribus  
pedunculato subverticillatis her. Suppl. p. 190  
Caulis simplicissimus teres sesquipedales nudus. folia acuta  
petala margine aequali basi superiore saepe minorum. Racemus  
terminalis longus. flores glomerati per se in pedunculis  
connatis oppositi quasi verticillati

~~folia oblonga~~ pedunculis villosis



perfoliata. C. foliis ovatis connatis approximatis acutis integerrimis ~~et~~  
Corymbis trichotomis

Crassula perfoliata foliis subrotundis, in medio a caule perfoliatis,  
approximatis rubro punctulatis Scop del fl. inf. 3. p. 12. tab. 6.

Habitat in cap. b. p. 5

Fruticosa basi nudata caespitosa in paucos ramos simplicia  
caespitosa folia carnosa subtus convexa supra plana  
patentiora punctulis rubris aspersa

Crassula perfoliata foliis connatis perfoliatis approximatis cordatis  
punctatis glaberrimis margine purpureo La Mart. Encycl.

repens Crasula foliis obovatis, pedunculis unifloris caule repente.

Habitat ad fretum magellanicum

Caulis filiformis ramosus. Rami oppositi ramati ad apicem foliosi. Folia ovata in petiolum angustate opposita distincta semina quicquidaria, glabra. Pedunculi capillares axillares usque ad folium, folio dimidio longiores

repens Brasilia foliis connatis ovatis integris glabris, corymbo supero decomperto Lin Suppl  
p. 109. N. V. pag. 307.  
Habitat in Cap. b. frei

muti yrota

Cultivata Crasula foliis oppositis obtuse ovatis integerrimis subultratis obliquis: paribus  
connatis remotis Len. S. V. pag. 305.

Crasula anacamperotis folio Dill. elth. 115: t. 97. fig. 114.

Habetat in Othopia ty.

Rami purpureo-florescentes <sup>teretes glaberrimi.</sup> Folia exigua lanata carnosa, perfoliata  
ultima lanceolata. ~~Corymbus~~ <sup>racemus</sup> terminalis trichotomis.

foliis obovatis obliquis integerrimis remotis, caule elongato, flori-  
bus paniculatis thuris

an me ~~lancei~~ <sup>non</sup> ~~non~~

2  
1  
Thyrsiflora Crasf. foliis perfoliatis ovatis ciliatis patulis, <sup>panicula</sup> ~~composito~~ <sup>subpicata</sup> composito attenua  
Dober. Suppl. 190 S.V. p. 307 Thunb. prod. v. pag. 53

Habitat in Cap. bono spei

imbriata Crasula foliis ovatis acutis quadrifariam imbricatis Corbus, floribus  
axillaribus sessilibus Hort. Kew. 1. p. 393.  
Habitat in cap. bono p. 1/2

non vidi

inanis P. foliis petiolatis ovatis, floribus quadrifidis ~~flore~~ <sup>lance</sup> simplicibus  
Thunb. prodr. pag 54.  
Habitat in C. b. per

non vidi

pubescens <sup>pubulati</sup> Crasf. foliis connatis ovatis acutis villosis, caule ramofo  
floribus <sup>compositis</sup> ~~crasfoliis~~ <sup>paniculato</sup> ~~paniculatis~~ Lin. Suppl. p. 190. t. V. 307 Thunb. prod.  
Habitat in Cap. bono spei



~~Crassula~~ <sup>1</sup> ~~caule~~ <sup>2</sup> ~~fruticosa~~, <sup>3</sup> foliis <sup>4</sup> ovatis basi attenuatis <sup>5</sup> connatis integerrimis <sup>6</sup> intra <sup>7</sup>  
marginem punctatis, <sup>8</sup> cymis <sup>9</sup> panicula <sup>10</sup> formibus Hort. Kew. t. p. 396.

Habitat in Cap. bona spei 7

~~Folia radi~~ Pedunculus longatus nudus, parvales oppositi patentissimi  
foliolum lanceolatum ad basin singulis, bifidi multiflori

foliis obovatis subulatis albo punctatis perfoliatis, panicula trifida  
Thurb. prod. pag. 56

non vidi

obovoides Crassula foliis ovatis distinctis acutis ciliatis, caule simplici pilosissimo  
lo, racemo composito: ramis paniculoformibus Hort. Kew. 1. p. 394 \*

Habitat in Cap. bono frei ♂.

Caulis vix spithameus. Folia carnosae, glabra, punctis minutis rubris impressis  
adpressa, unicalia. Racemus terminalis. Pedunculi distantes, patentes,  
subdivisi in Cymastriehotomas. Calycis foliola glabra, lineam longa. Pe-  
tala Calycis duplo longiora, alba, inferne viridifera. Nectaria lutea,  
filamenta alba. Anthera subrotunda, parva Hort. Kew. l. c.

montana Crasf. foliis connatis ovatis acutis ciliatis, caule fibrato, floribus aggregatis.  
Linn. Suppl. 109. N. V. p. 307

Habitat in Cap. bono spec.

non v. d.

punctata Craspula foliis oppositis ovatis punctatis ciliatis: inferioribus oblongis Lin. Spec:  
pl. pag. 406. & L. V. pag. 305

Craspula caule glabro, foliis connatis cordatis succulentis, floribus confertis  
Mell. Diet

Habitat in Othopia

Caulis teres, lavis, simplex. Folia opposita disticha, oblonga, sessilia, carnosa, exi-  
rata, punctata, subtus convexa, venis nervis ciliata. Floralia ovata. Corymbi axil-  
lares, brevissimi, fastigiati. Corolla campanulata, alba, limbo reflexo.  
Anthera purpurea Lin. Spec. l. l.

non vidi

umbellata

C. foliis ovatis subpetiolatis obtusis integris caule capillari erecto  
floribus umbellatis Thunb. prodr. pag. 54  
~~Habitat in C. b. per~~

non Ed.

obliqua Craspula foliis oppositis ovatis oblongis integerrimis acutis distinctis; mar-  
gine subcartilagineis Hort. Kew. 1. p. 393.

Cotyledon ovata Mill. Dict.

Habitat in Cap. bonae spei t.

alpestris Crasf. foliis connatis ovatis acutis quadrifariam imbricatis, capitulis pedun-  
culatis, Caule folioso Lin-Suppl. p. 109. L.V. p. 307

Habitat in Cap. b. pei

non vidi

pyramidata Crasf. foliis connatis ovatis obtusis quadrifariam incumbentibus; capitulis  
sessilibus Lin. suppl. pag. 109 \* S.V. p. 307.

Habitat Cap. b. p. ei

Planta pulchra quadrangularis, foliis ante approximatis; Lin

non vidi



testa Crasula foliis connatis ovatis obtusis cinereo lamellosis, scapo nudo, capitulo  
terminali Lin. Suppl. pag. 190. L.V. p. 307.

Habitat in Cap. bono spei

non vidi

argentea Crasula foliis connatis ovatis integris argenteis, caule tereto, corymbo supra  
decomposito Len. Suppl. p. 180. S. Sp. 306.

Habitat in Cap. b. p. ei

pellucida Crasula caule flavido repente, foliis ovatis sessilibus

Crasula, portularia facie repens Dill. Elth. 119. t. 100. fig. 119.

Crasula pellucida: Caule flavido repente foliis oppositi Len. l. v. pag.

306

Habitat in Cap b. pui

non ordi

muscosa Crasula caule herbaceo prostrato, foliis oppositis ovatis gibbis umbilicatis  
floribus sessilibus solitariis Lin. Bot. Acad. 6. p. 86 no 10\*

Habitat in C. b. p. c.

Caules filiformes, rarius <sup>terrestres</sup> ramosi, secti foliis minimis ovatis, oppositis,  
sessilibus, carnosis obtusiusculis; <sup>stipulati</sup> in exsertatis sub foliorum margi-  
ne. ordo punctorum. Flores ex alis solitarii, omnium minutissimi, lon-  
gitudine foliorum. Lin. l. c.

marginata

*C. folio ovato connatis margine membranaceo, pedun-  
culi unifloris, caule debili* Thunb. prod. per. 56

Habitat in Cap. b. prae

Caule debili glaber apice dichotome divisus. Folia remota  
unquicquidaria inferiora basi saepe angustata obtusa florata  
subcordata acuta, omnia margine tenuissimo membranaceo  
pedunculi in dichotomia pollicari filiformes calyces  
et petala caricolata

Dichotoma hort. kew. an huius pertinet

Coccinea Crassula foliis crassis planis cartilagineo ciliatis basi connatis vaginantibus Lin. Mart. 360.

S. J. p. 304

Cotyledon africana frutescens, floe umbellatis coccineo Bon. rar. 24. Tab. 24. Bradl.  
Juss. 5. p. 4. t. 50.

Cotyledon africana frutescens, floe carnes amplo Breyn. prod. 3. p. 30. t. 20. fig. 1.

Habitat in Cap. b. p. 5

Cauli ramosis. Folia carnea, confertim opposita hinc quasi quadryfaria  
glabra, margine albo unguiculari, suprema longiora lanceolata.  
Raspidium terminalis bis dichotomis, floribus sessilibus alternis, coccineis  
sempiternis, monopetalis limbo spatioso reflexo. Foliola calycis  
linearis lanceolata attenuata ciliata. cilia tenuissima Corolla sepius quilla  
cano laciniis oblongis

— lanceolato-oblongis