

# Cupania

Cal. 5phyllus. Cor. 5petala. Petala apice cucullata. Stylus trifidus. Caps. trilocularis. Sem.  
solitaria

1. Tomentosa C. foliis simplicibus obovatis dentato-lobatis
2. Pyracifolium C. foliis simplicibus ovato-subrotundis integerrimis
3. Tomentosa C. foliis pinnatis obovatis obtusis subtus ferrugineo-tomentosis
4. Saponarioides C. foliis pinnatis oblongis integris, subtus scabris
5. glabra C. foliis pinnatis ovatis obtusis crenatis glabris

Cup. laevis et dichrylla v. d. Ecl. 4

Dent. Cupania ~~filipponensis~~ folioli <sup>obovato</sup> oblongo, <sup>obovato</sup> abae imbricatis glabris teri-  
bus, capsulis tomentosis

Habitat in Guinea Thonning †

Arbor ramossima, magnitudine Quercus mexicana. Rami inferne teretes,  
superne angulati, tomentosi. Folia petiolata, pinnata cum pari, 2. 3 vel quadri-  
# sel obtusis pinnis absque acuminis simplicibus vel basivariis sub compositis  
pinnis: foliola petiolata, opposita, tripollicaria, interiora minora, oblonga, basi a-  
cuta, apice acumine brevi terminata, integerrima, utrinque glabra, costis  
subtus exceptis, supra avenia, subtus venosa. Petioli teretes, tomentosi,  
pennales brevissimi. Racemi versus summities ramorum, axillares, so-  
litarii, semipollicares, lani. # Pedunculus angulatus subtomentosus: pedicelli  
dispersi, solitarii interdum gemini, <sup>vel plures</sup> patentes, semipollicares uniflori  
flores polygami, masculi rudimento germinis, feminei antheris exsertis

Calyx quinquepartitus. Petala calycis duplo majora, lanceolata, basi  
versus acuta squama creta, concava. Nectarium receptaculum craspi-  
supra planum. Filamento corolla longiora, inferne villosa, nectario inserta.  
# Stylus contortus, obscure triplicatus. Capsula magnitudine pomi, carnosa  
pallida flava, hinc nubicunda, benecio tomentosa. Semina solitaria, magna

Germon obtusis  
gem, triq  
nium.

<sup>anellat</sup>  
cerasi, receptaculo papirae insidentia. Anellus cordiformis, externe convexus,  
interne planiusculus, vane incisus, interne foveola pro funiculis umbilicalibus  
carnosus, albidus, dulcis. Funiculi umbilicales gemini, coracii, utraque a dissepimento  
enati, venenati. Thonnung.

lanceolata Cupania foliis geminatis oblongis venosis, panicula terminali, capsulis  
glabris

Habitat in Guinea Gert 7

Rami teretes, glabri, juniores ferrugineo subtomentosi. Folia breviter petiolata,  
foliolo subsessilibus, tripolliciana, tenuissime venosa, supra nitida,  
integerrima. Pedunculi angulati, ferruginei.

confer cum D. li



*Ruparia tomentosa*

tomentosa Cupania foliis pennatis obovatis retusis inferne ferrugineo tomentosis  
Sw. prodr. pag. 61

Habitat in insula Trinitatis de Ryan.

Rami teretes, superne pulverulento-vellofi.

Folia petiolata, alterna, pennata impari bis. Trijuga: foliola petio-  
lata, intima alterna, reliqua opposita, cuneata, extima 4. 5. polli-  
cari intima triplo minor, remote dentato-ferrata supra coria sub-  
nitida, glabra, avenia, secundum nervos lineata: subtus venoso-  
reticulata, vellofa.

Petiolus communis tripollicaris: partiales brevissimi <sup>trinitati</sup>

Pedunculi axillares, subangulati, folio longiores. Flores compositi in  
racemo composito: rami patentes, alterni, sensim decrepentes, infi-  
mi ramosi. Pedicelli brevissimi 3-5. ex eodem puncto. Bractea  
sub ramificationibus ad pedicelli subulati, uti pedunculi, color pedi-  
celli et calycis. Et capsula vellofo-tomentosa.

Calyx pentaphyllus: folioli oblongi obtusi extus convexi in plicis, persistentibus  
Corolla petala 5: cuneis formis, vellofo, profertim marginis calycis vel longiora

1864

Cat. Rypan. n. 37.

Stamina filamenta 8, corolla parum longiora, basi vellofo. Anthera oblonga

germen superum, oblongum, tomentosum. Stylus brevis tomentosus. Stigmata tria, glabra, acuta.  
Capsula turbinato trigona magnitudine nucis, cori, involucelani, trivalvis salvis  
obcordati  
Semen unicum in fundo pericly coriuli, <sup>subrotundum</sup> ~~oblongum~~ <sup>nitidum</sup> leve, anillo carnos, excavato  
to infidens

<sup>vel monoua</sup>  
Polygame Divia videtur necesse est germen absque style in nonnullis inveniri  
Differentia generis anll. addenda.

*Puponia*

Habitat Cayana le Blond herb la Mantie

Racemi axillares et terminales, pedales, decemprophi patentes

Vidit la Mantie



Cupania

Habitat in insula Barbore Commesf. herb. Musf.

Tota glabra. Rami teretes. Folia petiolata, alterna, conjugata, vel bipuga, foliola opposita, subsessilia, bif. tri. pollicaria, <sup>interiorum minorum</sup> oblonga, obtusa, nitida, utraque glabra, venosa, subcoriacea, integerrima. Petioli linearis inter foliola paucum marginatis. Racemi axillares, compositi, foliis longiores

Tota abortu monoica ♂ Calyx 5que partitus: foliola 2 exteriora <sup>ovata</sup> minuta. Petala 5. ungue suballop. Stamina 8. filamenta basi villosa, corolla longior. Pistillum glandulosum. ♀ Cal. et Cor. ut in mare. Stamina brevissima, an sterilia? Germen subpedicellatum, trigonum, trilobum. Stigma simplex. Cupula trilobularis, loculis monospermis Musf. an Molucca  
a ludo specimen foliis conjugatis, saepe 5que jugis

• *macrophylla* Cupania

Habitat in insula Borbonia Commerçon herb. Mus. 7

• *Folia* bipinna: foliola subpetiolata, pithamœa, elliptica, utrinque venis pinnis, <sup>glabra</sup> al-  
teris margine bases breviora, obtusiuscula, integerrima. Petiolus glaber  
• Cinereo punctatus, uti pedunculi. Racemi subcompacti. Fructus oblongus,  
glaber

*Corygata Cupania* petiolis versutibus, foliis subtriplicis subsessilibus oblongo  
ovatis glabris venis nervatis floribus caducis Act. Bot. h. n. p. 109

P. 1. pag. 109

Habitat in Am. merid.  
ad decandram relictam

Sono - Rona quercifolia Aubl. R. h. p.

namidi

109  
fimbriolata Cupania folioli ovatis obtusi cum brevi acumine superne repan  
culis glabris alis nervorum fimbriulopetuis Act. Soc. N. P. 1. pag.

Habitat in Amer. merid.  
Secandra

<sup>Stygnis folium</sup>  
2 *Simpliifolia* *Cyrtaria*? foliis simplicibus ovalibus <sup>reticulatis</sup> subtus <sup>negatis</sup> pubescentibus

Habitat in Madagascar & Thonin

Rami <sup>flavido</sup> superiore tomentosi. Folia <sup>reticulata</sup> petiolata, alterna ovalia, <sup>marginibus ciliatis</sup> mucro

acumine brevi <sup>terminata</sup> <sup>subtus venis reticulatis constantibus negatis</sup> <sup>seabris ciliatis</sup> <sup>seabris ciliatis</sup> <sup>seabris ciliatis</sup> <sup>seabris ciliatis</sup>

<sup>stipulae parva subula</sup> <sup>caerulea</sup> <sup>caerulea</sup> <sup>caerulea</sup> <sup>caerulea</sup> <sup>caerulea</sup> <sup>caerulea</sup> <sup>caerulea</sup> <sup>caerulea</sup> <sup>caerulea</sup>  
Lateralibus <sup>caerulea</sup> <sup>caerulea</sup> <sup>caerulea</sup> <sup>caerulea</sup> <sup>caerulea</sup> <sup>caerulea</sup> <sup>caerulea</sup> <sup>caerulea</sup> <sup>caerulea</sup> <sup>caerulea</sup>  
~~petiolaribus~~. Fructus globosus, tomentosus, magnitudine <sup>caerulea</sup>

americana Cupania foliis simplicibus obovatis serrato dentatis

Cupania castanea folio, fructu sericeo & ramoso Plum. gen. 45.

<sup>46. 110.</sup> Cupania americana Lin. S. V. p. 864. Spec. pl. rar. 290\*

Habitat in America calidioris & herb. fr.

Rami cortice cinereo punctato Ramuli in apicis tergis

Folia opposita sessilia ovata integerrima venosa glabra

submarginata bipolliciana. Pedunculus terminalis solita  
rius glaber teris foliis brevior. Pedicelli umbellati 4-5. Flores  
glabri

Folia alterna, obovata, oblonga, obtusa, dentato serrata: nervis  
transversalibus crassis. Flores racemosi her.

rami teris, cucurbitatis, crispis fer penno corymbis, apice flavescen  
tia, tomentosa. Folia <sup>cu</sup> picholatis alterna, spithamea, internodiis minor  
remota quosque acutius serrata, oblonga, obtusa basi acuta  
venis pinnatis utrinque glabra. Petiolus teris, subtomentosus  
sur. Racemi compositi. Fructus tomentoso aures

creditur a Sw. eadem esse cum tomentosa

4 glabra Cupania foliis pinnatis ovatis obtusis crenatis glabris Schw. prod r. p. 61  
Cupania Brown. Jam. 170.  
Habitat in Jamaica

non vidi

3. *Laponarioides Lupania foliispennatis oblongis integris, subtus scabris Schum*  
prodr. p. 62.  
Habitat in Andra Occidentali

non v.



5 tomentosa Cupania foliis pennatis obovatis <sup>retusis</sup> inf. subtus ferrugineo-tomentosi Schw. prod.

pag. 61.

Trigonis tomentosa Jacq. Amer. 20102.1.

Habitat in Hispaniola,

arborum; rami teretes crassitate penna asperius superne uti petioli folia  
subtus, Pedunculi et calycis ferrugineo villosi tomentosi. <sup>caetero</sup> folia penna  
la pithamosa trijuga. <sup>altissima breviter petiolata</sup> foliola obovata basi acuta ~~ret~~ emarginata  
et retusis, supra glabra remote dentata basi integerrima exstima  
majori. Pedunculata Panicula terminalis racemosa pithamosa  
floribus confertis. Petala <sup>oblonga</sup> ~~oblonga~~ orbiculata villosa. Stamina longitudine  
corolla

de ser. ind. inter american