

Cyperus

umbellis effusis, pappulis alternis.

~~Cynopterus~~ C. culmo trigono aphyllis, umbella decomposita involucrio
Cineans brevior, pediculis lanceatis Thunb. ~~fl. rar.~~ fl. cap. pag. 18. Comby
Habitat in cap. b. ppe.

non vid.

55 46 55
Diffusus 56

Subgeminatis

Cyano ~~speculis digitalis subterni~~ lineari lanceolatis, Umbella diffusa,
~~foliis~~ foliis culmo altioribus.

Habitat in India orientali &

Culmi sesquipedales, tri^{quetri} ~~foliis~~. Folia culmo altiora, unguem fere longa, margine
scabra. Inducium subheptaphyllum, umbella longius, ~~penicillum~~ culmo lon-
gius; partialia tri-~~pentaphyllum~~ brevissimum, setaceum. Chroa oblique
truncata, obtusa. Umbella diffusa, ~~racem.~~ ~~subaequali~~ radiata; radii

~~longioribus~~ ~~epithemai~~, filiformes; partialia ~~etiam~~ radiis octo ~~secum~~ capillaribus,
~~potentissimis~~ ~~ultimis~~ ~~tribus~~ ~~quaque~~ ~~radice~~ ~~longioribus~~ ~~speculis~~

Supradecomposito

golitariarum, bina ~~terna~~ in singulis ~~radicibus~~ ultimis, laterales sessilibus
intermedia pedicellata, intermedia ~~sessilibus~~ ~~lineam~~ longa, angusta,

decemflos

ovata lanceolata, pennis convexa, acuta, fusco-virescentes: valvula
dors ovata, mucronata, dorso striata, dorso convexa, virides, margine
fusco, fructifera patula.

alternifolius 56
alternifolius 57

umbellae umbellatisque corymbosis

Cyperus ^{alternus} umbellae corymbosis, spiculis linearibus, ^{que} culmo ^{sub} striato, involucri involucri ^{sub} striato lobis.

56-57 57

Cyperus alternifolius, culmo triquetronudo apice alternatim folioso, pedunculis lateralibus prolixis Lin. Manit 20. Fauci collect.
2. p. 306. 4c. var. 2. t. 298.

Habitat in Madagascari.

~~Culmo teretibus, spiculis diffusi-
tis, subtilissime striato~~

Valde affinis, ~~spiculis linearibus, involucri foliolis angustis~~
sequenti differ. ^{si} ~~spiculis linearibus, involucri foliolis angustis~~ ^{pagina} ~~spiculis linearibus, involucri foliolis angustis~~
spiculis linearibus, rigenti-8x floribus. valvulis lanceolatis.

Culmus basi crassitie fere penno Cygnae, vaginis longe ^{inferne} vestitis

Ina Cyperus culmo triquetro seminudo, umbella foliosa decomposita, spiculis
alternis grandis distinctis Lin. Spec. pl. p. 67* S.V. pag. 97. Bot. gramp. 41 &
Gramen cyperoides Indiae orientalis alatus, panicula sparsa palea
pente Pluk. alm. 174. t. 101. fig. 7. rest. ned. ad fontonici de
Ina Balar Rheed. Mal. 12. p. 105. t. 56. figurum per firm. ad ibum

Habitat in India

Peduncululi in umbella plurimi apice umbelliferi, umbella ex tribus
quatuorve radiis, qui alternatim spiculas gerunt ex granis triquetris
obtusis, vix manifesto palea vestitis, parum distantibus Lin. Spe.

l. c.

In herb. Linn. est in n. fontonici Roth. et sic syn. Pluk. omnino in ten

2. *formosus* Cyprip.

~~Habitat in Louisiana Herb. Mus.~~

~~Petalus triquetus, angulis levibus; glaber, cuspide penna anserina. Umbella
composita: radii circiter 10. parvales pauciores. Involucrum 5-6 phyllum
longissimum, margine scabrum. parvales 2-3 phyllum, umbella brevior. Ochea
breves, truncatae, margine membranaceae. Spicula plures, sespales, in capitulo
ovato lanceolatae, unguiculari et ultra, compressae, multiflorae. Squamae
laeviores, lanceolatae, acutae spula, parum distantes~~

59 60 61
Elegans 60

Cyperus spiculis ~~digitatis~~ ^{linearibus} subternis, valvulis obcordatis mucronatis ~~dis~~ ^{line}

62 ~~spiculis~~, umbella ^{maxime} laxa
Cyperus panicula ^{maxime} sparsa ferruginea compressa elegantis sima. Sloan.
cat. 35 hist. 1. p. 117. t. 75. f. 1.

Cyperus elegans; culmo triquetro nudo, umbella foliosa, pedunculis nudis
prolixis, spicis confertis, mucronibus patulis Len. 845. v. 97.

Cyperus elegans; umbelli subradiatis & subtriplicatis, spiculis
linearibus, squamis obcordatis subdistantibus mucronatis patulis;
involucris planis Et. amer. 2. p. 7.

Cyperus laxus; culmo triquetro nudo, umbella foliosa laxissima,
spiculis subaggregatis raris & squarosis Lamark. illustr. 1. p. 146.

Habitat in Cayenna, Brasilia von Rohr

Culmus acute triquetrus; penna columbina parum tenuior. Folia
semirigida lata, margine scabius. Involucrum subheptaphyllum
foliolam unicam umbella longius, reliqua ^{um} aequalia vel breviora
rara extorsum, uti folia, margine scabiuscula; partiale di-Triphyl
cum umbellula brevius, setaceum. Petiole brevissimus. Umbella

composita, obcordata, patentissima. radii longiores quadrupoliarum, ^{sub}particulis
subtriradiata: radii pollicares vel breviores. Spicula subterna

linearis, tridecim - quatuordecim floris: ^{utrinque} valvulae prope carinam bis tripartae, margine
membranaceo. Semina triquetra
specimens mei Optime cum figuris Sloanii conveniunt, nisi quod spicula pauciores.

~~62~~ 61 Demidatus 61 Pyrenis spiculis Digitatis Subsenis linearibus, valvulis apice patulis, linearibus lanicollatis,
involucro submonophyllo brevissimo.
60 61 Pyrenis demidatus; calmo triquetro, involucro subnullo Lin. Suppl.
90.

Habitat in Cap. b. Spei. Osbeck. 7

Pulmus sesquipedalis, penna colymbina crassior, triqueter.
Involucrum submonophyllum, semipollicare: partibus nullis. Octrea brevissima.
Umbellae radii septem- octo, bipollicares; partialis subirradiata; radii
vix unguiculares. Spicula tres sex, unguicularis, viginti sex glora,
fusca: valvula enervis, non lanicollata, acuta, margine membranacea,
enerviae.

Composit.

biceps Cyperus

Habitat Golam h. b. Gusf.

Culmus rigidus aphyllus, erectus, pedalis, obtusa triquetris, laevis. Folia radicalia, carinata, glabra. Spicula in apice aggregata in capitulo majoritate nuc. a. Guylanae, major, cum pedunculo longitudine capituli, capitulo duplo minore. Involucrum hexa phyllum, umbella 3 plarea quadruplo longiore, margine scabrum. Spicula lineares, semipollucaris, obtusa fuscula, compressa, multi flore, nitida, flavescenti. Squamae oblongae, acuminatae.

~~94~~
vixiopsis C. culmo triquetro nudo, foliis convolutis vestibus; umbella foliis 2 admodum
posita, pinnis confertis, pinnulis compressis ovatis persistentibus. Sw. pr. p. 20.
Cyperus 11. Brown. Jam. 179.
Habitat in Jamaica. 4

sed duo foliola saepe tam brevia et facila, diphyllum si caute non attendatur,
pro diphylo ~~sumitur~~ ^{habentur}. Radii omnes interdum brevis & imae ^{et hinc} ~~et hinc~~ ^{et hinc} ~~et hinc~~
~~exone~~ ^{40 ne} glomerae in capitulo globoso magnitudine fere juglandis glomeratae
apparent. Cyperus geminatus Retzii obs. 4. p. 11. ~~et~~ absque dubio ^{est} diversus

~~Idem~~
an eadem cum vixiopsis herb. New.
descript. vide inter Amer. Puyanii

63 ~~64667~~ 6

ochraceo ~~64~~ Cyperus ^{oblongis} ~~terminatis~~ ^{laevibus} spiculis linearibus ~~globo~~ ^{globo} capitulo ^{patentibus}
sorbulis oratis, involucrio pentaphyllo umbella longiore.

Habitat in insula St. Anni West. ⁷⁴ ~~75~~ ⁷⁶ ~~77~~ ⁷⁸ ~~79~~ ⁸⁰ ~~81~~ ⁸² ~~83~~ ⁸⁴ ~~85~~ ⁸⁶ ~~87~~ ⁸⁸ ~~89~~ ⁹⁰ ~~91~~ ⁹² ~~93~~ ⁹⁴ ~~95~~ ⁹⁶ ~~97~~ ⁹⁸ ~~99~~ ¹⁰⁰ ~~101~~ ¹⁰² ~~103~~ ¹⁰⁴ ~~105~~ ¹⁰⁶ ~~107~~ ¹⁰⁸ ~~109~~ ¹¹⁰ ~~111~~ ¹¹² ~~113~~ ¹¹⁴ ~~115~~ ¹¹⁶ ~~117~~ ¹¹⁸ ~~119~~ ¹²⁰ ~~121~~ ¹²² ~~123~~ ¹²⁴ ~~125~~ ¹²⁶ ~~127~~ ¹²⁸ ~~129~~ ¹³⁰ ~~131~~ ¹³² ~~133~~ ¹³⁴ ~~135~~ ¹³⁶ ~~137~~ ¹³⁸ ~~139~~ ¹⁴⁰ ~~141~~ ¹⁴² ~~143~~ ¹⁴⁴ ~~145~~ ¹⁴⁶ ~~147~~ ¹⁴⁸ ~~149~~ ¹⁵⁰ ~~151~~ ¹⁵² ~~153~~ ¹⁵⁴ ~~155~~ ¹⁵⁶ ~~157~~ ¹⁵⁸ ~~159~~ ¹⁶⁰ ~~161~~ ¹⁶² ~~163~~ ¹⁶⁴ ~~165~~ ¹⁶⁶ ~~167~~ ¹⁶⁸ ~~169~~ ¹⁷⁰ ~~171~~ ¹⁷² ~~173~~ ¹⁷⁴ ~~175~~ ¹⁷⁶ ~~177~~ ¹⁷⁸ ~~179~~ ¹⁸⁰ ~~181~~ ¹⁸² ~~183~~ ¹⁸⁴ ~~185~~ ¹⁸⁶ ~~187~~ ¹⁸⁸ ~~189~~ ¹⁹⁰ ~~191~~ ¹⁹² ~~193~~ ¹⁹⁴ ~~195~~ ¹⁹⁶ ~~197~~ ¹⁹⁸ ~~199~~ ²⁰⁰ ~~201~~ ²⁰² ~~203~~ ²⁰⁴ ~~205~~ ²⁰⁶ ~~207~~ ²⁰⁸ ~~209~~ ²¹⁰ ~~211~~ ²¹² ~~213~~ ²¹⁴ ~~215~~ ²¹⁶ ~~217~~ ²¹⁸ ~~219~~ ²²⁰ ~~221~~ ²²² ~~223~~ ²²⁴ ~~225~~ ²²⁶ ~~227~~ ²²⁸ ~~229~~ ²³⁰ ~~231~~ ²³² ~~233~~ ²³⁴ ~~235~~ ²³⁶ ~~237~~ ²³⁸ ~~239~~ ²⁴⁰ ~~241~~ ²⁴² ~~243~~ ²⁴⁴ ~~245~~ ²⁴⁶ ~~247~~ ²⁴⁸ ~~249~~ ²⁵⁰ ~~251~~ ²⁵² ~~253~~ ²⁵⁴ ~~255~~ ²⁵⁶ ~~257~~ ²⁵⁸ ~~259~~ ²⁶⁰ ~~261~~ ²⁶² ~~263~~ ²⁶⁴ ~~265~~ ²⁶⁶ ~~267~~ ²⁶⁸ ~~269~~ ²⁷⁰ ~~271~~ ²⁷² ~~273~~ ²⁷⁴ ~~275~~ ²⁷⁶ ~~277~~ ²⁷⁸ ~~279~~ ²⁸⁰ ~~281~~ ²⁸² ~~283~~ ²⁸⁴ ~~285~~ ²⁸⁶ ~~287~~ ²⁸⁸ ~~289~~ ²⁹⁰ ~~291~~ ²⁹² ~~293~~ ²⁹⁴ ~~295~~ ²⁹⁶ ~~297~~ ²⁹⁸ ~~299~~ ³⁰⁰ ~~301~~ ³⁰² ~~303~~ ³⁰⁴ ~~305~~ ³⁰⁶ ~~307~~ ³⁰⁸ ~~309~~ ³¹⁰ ~~311~~ ³¹² ~~313~~ ³¹⁴ ~~315~~ ³¹⁶ ~~317~~ ³¹⁸ ~~319~~ ³²⁰ ~~321~~ ³²² ~~323~~ ³²⁴ ~~325~~ ³²⁶ ~~327~~ ³²⁸ ~~329~~ ³³⁰ ~~331~~ ³³² ~~333~~ ³³⁴ ~~335~~ ³³⁶ ~~337~~ ³³⁸ ~~339~~ ³⁴⁰ ~~341~~ ³⁴² ~~343~~ ³⁴⁴ ~~345~~ ³⁴⁶ ~~347~~ ³⁴⁸ ~~349~~ ³⁵⁰ ~~351~~ ³⁵² ~~353~~ ³⁵⁴ ~~355~~ ³⁵⁶ ~~357~~ ³⁵⁸ ~~359~~ ³⁶⁰ ~~361~~ ³⁶² ~~363~~ ³⁶⁴ ~~365~~ ³⁶⁶ ~~367~~ ³⁶⁸ ~~369~~ ³⁷⁰ ~~371~~ ³⁷² ~~373~~ ³⁷⁴ ~~375~~ ³⁷⁶ ~~377~~ ³⁷⁸ ~~379~~ ³⁸⁰ ~~381~~ ³⁸² ~~383~~ ³⁸⁴ ~~385~~ ³⁸⁶ ~~387~~ ³⁸⁸ ~~389~~ ³⁹⁰ ~~391~~ ³⁹² ~~393~~ ³⁹⁴ ~~395~~ ³⁹⁶ ~~397~~ ³⁹⁸ ~~399~~ ⁴⁰⁰ ~~401~~ ⁴⁰² ~~403~~ ⁴⁰⁴ ~~405~~ ⁴⁰⁶ ~~407~~ ⁴⁰⁸ ~~409~~ ⁴¹⁰ ~~411~~ ⁴¹² ~~413~~ ⁴¹⁴ ~~415~~ ⁴¹⁶ ~~417~~ ⁴¹⁸ ~~419~~ ⁴²⁰ ~~421~~ ⁴²² ~~423~~ ⁴²⁴ ~~425~~ ⁴²⁶ ~~427~~ ⁴²⁸ ~~429~~ ⁴³⁰ ~~431~~ ⁴³² ~~433~~ ⁴³⁴ ~~435~~ ⁴³⁶ ~~437~~ ⁴³⁸ ~~439~~ ⁴⁴⁰ ~~441~~ ⁴⁴² ~~443~~ ⁴⁴⁴ ~~445~~ ⁴⁴⁶ ~~447~~ ⁴⁴⁸ ~~449~~ ⁴⁵⁰ ~~451~~ ⁴⁵² ~~453~~ ⁴⁵⁴ ~~455~~ ⁴⁵⁶ ~~457~~ ⁴⁵⁸ ~~459~~ ⁴⁶⁰ ~~461~~ ⁴⁶² ~~463~~ ⁴⁶⁴ ~~465~~ ⁴⁶⁶ ~~467~~ ⁴⁶⁸ ~~469~~ ⁴⁷⁰ ~~471~~ ⁴⁷² ~~473~~ ⁴⁷⁴ ~~475~~ ⁴⁷⁶ ~~477~~ ⁴⁷⁸ ~~479~~ ⁴⁸⁰ ~~481~~ ⁴⁸² ~~483~~ ⁴⁸⁴ ~~485~~ ⁴⁸⁶ ~~487~~ ⁴⁸⁸ ~~489~~ ⁴⁹⁰ ~~491~~ ⁴⁹² ~~493~~ ⁴⁹⁴ ~~495~~ ⁴⁹⁶ ~~497~~ ⁴⁹⁸ ~~499~~ ⁵⁰⁰ ~~501~~ ⁵⁰² ~~503~~ ⁵⁰⁴ ~~505~~ ⁵⁰⁶ ~~507~~ ⁵⁰⁸ ~~509~~ ⁵¹⁰ ~~511~~ ⁵¹² ~~513~~ ⁵¹⁴ ~~515~~ ⁵¹⁶ ~~517~~ ⁵¹⁸ ~~519~~ ⁵²⁰ ~~521~~ ⁵²² ~~523~~ ⁵²⁴ ~~525~~ ⁵²⁶ ~~527~~ ⁵²⁸ ~~529~~ ⁵³⁰ ~~531~~ ⁵³² ~~533~~ ⁵³⁴ ~~535~~ ⁵³⁶ ~~537~~ ⁵³⁸ ~~539~~ ⁵⁴⁰ ~~541~~ ⁵⁴² ~~543~~ ⁵⁴⁴ ~~545~~ ⁵⁴⁶ ~~547~~ ⁵⁴⁸ ~~549~~ ⁵⁵⁰ ~~551~~ ⁵⁵² ~~553~~ ⁵⁵⁴ ~~555~~ ⁵⁵⁶ ~~557~~ ⁵⁵⁸ ~~559~~ ⁵⁶⁰ ~~561~~ ⁵⁶² ~~563~~ ⁵⁶⁴ ~~565~~ ⁵⁶⁶ ~~567~~ ⁵⁶⁸ ~~569~~ ⁵⁷⁰ ~~571~~ ⁵⁷² ~~573~~ ⁵⁷⁴ ~~575~~ ⁵⁷⁶ ~~577~~ ⁵⁷⁸ ~~579~~ ⁵⁸⁰ ~~581~~ ⁵⁸² ~~583~~ ⁵⁸⁴ ~~585~~ ⁵⁸⁶ ~~587~~ ⁵⁸⁸ ~~589~~ ⁵⁹⁰ ~~591~~ ⁵⁹² ~~593~~ ⁵⁹⁴ ~~595~~ ⁵⁹⁶ ~~597~~ ⁵⁹⁸ ~~599~~ ⁶⁰⁰ ~~601~~ ⁶⁰² ~~603~~ ⁶⁰⁴ ~~605~~ ⁶⁰⁶ ~~607~~ ⁶⁰⁸ ~~609~~ ⁶¹⁰ ~~611~~ ⁶¹² ~~613~~ ⁶¹⁴ ~~615~~ ⁶¹⁶ ~~617~~ ⁶¹⁸ ~~619~~ ⁶²⁰ ~~621~~ ⁶²² ~~623~~ ⁶²⁴ ~~625~~ ⁶²⁶ ~~627~~ ⁶²⁸ ~~629~~ ⁶³⁰ ~~631~~ ⁶³² ~~633~~ ⁶³⁴ ~~635~~ ⁶³⁶ ~~637~~ ⁶³⁸ ~~639~~ ⁶⁴⁰ ~~641~~ ⁶⁴² ~~643~~ ⁶⁴⁴ ~~645~~ ⁶⁴⁶ ~~647~~ ⁶⁴⁸ ~~649~~ ⁶⁵⁰ ~~651~~ ⁶⁵² ~~653~~ ⁶⁵⁴ ~~655~~ ⁶⁵⁶ ~~657~~ ⁶⁵⁸ ~~659~~ ⁶⁶⁰ ~~661~~ ⁶⁶² ~~663~~ ⁶⁶⁴ ~~665~~ ⁶⁶⁶ ~~667~~ ⁶⁶⁸ ~~669~~ ⁶⁷⁰ ~~671~~ ⁶⁷² ~~673~~ ⁶⁷⁴ ~~675~~ ⁶⁷⁶ ~~677~~ ⁶⁷⁸ ~~679~~ ⁶⁸⁰ ~~681~~ ⁶⁸² ~~683~~ ⁶⁸⁴ ~~685~~ ⁶⁸⁶ ~~687~~ ⁶⁸⁸ ~~689~~ ⁶⁹⁰ ~~691~~ ⁶⁹² ~~693~~ ⁶⁹⁴ ~~695~~ ⁶⁹⁶ ~~697~~ ⁶⁹⁸ ~~699~~ ⁷⁰⁰ ~~701~~ ⁷⁰² ~~703~~ ⁷⁰⁴ ~~705~~ ⁷⁰⁶ ~~707~~ ⁷⁰⁸ ~~709~~ ⁷¹⁰ ~~711~~ ⁷¹² ~~713~~ ⁷¹⁴ ~~715~~ ⁷¹⁶ ~~717~~ ⁷¹⁸ ~~719~~ ⁷²⁰ ~~721~~ ⁷²² ~~723~~ ⁷²⁴ ~~725~~ ⁷²⁶ ~~727~~ ⁷²⁸ ~~729~~ ⁷³⁰ ~~731~~ ⁷³² ~~733~~ ⁷³⁴ ~~735~~ ⁷³⁶ ~~737~~ ⁷³⁸ ~~739~~ ⁷⁴⁰ ~~741~~ ⁷⁴² ~~743~~ ⁷⁴⁴ ~~745~~ ⁷⁴⁶ ~~747~~ ⁷⁴⁸ ~~749~~ ⁷⁵⁰ ~~751~~ ⁷⁵² ~~753~~ ⁷⁵⁴ ~~755~~ ⁷⁵⁶ ~~757~~ ⁷⁵⁸ ~~759~~ ⁷⁶⁰ ~~761~~ ⁷⁶² ~~763~~ ⁷⁶⁴ ~~765~~ ⁷⁶⁶ ~~767~~ ⁷⁶⁸ ~~769~~ ⁷⁷⁰ ~~771~~ ⁷⁷² ~~773~~ ⁷⁷⁴ ~~775~~ ⁷⁷⁶ ~~777~~ ⁷⁷⁸ ~~779~~ ⁷⁸⁰ ~~781~~ ⁷⁸² ~~783~~ ⁷⁸⁴ ~~785~~ ⁷⁸⁶ ~~787~~ ⁷⁸⁸ ~~789~~ ⁷⁹⁰ ~~791~~ ⁷⁹² ~~793~~ ⁷⁹⁴ ~~795~~ ⁷⁹⁶ ~~797~~ ⁷⁹⁸ ~~799~~ ⁸⁰⁰ ~~801~~ ⁸⁰² ~~803~~ ⁸⁰⁴ ~~805~~ ⁸⁰⁶ ~~807~~ ⁸⁰⁸ ~~809~~ ⁸¹⁰ ~~811~~ ⁸¹² ~~813~~ ⁸¹⁴ ~~815~~ ⁸¹⁶ ~~817~~ ⁸¹⁸ ~~819~~ ⁸²⁰ ~~821~~ ⁸²² ~~823~~ ⁸²⁴ ~~825~~ ⁸²⁶ ~~827~~ ⁸²⁸ ~~829~~ ⁸³⁰ ~~831~~ ⁸³² ~~833~~ ⁸³⁴ ~~835~~ ⁸³⁶ ~~837~~ ⁸³⁸ ~~839~~ ⁸⁴⁰ ~~841~~ ⁸⁴² ~~843~~ ⁸⁴⁴ ~~845~~ ⁸⁴⁶ ~~847~~ ⁸⁴⁸ ~~849~~ ⁸⁵⁰ ~~851~~ ⁸⁵² ~~853~~ ⁸⁵⁴ ~~855~~ ⁸⁵⁶ ~~857~~ ⁸⁵⁸ ~~859~~ ⁸⁶⁰ ~~861~~ ⁸⁶² ~~863~~ ⁸⁶⁴ ~~865~~ ⁸⁶⁶ ~~867~~ ⁸⁶⁸ ~~869~~ ⁸⁷⁰ ~~871~~ ⁸⁷² ~~873~~ ⁸⁷⁴ ~~875~~ ⁸⁷⁶ ~~877~~ ⁸⁷⁸ ~~879~~ ⁸⁸⁰ ~~881~~ ⁸⁸² ~~883~~ ⁸⁸⁴ ~~885~~ ⁸⁸⁶ ~~887~~ ⁸⁸⁸ ~~889~~ ⁸⁹⁰ ~~891~~ ⁸⁹² ~~893~~ ⁸⁹⁴ ~~895~~ ⁸⁹⁶ ~~897~~ ⁸⁹⁸ ~~899~~ ⁹⁰⁰ ~~901~~ ⁹⁰² ~~903~~ ⁹⁰⁴ ~~905~~ ⁹⁰⁶ ~~907~~ ⁹⁰⁸ ~~909~~ ⁹¹⁰ ~~911~~ ⁹¹² ~~913~~ ⁹¹⁴ ~~915~~ ⁹¹⁶ ~~917~~ ⁹¹⁸ ~~919~~ ⁹²⁰ ~~921~~ ⁹²² ~~923~~ ⁹²⁴ ~~925~~ ⁹²⁶ ~~927~~ ⁹²⁸ ~~929~~ ⁹³⁰ ~~931~~ ⁹³² ~~933~~ ⁹³⁴ ~~935~~ ⁹³⁶ ~~937~~ ⁹³⁸ ~~939~~ ⁹⁴⁰ ~~941~~ ⁹⁴² ~~943~~ ⁹⁴⁴ ~~945~~ ⁹⁴⁶ ~~947~~ ⁹⁴⁸ ~~949~~ ⁹⁵⁰ ~~951~~ ⁹⁵² ~~953~~ ⁹⁵⁴ ~~955~~ ⁹⁵⁶ ~~957~~ ⁹⁵⁸ ~~959~~ ⁹⁶⁰ ~~961~~ ⁹⁶² ~~963~~ ⁹⁶⁴ ~~965~~ ⁹⁶⁶ ~~967~~ ⁹⁶⁸ ~~969~~ ⁹⁷⁰ ~~971~~ ⁹⁷² ~~973~~ ⁹⁷⁴ ~~975~~ ⁹⁷⁶ ~~977~~ ⁹⁷⁸ ~~979~~ ⁹⁸⁰ ~~981~~ ⁹⁸² ~~983~~ ⁹⁸⁴ ~~985~~ ⁹⁸⁶ ~~987~~ ⁹⁸⁸ ~~989~~ ⁹⁹⁰ ~~991~~ ⁹⁹² ~~993~~ ⁹⁹⁴ ~~995~~ ⁹⁹⁶ ~~997~~ ⁹⁹⁸ ~~999~~ ¹⁰⁰⁰ ~~1001~~ ¹⁰⁰² ~~1003~~ ¹⁰⁰⁴ ~~1005~~ ¹⁰⁰⁶ ~~1007~~ ¹⁰⁰⁸ ~~1009~~ ¹⁰¹⁰ ~~1011~~ ¹⁰¹² ~~1013~~ ¹⁰¹⁴ ~~1015~~ ¹⁰¹⁶ ~~1017~~ ¹⁰¹⁸ ~~1019~~ ¹⁰²⁰ ~~1021~~ ¹⁰²² ~~1023~~ ¹⁰²⁴ ~~1025~~ ¹⁰²⁶ ~~1027~~ ¹⁰²⁸ ~~1029~~ ¹⁰³⁰ ~~1031~~ ¹⁰³² ~~1033~~ ¹⁰³⁴ ~~1035~~ ¹⁰³⁶ ~~1037~~ ¹⁰³⁸ ~~1039~~ ¹⁰⁴⁰ ~~1041~~ ¹⁰⁴² ~~1043~~ ¹⁰⁴⁴ ~~1045~~ ¹⁰⁴⁶ ~~1047~~ ¹⁰⁴⁸ ~~1049~~ ¹⁰⁵⁰ ~~1051~~ ¹⁰⁵² ~~1053~~ ¹⁰⁵⁴ ~~1055~~ ¹⁰⁵⁶ ~~1057~~ ¹⁰⁵⁸ ~~1059~~ ¹⁰⁶⁰ ~~1061~~ ¹⁰⁶² ~~1063~~ ¹⁰⁶⁴ ~~1065~~ ¹⁰⁶⁶ ~~1067~~ ¹⁰⁶⁸ ~~1069~~ ¹⁰⁷⁰ ~~1071~~ ¹⁰⁷² ~~1073~~ ¹⁰⁷⁴ ~~1075~~ ¹⁰⁷⁶ ~~1077~~ ¹⁰⁷⁸ ~~1079~~ ¹⁰⁸⁰ ~~1081~~ ¹⁰⁸² ~~1083~~ ¹⁰⁸⁴ ~~1085~~ ¹⁰⁸⁶ ~~1087~~ ¹⁰⁸⁸ ~~1089~~ ¹⁰⁹⁰ ~~1091~~ ¹⁰⁹² ~~1093~~ ¹⁰⁹⁴ ~~1095~~ ¹⁰⁹⁶ ~~1097~~ ¹⁰⁹⁸ ~~1099~~ ¹¹⁰⁰ ~~1101~~ ¹¹⁰² ~~1103~~ ¹¹⁰⁴ ~~1105~~ ¹¹⁰⁶ ~~1107~~ ¹¹⁰⁸ ~~1109~~ ¹¹¹⁰ ~~1111~~ ¹¹¹² ~~1113~~ ¹¹¹⁴ ~~1115~~ ¹¹¹⁶ ~~1117~~ ¹¹¹⁸ ~~1119~~ ¹¹²⁰ ~~1121~~ ¹¹²² ~~1123~~ ¹¹²⁴ ~~1125~~ ¹¹²⁶ ~~1127~~ ¹¹²⁸ ~~1129~~ ¹¹³⁰ ~~1131~~ ¹¹³² ~~1133~~ ¹¹³⁴ ~~1135~~ ¹¹³⁶ ~~1137~~ ¹¹³⁸ ~~1139~~ ¹¹⁴⁰ ~~1141~~ ¹¹⁴² ~~1143~~ ¹¹⁴⁴ ~~1145~~ ¹¹⁴⁶ ~~1147~~ ¹¹⁴⁸ ~~1149~~ ¹¹⁵⁰ ~~1151~~ ¹¹⁵² ~~1153~~ ¹¹⁵⁴ ~~1155~~ ¹¹⁵⁶ ~~1157~~ ¹¹⁵⁸ ~~1159~~ ¹¹⁶⁰ ~~1161~~ ¹¹⁶² ~~1163~~ ¹¹⁶⁴ ~~1165~~ ¹¹⁶⁶ ~~1167~~ ¹¹⁶⁸ ~~1169~~ ¹¹⁷⁰ ~~1171~~ ¹¹⁷² ~~1173~~ ¹¹⁷⁴ ~~1175~~ ¹¹⁷⁶ ~~1177~~ ¹¹⁷⁸ ~~1179~~ ¹¹⁸⁰ ~~1181~~ ¹¹⁸² ~~1183~~ ¹¹⁸⁴ ~~1185~~ ¹¹⁸⁶ ~~1187~~ ¹¹⁸⁸ ~~1189~~ ¹¹⁹⁰ ~~1191~~ ¹¹⁹² ~~1193~~ ¹¹⁹⁴ ~~1195~~ ¹¹⁹⁶ ~~1197~~ ¹¹⁹⁸ ~~1199~~ ¹²⁰⁰ ~~1201~~ ¹²⁰² ~~1203~~ ¹²⁰⁴ ~~1205~~ ¹²⁰⁶ ~~1207~~ ¹²⁰⁸ ~~1209~~ ¹²¹⁰ ~~1211~~ ¹²¹² ~~1213~~ ¹²¹⁴ ~~1215~~ ¹²¹⁶ ~~1217~~ ¹²¹⁸ ~~1219~~ ¹²²⁰ ~~1221~~ ¹²²² ~~1223~~ ¹²²⁴ ~~1225~~ ¹²²⁶ ~~1227~~ ¹²²⁸ ~~1229~~ ¹²³⁰ ~~1231~~ ¹²³² ~~1233~~ ¹²³⁴ ~~1235~~ ¹²³⁶ ~~1237~~ ¹²³⁸ ~~1239~~ ¹²⁴⁰ ~~1241~~ ¹²⁴² ~~1243~~ ¹²⁴⁴ ~~1245~~ ¹²⁴⁶ ~~1247~~ ¹²⁴⁸ ~~1249~~ ¹²⁵⁰ ~~1251~~ ¹²⁵² ~~1253~~ ¹²⁵⁴ ~~1255~~ ¹²⁵⁶ ~~1257~~ ¹²⁵⁸ ~~1259~~ ¹²⁶⁰ ~~1261~~ ¹²⁶² ~~1263~~ ¹²⁶⁴ ~~1265~~ ¹²⁶⁶ ~~1267~~ ¹²⁶⁸ ~~1269~~ ¹²⁷⁰ ~~1271~~ ¹²⁷² ~~1273~~ ¹²⁷⁴ ~~1275~~ ¹²⁷⁶ ~~1277~~ ¹²⁷⁸ ~~1279~~ ¹²⁸⁰ ~~1281~~ ¹²⁸² ~~1283~~ ¹²⁸⁴ ~~1285~~ ¹²⁸⁶ ~~1287~~ ¹²⁸⁸ ~~1289~~ ¹²⁹⁰ ~~1291~~ ¹²⁹² ~~1293~~ ¹²⁹⁴ ~~1295~~ ¹²⁹⁶ ~~1297~~ ¹²⁹⁸ ~~1299~~ ¹³⁰⁰ ~~1301~~ ¹³⁰² ~~1303~~ ¹³⁰⁴ ~~1305~~ ¹³⁰⁶ ~~1307~~ ¹³⁰⁸ ~~1309~~ ¹³¹⁰ ~~1311~~ ¹³¹² ~~1313~~ ¹³¹⁴ ~~1315~~ ¹³¹⁶ ~~1317~~ ¹³¹⁸ ~~1319~~ ¹³²⁰ ~~1321~~ ¹³²² ~~1323~~ ¹³²⁴ ~~1325~~ ¹³²⁶ ~~1327~~ ¹³²⁸ ~~1329~~ ¹³³⁰ ~~1331~~ ¹³³² ~~1333~~ ¹³³⁴ ~~1335~~ ¹³³⁶ ~~1337~~ ¹³³⁸ ~~1339~~ ¹³⁴⁰ ~~1341~~ ¹³⁴² ~~1343~~ ¹³⁴⁴ ~~1345~~ ¹³⁴⁶ ~~1347~~ ¹³⁴⁸ ~~1349~~ ¹³⁵⁰ ~~1351~~ ¹³⁵² ~~1353~~ ¹³⁵⁴ ~~1355~~ ¹³⁵⁶ ~~1357~~ ¹³⁵⁸ ~~1359~~ ¹³⁶⁰ ~~1361~~ ¹³⁶² ~~1363~~ ¹³⁶⁴ ~~1365~~ ¹³⁶⁶ ~~1367~~ ¹³⁶⁸ ~~1369~~ ¹³⁷⁰ ~~1371~~ ¹³⁷² ~~1373~~ ¹³⁷⁴ ~~1375~~ ¹³⁷⁶ ~~1377~~ ¹³⁷⁸ ~~1379~~ ¹³⁸⁰ ~~1381~~ ¹³⁸² ~~1383~~ ¹³⁸⁴

64 65 66

Vegetus 67

68

69

Cyperus spiculis lanceolatis subgloboso capitatis praeterea compactis
Salsulis oratis unineris, in volucrio umbella longiore.
Cyperus

Cyperus compressus Jacq. bot. vind. 3. t. 12.

Cyperus eragrostis; Culmo triquetro nudo, umbella composita foliosa, spi-

Cyperus glome- culis glomeratis, squamis acutiusculis Lamarck illustr. 1. p. 146 excluda
ratur; Culmo tri-
quetro nudo, umbella
tetraphylla, pedun-
culis papillatis

Cyperus declinatus Moench

glomeratis Walt.
fl. car. 70

Cyperus monandrus Culmo inferna folioso obtuse triangulo, umbella
polystachya composita, spicis lanceolatis confertis sessilibus, flori-
bus monandris Koth. catal. 1. p. 3.

Habitat in America meridionali, in Carolina Bosc 4

Glauco-virescens Culmus sesqui-pedalis, acute trigonus, inferna foliosis. Solia culmo bre-
viora, linear lanceolata, margine serrulato & caltrae Vaginae canina
laevia: vaginae uti folia subtus in exsiccato inter strias punctatae. Anvolucri
foliola quinque, umbella longiora, praeter nonnulla breviora ad capitulum

partia le sub-phyll centrale: Ochreas oblique truncatae, pallidae, purpureo subpunctatae, ore mem-
vert-

65 ~~69~~ 677066

formosus ~~69~~ Cyperus spiculis ~~terminatis~~ ovato-lanceolatis, ^{Capitatis} involucrio sub hexaphyllo

~~69~~ ~~67~~ longissimo margine scabro et ~~capitatis~~ ^{capitatis} ~~capitatis~~

Habitat in Louisiana. ex herbario Gussiae

Palmis trigonus, angulis 2 oppositis pennae asperinae, Trigones, ^{acuta} angulis ^{basi foliosus} lae

Bolia lineari-rubra. Involucrum ^{penta} ~~quingua~~ hexaphyllum: ^{partiale} umbellula 6 di-tri

phyllum, umbellula brevior. Ochraceae breves, ^{truncata} ore membrana

natae. Umbellula ⁴ ~~universali~~ radii ^{3, 2, 2, 2, 2, 2 pollicares} circiter decem, ^{partialis} paulis

res breves. Spiculae circiter octo, ~~sessile~~ unguiculares, ^{sigati} ~~trigati~~ ^{trigati}

florae, flavescentes: valvulae lineari-lanceolatae, ^{acuta} ~~latere~~

laesae, ~~stigma~~ ^{stringua} unia, ^{ad} ~~caninam~~ ~~longa~~, decidua.

4 Composita:

5

umbellula ~~trian~~ umb. Comp. longissima marginata

*Divaniatus Cyperus calms Triquetro umbella composita conferta subtiri
phylla, fructibus lanceolatis complanatis divanatis* L. M. T. La-poy

145.

Habitat in India orientali

michi cognatus

66 70-69

Surinamensis 69 *Cyperus* spiculis acatis capitulis subglobosis glomeratis; spiculis ovatis, in

~~volucro~~ longis binis.

Cyperus Surinamensis; Culmo triquetro nudo, umbella daciomposita, spicis oblongo-ovatis conglomeratis, mucronibus incumbentibus Rehb.

Cyperus
Graminis tertio spe
cies Martini hist.

~~Graminis tertio spe~~ 2. gram. 35. t. 6. f. 5

1. fide Martini: Habitat in Surinamo Holandae, v. Roer, in Porto ricco West 7

Radix fibrosa. Culmus semi- sesquipedalis, crassitie pennae columbinae, rigidus, obtuse trigonus. folia radicalia, nonnulla altitudine culmi, linearia ovata convoluta-subulata extrorsum margine carinaque serrulato-scabra, uti culmi glauca v. indiglauescentia. involucrium pentaphyllum, omnia umbella longiora, duo fere longitudine culmi: partialia diphyllum, umbella breviora, patentis binum, foliola ovata basi dilatata, in acumine

48 structura foliorum: unguiculata desinentia, margine basi membranacea. Umbella quae tri-daciom, longior bipollicaris: partialia tri- sex radiata. Capitula

Composita;

glab. subglosa, semina Coniandri duplo majora, a tribus- sex minoribus conglomerata; spiculis minimis ^{decem circiter} seu multis ^{sub} ovatis, acutis, involu-
crata. Spicula minima, plerumque in singulis capitulis, p. & tramineis

dem florae, & tramineis: valvula ovata, acuta, striato-

Cum spicula in capitulis confertissima sint, scirpium ~~capitulis undique confertis & spiculis confertis in undique imbricatis in~~
minimo intuitu facile assequi. ~~capitulis primis undique confertis~~
sanat umbellis subcompositis, compositis et daciompositis.
cat. -

planifolius Cyreni Inquies foliis prolongis planifolius umbella
composita, fructu illi confectis lineamentis acuti saturate capso
nei Act. Soc. par. 1. nov. 106
Habitat in Am. merid.

michi ignoto

Luzula 68 # 67
polycephalus 74

Cyperus capitulis simplicibus glomeratisque ovatis, spiculis oblongis,
involucro longissimis

Cyperus umbella simplicissima, polyphylla; capitulis varmina-
libus, solitariis, oblongo-ovatis, spicis ovatis, concavis, densissi-
me conglomeratis Rottb. progr. gram. 23. t. 13. p. 2.

Cyperus polycephalus; culmo triquetris, umbella polyphylla,
capitulis ovatis pedunculatis, spiculis densissime conges-
tis Lamour. illust. v. 1. p. 147.

Cyperus conoides; triquetris foliosus; umbella longissi-
me involuata simplicis; capitulis compacte compositis
obtusis conoides Ait. bot. par. h. n. par. 1. p. 106 fide herbarii
Lamour. et Richardii.

Habitat in Surinamo, Cayenna &

Culmus bipedalis et altior, crassitie pennae columbinae
superne, acute angulatus, basi foliosis & solis culmo bre-
viora, pall. linearia, pallide viridia. Involucrum subhepta-

phyllum: foliola omnia umbella longiora, margine carinaque striata, duo interdum sesqui pedalia. Ochreae oblique truncatae, ferrugineae. Umbella simplex, subseptem radiata; radii longiores bipollicares vel ultra: Capitula primo duplo vel quadruplo majora, interdum e tribus vel quatuor minoribus agglomerata. Spiculae numerosis binis, oblongae, pallide virescentes vel lateribus ferrugineis, parvae, confectis binis, oblongae, pallide virescentes, interdum lateribus ferrugineis: valvulae oblongae, unistriatae.
Synonyma a Rothbollio huc allata minime huc pertinent.

~~50 66~~
~~fili culms 54~~
~~57 60~~
~~57 68~~
~~64 68~~

^{capitulis}
~~Spiculis globosis capitatis patentibus~~
~~Cyperus umbella subuniradiata radice capillari, spiculis globo-~~
~~sis, involucrii triphylli, culm. que seta~~
~~cei.~~

~~Cyperus trigonus, Culmo triquetro nudo, umbella simplici, spiculis~~
~~linearibus confertis, basi horizontalibus. Lin. spec. pl. 69.~~

~~Habitat in Virginia, Carolina.~~
~~Culmus pedalis, obsoletus trigonus, folia spithamea et ultra, culmo~~
~~breviora. Involucrum triphyllum, erecto-patens, foliola duo um-~~
~~bella longiora. Radius bipollicaris. Spiculae confertae in capitulis~~

~~# Cyrtia oblique~~
~~truncata, obtusa. globosa, pedo duplo major, unicum in apice radii, altera ad eju-~~
~~tas in, copioso, vix anguiculari, lineares angusta, lineares, tera vinctu-~~
~~la, simpli flore: valvulae parum distantes, laxa imbricata, lineari-oblon-~~
~~ga, obtusiusculae, striatae, carina virides, lateribus flavo-estentibus.~~

~~De Synonymo, non dico omnino certus sum, specimen enim e proprio~~
~~herbario donavit nomine nomine ab ipso inscriptum, voto caelo~~
~~discrepano a Sloanei et Rottbolli. P. trigono, quod quoque~~
~~abundat, potest conferenti descriptionem Linnaei cum iconibus~~
~~et descriptionibus virorum nominatorum.~~

~~et~~
~~hinc trigonum et nomen trinitate~~

Cuculodes cephalus C. Culmo setaceo torquato nudo, capitulo lepti, involucrio
late trophyllo patentissimo, floribus nudiore. Reti. off. s. p. 114
Habitat in nemorosis Schaendranione.

Michelianus
Cephalodes

69 71
hyalinus 72

--- --- --- --- --- umbello radius elongatis, spiculis alternis.

Cyperus ^{multifloris} spiculis oblongis ~~remotis~~ & culis sub & squarrosis, valvulis ovatis
~~attenuatis~~ ^{involucris} hexaphyllo longis sermo.

Habitat in India orientali Rottler. O

Affinis sequenti, Differt: spiculis distantibus, latioribus, paucioribus, & squarrosis. valvulis lateribus ~~membranaceis~~, minime striatis.

Culmus digitalis - semipedalis. ~~Involucris~~ Folia et involucri
Culmo saepe longiora. Ochreae truncatae. Umbella tri-quadrifida.
Spiculae sex-octo, sex-octoflora: valvulae complanatae, carina
superne curvata.

Affinis sequenti, Differt: foliis ~~et~~ involucris latioribus, copio-
rioribus: spiculis paucioribus, distantibus, latioribus, & squarrosis;
valvulis attenuatis, nec mucronatis, lateribus membranaceis, lateralibus, mini-
me striatis, obsolete remote trinerviis, hyalinis.

Compactus ~~14~~ ¹⁷ ~~Cypripis~~ ^{compactus} ~~spicis~~ ^{subulatis} ~~capitata~~ ^{sub} ~~spiralibus~~, involucrio pen-
taphyllo ^{compactus}
~~compactus~~ ¹⁷ ~~Cypripis~~ ^{umbellata} ~~decomposita~~, involucrio pentaphyllo, spiculis congestis
subspiralibus. Retz. obs. 5. p. 10. 7

Habitat in China

Stemus bipedalis, striatus in parte foliosus. Folia altitudine fere cubiti, linea-
ria, margine carinae laevia: vaginae purpurascens. Umbellae radii

tres-quinque

striati. Involucrii foliolaem unicum sesquipedali, brevissimo tripolli-
cani, omnino margine scabra. involucella pro radiorum numero, capitulo
breviora, ciliata. Capitula compacta & spiculis ^{sub} ~~spiralibus~~ ^{semipetentibus} ~~linearibus~~

umbella decomposita

sessilibus, linearibus subulatis, ferrugineis, sexfloris: Valvulae subulatae, striatae, margine
convergentes, purpurascens, carina viridi Retz. l. c.

Confertus 72
70

^{oblongis confertis subulosis capitatis, patentibus}
Capitulo ^{subulosis} ~~subulosis~~ ~~confertissimo~~ ~~oblongis~~, valvulis ~~oblongis striatis~~, foliis
lineariibus.
Cyperus confertus; culmo triquetro nudo, umbella simplici foliosa, spiculis
ovatis conglomeratis, glumis apice recurvis acuminatis Swartz prod. 20
Habitat in Caribaeis. West.

^{acute angulati}
Culmi palmares, altiores usque ad bipedales, ~~triquetri~~, basi foliosi. Folii
altitudine culmi vel paulo longiora. Involucrum triphyllum: foliolar
~~umbella longiora longis simis; tria umbella breviora, petalica ad glomem~~
in capitulo centrali. Un. Ochreae truncatae. Umbella ^{simpliciter} radii quatuor; et
spiculae circiter quatuordecim, breves, octo decemflorae: valvulae
striatae, flavo, dorso virides.
Stamena duo vel tria. Semina trigona Swartz. l. c.
Specimina interdum habui, absque culmo, radio uno vel altero radi
cali.

Triquetri.

~~Productus Swartz. Specimen intermedium 70
Habitat Galam West. Ind. 70~~

Dyrrhus

bulbosus

Cyperus ^{sub} spiculis geminatis spicatis, culmo foliisque setaceis, radice
bulbosa.

Habitat ad Senegal Guineam, in Guinea Thonning, in India orientali

Roettler Nonis et Pictler

Bulbus ^{magnitudine pisi} ovalis ^{valvulae} a hujus anni bivalvis; apice
in setis plurimis distinctus,

^{Letacei}
tenuis *Hyllinga* *Calmo* ~~*foliis*~~ ~~*filiformibus*~~ ~~*in*~~ ~~*specie*~~ ~~*umbellatis*~~
~~*lesplibus pedunculatisque depauperatis, spiculi patulis*~~
~~*Habitat in nemorosis Indis Congo*~~
~~*an modo varietas panicis differat quod omnibus partibus*~~
~~*multo tenerior. Spiculis interdum 3-4 interdum plures*~~
~~*longius pedunculatis*~~

Chrysanthemum

num. 66 ⁶⁶ ^{Spiculis} *Cyperus spiculis* ^{valvulis mucronatis} *digitatis lanceolatis*, in volucri *Diphylla* umbel-
la longiore t.

173 ⁷⁵ 44

Gramen cyperoides pumilum elegans e madao spatana
Pluken. alm. 179. t. 191. f. 8.

Cyperus pumilus; culmo triquetro nudo, umbella *Diphylla* com-
posita, spiculis alterno *digitatis lanceolatis*, glumis mucrona-
tis Lin. am. acad. 4. p. 302.

Habitat in Gordia

Similis *C. compresso*, sed brevior, distinctus: glumis minus em-
bricatis et apice mucronatis. Spicula vix alterna per a digi-
tatae Lin. l. c.

Quamvis mihi ^{ignotus} ~~non~~, a *C. cyperoides* ^{pumilo} tamen *Rotböllii*
tamen desit separari cum Descriptio Linnei non bene
et figura Plukenetii ^{minime} ~~non bene convenit~~ cum Rotböllii spe-
cimina conveniat.

inflatus C.

culmo triquetris nudo, umbella simplicii ^{dominati} spici ^{congestis} capituli linearibus,
Squamis subulatis reflectis recurvatis
gramen hirsutum indicum, glomeratis capituli Coryphi coenocles
gluma e squamulis plurimis perspicuis antabris, ansta uncin
donatis compactis Pluk. Phyt. 170 Tab. 190. fig 7.
Stipus capitatis culmo seseti. foliis, capitulis axillari
bus, pedunculatis involucri diphyllo linearibus Burm

se. ind. p. 21.
Stipus intricatis culmo triquetris nudo, umbella foliosa
simplicii: flosculorum squamis subulatis recurvatis lin
ment. al. p. 102

Cyperus anstatis capitulis sessilibus pedunculatis que; spici
lis linearibus minimis; flosculis subulatis reflectis
Prothb. gram. p. 23. Tab 6. fig.

Vonat interdum
capitulo unico
sessili absque
lateralibus

Habitat in India orientali
Pulmi digitales vel parvis vestra basi vaginatis et foliis
folia cominata altitudine culmi involucri subtetraphyllum
umbella longior pedunculi 3-4. in pollicari inaequales
cum capitulo sessili in medio. spicula congesta in capitulis oblon
go magnitudine pisi congesti linearibus parvis squamis striatis pedic
cumine reflexo attenuatis hinc capitula quasi pilosa

~~6~~ ⁶ ~~75~~ ⁷⁶ ~~74~~ ⁷⁴ *Cyperus* ^{concoloratus} ~~concoloratus~~ ^{glomeratus} ~~concoloratus~~ ^{sub squarrosus} ~~concoloratus~~ ^{mucronatus}
^{nitens} *Cyperus* ^{spiculis terminalibus} ~~concoloratus~~ ^{sub squarrosus}, ^{valvulis obtusis}
^{involucro triphylo} ~~concoloratus~~ ^{longitudine culmi foliis quo longitudine culmi}
^{# umbella} *Cyperus* ^{nitens}; ^{culmi triquetris basi basi foliolis}, ^{umbellae simplicibus},
^{longiore} ^{involucro triphylo}, ^{floribus monandris}. Retz. obs. 5. p. 13

Habitat in India orientali ^{King} ^{foliiformis} ^{fl.}
^{nitens} *Culmis* ^{bi-quadrifloris}, ^(angulis acutiusculis). Folia radicalia,
basium unum vel alterum culmo longius, linearia, arista. Involucrum
foliola patentia: ~~duo~~ unum unicum longitudine culmi, tertium
umbellae brevius. Umbella [#] bi-quadriflora: radii vix pollicares.
^{Decem} Spiculae circiter sex in apice radiorum, obtusiusculae quatuor
decim-viginti floris interdum cinereo varicenter, nitidae: valvulae
ovatae, obtusae cum mucrone petulo, hyalinae, interdum griseo vi
rescentes, ut saepius versus marginem ferrugineae, dorso con-
carinatae, utrinque ad carinam bi-striatae.
Varia altitudine culmi polle

Simples

Culmus interdum nullus spiculis inter folia adradicem 80, 80 libis
Accedit ad squarrosam. Saepius sub nomine Cyp. pumili
lin. ex India orientali; ^{mus 800} an eadem idem an diversus?
Differat Cq: squarroso spiculis latioribus, nitidis disco non flexus
80 minus squarrosus.

Seneculus
compactus

Cyperis culmo triquetro nudo, capitulo terminali subtriphylo
Squamis striatis obtusifoliis La Mart. T. Enc. pag 144.

Habitat in Madagascaria

Speculo ovato compresso

75-167
Squarrosus

Cyperus spiculis alternis ^{linearibus} ~~conferentibus~~, ^{valvulis aristatis recurvis} ~~foliis involucri~~ ^{que} ~~linearibus~~.

¶ Culmi digitales,
foliis sopre brevioribus, gracilibus filiformibus, umbella simpliciter

Gramen cyperoides maderaspatanum minimum ischaemi paniculis cum aristis Pluch. ameth. 114. t. 397. f. 2.

Umbella simpliciter trijugata. Spiculae angustae, linearibus semifoliolosa

Cyperus squarrosus; Culmo triquetro foliosis, umbella foliosa glomerata, spicis striatis squarrosis Lin. Con. acad. 4. p. 303. Kottb. gram. 25. t. 6. f. 3.

angustae; disco flexuosa; valvulae lineares, latenibus membranaceis minutis striatis.

Cyperus maderaspatanus; Culmo triquetro, umbella composita, spiculis linearibus, glumis lanceolatis uncinato-mucronatis, involucrio triphylo longioribus lanceolato Wild. Spec. pl. 1. p. 270.

Habitat in India orientali - Kerig, Rottler 0

~~Similis~~ ~~affinis~~ sequenti, differens differens: foliis et foliolis ~~invalens~~ linearibus canalicatis, nec involuto-petalis; spiculis brevioribus, ad lentem flexuosis a flosculo ad flosculum ~~quod non in C. costaricensi observo~~. valvula lateris flava ~~et venata~~.

¶ non in apice sessilis sed alterna conferentibus linearibus

Variat eodem modo ac reliquae hujus sectionis. Jam antea monui quod C. pygmaeus Rottbollii omnino ~~differt~~ a C. squarroso Linnaei ^{usq. diversum} ~~per se~~ ^{cum} ~~non~~ ^{verte}

convenit

~~ad figuram Pleukenetii nec ad descriptionem squarrosi in amenitatibus Linnaei~~
 Specimen autem ^{herbarii Rothb. illi} ~~Cyperus~~ ^{quarrosus} dictus non modo cum ^{quadrat} ~~ad figuram citatam~~ ^{et descriptione} ~~Linnaei~~, ^{quadrat} ~~sed etiam~~ ^{speciminibus} ~~tae cum Widenowii~~ ^{descriptione} ~~C. maderassatensis Widenowii~~ ^{ae} ~~que ac specimenibus a Koenigio, Klavinis et Koetlero a Tranquebaria~~
~~misit, unde varosi Nulle Cal. Widenowiae quoque obtinuit. Quod Linnaeus~~
~~speculae Cus Linnaeus absque dubio habuit specimenque in radiis et tantum~~
~~modo speculibus terminalibus glomeratis, ^{infructum} ~~ideoque ac~~ ^{et} ~~cum non umbellam~~
~~tum deesse non habere dixit.~~~~

~~ad figuram Pleukenetii nec ad descriptionem C. squarrosi in amenitatibus Lin. conveniat. Cyperus in herbario Rothb. illi squarrosus dictus non modo ad figuram citatam Pleukenetii et descriptionem Linnaei omnino quadrat, sed etiam ad C. maderassatensem Widenowii et specimen a Koenigio, Klavinis et Koetlero misit, unde verisimile Cyperum suum esse Widenowius obtinuit. sine dubio specimen radiis destitutum et speculis terminalibus glomeratis infructum non habuit, ideoque cum umbella carere dixit.~~

Herbar. Koen. 1. pag. 146.
 1755. 4. 1.

Ceterum Cyperus velis frequenter modo umbella foliosae speculibus
 subunguibus non squarrosus a Mark. 1. pag. 146.

filices ~~76~~ 77

~~Cyperus spicatus~~ ^{ramotiusculus} spiculis linearibus lanceolatis distantibus, laxa imbricatis, involucri triphylo umbella longiora foliisque laxis
Habitat in Virginia

Culmi semi-pedales, filiformes. ^{acutangula} Sola radicalia, longitudine culmi, linearia, laxa, ~~triac~~ ^{triac} margine scabra. Involucri foliola cum foliis

patiale nullum. Similia: ⁴ Cyrea truncata. Umbellae ^{superplex} tri-quadriradiata: radius longior pollicaris: partialis tri-radiata: radii brevissimi. Spicula quatuor-

Subcomposita

octo, distanter, vix unguicularer, ^{duodecimflora} ^{patentespina} ~~complanata~~, flavo-ferruginea: ~~valvula~~ ^{oblonga} ~~laxa imbricata~~, flavo-ferruginea, ~~valvula~~ ^{oblonga} ~~laevigata~~: valvula ^{oblonga} ~~lanceolata~~, acuta, prope carinam utrinque unistriata. -

Composita

D. Hans Culmo Inquieto nudo, umbella foliosa supra decomposita, spicis alternis
filiformibus, flosculis distantibus. An suppl. p. 103. L. 8. pag. 90.

Cyperus culmo Inquieto nudo; umbella foliosa supra decomposita;

partim spicis digitalibus intricatis, spicis alternis congestis filiformibus
descriptis. ~~seca~~ ~~et~~ ~~subulatis~~ ~~Robt. Gram. pag. 37. Tab. 10. & ex clava Linn. L.~~
~~nam elatum pertinet~~ ~~figura elati Robt. ad elatum An pertinet~~
Habitat in India

Radix tuberosa Culmus Inquietus saepe bipedalis striatus basi foliosus.

Stola graminea margine leviter striata. An involucrium
particula triphylla 5-6 phyllum foliolis 3 bus umbella ^{maxima} Congiombus ⁴ non
umbellae radii 5-6 praeter 3 in centro brevissimo reliquis

umbella composita subspithamea filiformes partiales 4-5. patentissima
et decomposita ^{ultima ternis} centrali laterales centrali Congiombus. Spicula alterna
memoto ^{patentissima} angustissima subulata. ^{dentata e squarum distan}
4-5 annis ^{4-5 ult. ra. oblonga} tibus, unguiculatae. Squamae lineares dorso striatae
ni fallor a Robt. ^{4-5 ult. ra. oblonga} vnde ei margine leviter subferruginea obtusa
michi misit ad

Descriptio *Cyperus* Robt. pag. 37. partim sciendum specimen
elati partim sciendum specimen distans composita
figura vero ad *Cyp. elatum* umbella decomposita que
pertinet

77 78
intactus 79

Cyperus spiculis ~~altis~~ ^{remotis} linearibus ~~intactis~~, involucrio ~~brachyphylo~~
~~et~~ amphyllo foliis que rigidis.

Habitat Galam. Est herbario Russiaci 7

Similis in omni ~~omnino~~ ^{omnino} ~~precedenti~~, ~~discessit~~. ~~Spiculis~~ ~~magis~~ ~~distan-~~
~~tibus~~, ~~triplo~~ ~~brevioribus~~, ~~valentis~~ ~~marginem~~ ~~versus~~ ~~laevos~~, ~~absque~~ ~~ma-~~
~~culo~~ ~~spiculae~~ ~~breviores~~ ~~et~~ ~~arctius~~ ~~imbricatae~~, ~~foliis~~ ~~rigida~~.

~~nomen spiculae~~

nomen triale mut.

~~notus Cyrenis~~

~~Habitat herb. ha. Mantua~~

~~Spica brevis, latior~~

78

~~77~~ 78

olivaceus.

Spicis corymbosis laxis ~~rectis, # corymbosis~~
 77 ~~Cyperus~~ ~~Spiculis linearibus lanceolatis compressis, radii umbellulae corymbosis~~
~~in involucrio~~ ~~umbellulae corymbosis~~ ~~radii umbellulae corymbosis fasciculatis~~

involucro Habitat in Cap. b. 3 p. 2.

umbellam equante.

Culmus sesquipedalis, ^{gracili} aculeo trigonus, nudus. Involucrum ~~triphyllum~~
~~triphyllum~~

partiala nullum: foliola duo longitudine umbellae, ^{tercium} ~~tertia breviora~~ # Ochrea brevissima.

Umbellae ^{composita} radii septem, ^{quinque} quatuor bipollicares, subquales, fl. 4 for ones: par-
 tiales, duo vel tres, brevissimi, bi-^{triflori}. Spicula unguiculatae, parum
 distantes, patentissima, sordide flavo-virantes, ~~et~~ viginti duo floris, acuta.
 valvula lanceolata, mucronatae, deciduae, ut in sequente.

Differt ab in sequente: involucro brevi; spiculis radiis umbellae
 congruentibus, pluribus; spiculae distantibus. —

~~Strophanthus C. Culmo basi tuberoso fctis fctis
Stoloniferis Culmo ^{libro} umbella subternata; speculis lanceolatis concavis, involucri
diphullo, radice repandibru~~

~~Habitat in India orientali~~

~~Radix tuberosa oblonga fctis negris longis veta stolonifera stolones
articulati, per intervalla culmos triquetros pedales striatos
basi vaginatos et folios emittentes. Folia angusta carina-
ta, culmo dimidio breviora. Umbella pedunculis geminis
vel ternis, pollicantibus vel brevioribus cum vel absque umbella
centra. Spica in apice conferta 5 quinque sex unguisula
res obtusa. Squama ovata obtusa carina et margine pallide
virides. involucrium diphyllum intermedium 3te umbella lon-
gus profertur unicum intermedium 3trum breve.~~

~~Cyperus stoloniferus culmo nudo triquetro, umbella composita
involucrata, speculi lanceolatis Rete obs. 4. p. 10 x~~

parryi C. Culmo triquetro basi folioso, involucrio triphyllis, umbella
uni-versali de Composita: partialibus subtrifidis Retz. obs. 4 p. 10
Cyperus parryi Culmo triquetro, involucrio triphyllis umbella uni-
versali de Composita; partialibus subtrifidis, spicis alternis feli-
formibus Rothb. gram. pag. 31. Tab. 7. fig. 3.

Habitat in India orientali occidentali

simili effidente at folia angustiora spicis gracilioribus et panem
brevis

variat umbellis simplicibus compositisque, involucrio umbella aequata
vel longiore, spicis magis purpureis vel minus, panem latioribus vel angustioribus

spicis subrotundis lineatis

composita squamis obtusis periculis

po lycophala Cyperus

Habitat herb. L. manthii

Culmus triquetus, obtusangulus, striatus, glaber, pedalis, basi foliosus. Soli
lineana, loricata, acuta, culmi breviora. Capitula plura, ovata, compacta
nonnulli breves pedunculati, magnitudine vix Coryli. Involucrum,
phyllum, umbella longior,

Similem ex Puerto Rico de Dry radiis longioribus, spiculis ovatis, com-
pressis, squamis linearibus patulis, capitulis majoribus. a loricato forte
non diversus eadem forte ac illa capituli minoribus proleseri

82-83
flavidus St.
St.

Cyperus spiculis alternis approximatis patentibus, in involucrio triplicato subulato.
patentibus, in involucrio embriicati
discrepantibus linear lanceolatis,

Cyperus flavidus; culmo triquetro striato, umbella composita triplicata, spiculis lanceolatis, aequa multis monandris Retz. obs. 5. p. 13.

Cyperus divaricatus; culmo triquetro, umbella composita, conferta subtriplicata, spiculis linearibus complanatis divaricatis Comar. illustr. 1. p. 145.

Habitat in India orientali Hong, Specimina culta in herbario La Martii. vidi

Culmi semipedales et ultra, obsolete trigoni. Folis radicalibus plerumque, unicum infera in culmo eoque dimidio brevius, angusta. Involucrum folioleto erecto-patentia, duo umbella longiora, unicum aequante.

Ochreae breves
obtusae.

Umbellae ^{subsimplae;} radii plerumque quatuor, longioribus tripollicaribus. Spiculae circiter octodecim, viginti sex - triginta florum, in cultis saepe quinquagintaflorae, acutiusculae, sordide flavescentes; ^{nitidae} involuculae linearibus oblongae, obtusae, margine tenui membranaceo, dorso striato, lateribus laevibus. Rachis inter flores flexuosa.

Semen pyriforme, triquetrum, nigrum.

Rachis in haec species et alii in capitulum conglomeratum ~~confertum~~ ~~confertum~~. Stamen unicum, uti Retz. vidi. Ab antecedente differt involucris non patentibus apicibus non mucronatis

~~Cyperus~~

~~Habitat in Madagascari~~

~~Culmi teretri, aphylli, acutanguli, saepe pedales, teres. Volea attenuata, carinata, culmo dimidio breviora. Anvolucrum triphyllum: foliola unicum bipollicare, tertium longitudine capituli. Spica plures, septem, pallida, multiflora, ovata, compressa; squama lanceolata, striata~~

~~vel spiculae duas infirmas geminatas vel ternas in infimo pedicello
 erat Praedita varis quod spiculae centrales non longiora sunt, sed
 aequales
 namque in galea imbedentes~~

Nec figura Ruicentii ne Descriptio Linnai ad hunc satis quadrant, ergo
 haec synonyma omisi. Omnia, quae vidi, specimina umbellam simpli-
 cem spiculis in infimo pedicello geminatis ternisve habuerunt.

Romae P. C. ultimo in quibus mado, sequitur
 tetrastichis de quibus sub quibus
 Habitat in frons montis de quibus
 Culmi tetra- et trigoni vel pentagoni
 grammas ad hunc modum. Capitulum
 eo numerosissimo spiculis oblongis
 in vertice tetra- et trigoni
 Corporulae vel in facie foliorum
 munitur

84 85 86

Flavescens 84

Cyperus spiculis linear-lanceolatis, umbella
triphyllo foliisque ~~setosis~~ umbello triplicata longiore.
Funicus exiguus proterensis Trag.

in involucrio

Gramen Cyperoides, minus, panicula sparsa subflava Bauh. Theatr.
88.

Cyperus minor pulcher, panicula lata compressa subflavescente
Monn. hist. 3. p. 239. 8. p. t. 11. f. 37.

Cyperus flavescens; Culmo triquetro nudo, umbella triphyllo,
pedunculis simplicibus inaequalibus, spicis confertis lanceolatis
Lin. Syst. veg. Pollich. palat. n. 41.

Habitat in Europae paludosis B.

Similis C. fusco. ~~Gramen~~ Culmus filiformis. Folia altitudinis culmi. Involucrum
Culmo brevius. Spiculae flavae: valvulae concavae, obtusae.
Spicula Termini interdum duo setae, involucrio triphyllo.

viridifolius. Culmo triquetro, umbella composita triphylla, spicis linearibus
confectis, glumis acutis patulis Wed-spec. p. 200

Cyperus compressus noster seu *viridis* Kunth & Schreb. Sicil. no. 69. tab. 83

Cyperus viridifolius

Hoffm. Germ.

Habitat in montosis Siciliae B

non vidi.