

Cyperus

147. ~~148~~

capitulis

aristatus ~~148~~ Cyperus Capitatus, valvulis calycinis subulatis recurvis.

Scirpus capitatus; Culmo tereti folioso, capitulis axillaribus pedunculatis, involucrio Diphylo linearum Burm. ind. 21.

# Scirpus Lappaceus; Scirpus intricatus, culmo triquetronudo, umbella foliosa simplicis, speciosus  
culmo triquetro "oblongis squamis calycinis recurvis Lin. mant. 182.  
pubescentibus, capitulis terminalibus solitariis  
terminalibus solitariis Cyperus aristatus, capitulis oblongis sessilibus et pedunculatis, spiculis  
nerviis involucrio, nervibus minimis, flosculis subulatis reflexis Roettl. gram. 23. t. 6. f. 1.

culmo triquetro  
pubescentibus, capitulis  
terminalibus solitariis  
nerviis involucrio,  
glumis striatis  
recurvis Lamarck.  
Illustr. 1. p. 139

# Habitat in India orientali et Cap. b. spec. 0

Radix capillaris. Culmi plures, fasciculati, <sup>Digitales,</sup> palmares vel parum  
ultra, acute triquetri. Folia linearia longitudine culmi, linearia,  
coninata, margine laevia. Involucrium <sup>foliosa</sup> triphyllosum; duo umbella  
duplex vel triplex longiora, quantum brevior. Ochreae truncatae, fer-  
rugineae. Umbella ~~trifida~~ bi-triradiata, radius longior interdum polli-  
caris. Capitula vix unguiculata, obtusissima, spiculae ferrugi-  
nea. Spiculae copiosae, lineam fere longa, alternae, confertissimae,  
patentes; octo- decem florae: valvulae in aristas uncinatas adde-  
muato, striatae.

# Simplex

Varia culmo pollicari vel desuper pollicari; capitulo globoso vel cylindrico,  
atque solitario absque umbella radiis lateralibus.

147

anstatu. Cyneru anstatu.

148 ~~148~~

glomeratus 154 *Cyperus* spicis glomeratis oblongis, <sup>imbricatis,</sup> spiculis <sup>valvulisque</sup> linearibus, involucri <sup>triphyllo</sup> umbella involucri <sup>cellis</sup> que triphylla umbellulis longioribus.

# *Cyperus* *cinna* *Cyperus* aquaticus italicus proterior, locustis venustissimis in ra-  
momaw; culmo eum dense congestis Mich. gen. 45. Segu. ver. 3. p. 68. t. 2. p. 2.

triquetro folioso *Cyperus* capitulis glomeratis congestis, seminibus aristatis Mont.  
umbella spiculis  
capitulis pedunc Gram. 14. t. 1. p. 1.

lati sessilibusque, *Cyperus* culmo triquetro nudo, umbella triphylla supradecomposita,  
involucro penta  
phyllis serrulato. spiculis glomerato rotundatis, spiculis subulatis Lin am. acad.  
scabro Act 2 obs. 4. p. 301.

4 p. 10

# Habitat in Italia paludosis, an ex India advena? Specimens  
~~debes Allionio et Brunellis super. Bononiensis Bononiensis~~

*Pulmus* bipedalis *Aultra*, eras sitia pensio *Cygnus*, acute <sup>angulatus</sup> *tripater*  
inferne foliosus. Folia duo <sup>10</sup> ~~tra~~, culmo altiora, longe vaginantia  
ex utroque margine canaque scabra. Involucrum foliola tria vel  
quatuor, proter interiora, umbella longiora: partialis duo vel

# *Subdecomposita* tria, umbellula longiora. Ochrea oblique truncata, ligula lan-  
142 ~~subdecomposita~~ *leolata*, integra. Umbellula <sup>non pariter</sup> ~~pradu~~ interdum novem, longiores spicula

*moi*. Spica omnes interdum sessiles, interdum tres vel quatuor pedicellatae, basi  
 glomeratae: Squamula numerosissima, patentes, lineares <sup>in semicirculares, compressae</sup> duae vel tres longae, quatuor-  
 dae in <sup>capitulo,</sup> florae, ferrugineae: valvulae lineares, angustae, obtusiuscula, vix <sup>interdum</sup> striatae.  
 A *P. alopecuroides* *Höftböllii* distinctissimus. Specimina culta vidi (Capitulo  
 solitario sessili, interdum umbella simplici, radio ~~monostachyis~~ unicapitatis.

glomeratae  
 Elytra  
 Elytra capitulis glomeratis conatis, serratis ensatis Mand. Gron. 14.  
 7.14.1.  
 Elytra capitulis glomeratis, conatis, serratis, ensatis, venis primis

150  
compactus ~~176~~ <sup>glomeratis, imbricatis, sessilibus, subulatis,</sup> Cyrenus spicis oblongis ~~sessilibus, subulatis,~~ spiculis ~~cylindricis~~ <sup>erectis,</sup> ligula  
ochrea bifida: laciniis divaricatis.

Cyrenus alexandrinus, Culmo trigono, umbella decomposita involucrio poly-  
phylo serrato breviora, spiculis cylindricis Thunb. prod. 18.

Habitat in Cap. b. sp. ~~Indiae orientalis~~ <sup>part. 4</sup>

umbella

Viridi-glaucus. Culmus altus, crassitie fere pennis anserinis, acuta an-  
gularis. Involucrum penta-hexaphyllum: foliola omnia ~~longis~~ <sup>longis</sup>  
sa, longius sesquipedale, margine carinaque serrulato scabra; partiale  
monophyllum ad singulum capitulum partiale, monophyllum, capitulo  
longius, linearis subulatum. Ochrea pollicares, trigona, obliqua, trunca

ligula serrata  
scabra

Umbella <sup>composita</sup> ~~part. quinque~~ - sex, longiorer quinque ~~sex~~ pollicares.

Spica quinque - octo, ~~sessiles~~, ~~intermedias~~ erectae pollicaris vel sesqui polli-  
cari, crassitie saepe digiti, ~~intermedias~~ ~~erectae~~, <sup>lateralis</sup> radiatum pater-  
ter, breviores, tenues, omnes obtuse ~~trigono~~ <sup>undique</sup> spiculis capis

vir. semi ungui-  
cularis

tris semis, ~~et~~ testa. Spiculae parvae, ~~trigono~~ <sup>trigono</sup>, ~~trigono~~ <sup>trigono</sup>, acuta: valvulae  
lineares, attenuatae, ~~strictae~~ <sup>strictae</sup> ~~concolorae~~, ~~dorso~~ <sup>dorso</sup> ~~et~~ ~~nato~~ <sup>virides</sup>, margine purpure  
rico

Hinc in herbario Lamarckii vidi sub nomine C. regularis Thunbergii & alio  
reverso com hinc retuli propter figuram spicularum.

~~176~~

compactus

~~Cypripus~~

Habitat in Cap. b. spec. 4

Habitus: <sup>o</sup> culmi nodos, basi et apice foliosi, in paucis praesertim pri-  
mo divisionis apylli. inflorescentia simplicissima - siccatione compo-  
sita.

<sup>o</sup> Radix fibrosa, in paucis tuberibus zonatis.



rostratus Schoereri

Habitat in America septentr. herb. Ruch.

Caulmi Iniquistri, glabri. Folia levia, plana, unguem lata, margine scabro, pinnis  
axillares, solitarii, serrati. Spiculis 6 flore, alterni, semipollicares, distan-  
tes, patenti sine. Squama ovata, ferruginea, exteriore minores. Semina glabra  
oblonga, levia, rostro longo subulato terminata

adest quoque in herb. Rusfi

\* Cúlcma terehi

xx = Cúlcma trigétra - spicis solitariis aggregatiue.

-- umbella contracta

--- umbella radiis elongatis, spiculis ~~umbellatis~~ terminalibus, umbellatis

---- umb. rad. elong. ---- spiculis alternis ----