

# Cypripedium

1. calceolus C. foliis caulinis ovato-lanceolatis alternis, petalis acuminatis
2. parviflorum C
3. ~~fructabile C~~
4. album C foliis ovato-lanceolatis caulinis, petalis obtusis
5. japonicum C. foliis caulinis subrotundis suboppositis
6. bulbiferum C. folio ovato radicali
7. acule C. foliis oblongis radicalibus

spectabile C. corolla labii superioris ovali basi retuso concavo sub  
tus carina obtusa, inferioris petalis longioribus grossis. Salisb.  
act. soc: linc. 1. pag 70. tab. 3. fig. 3.

Helleborine flore majore purpureo Mons. hist. 3. p.  
400 s. 12. t. 11. fig. 17.

Helleborine calceolus dicto, mariana, flore gemello  
candido, semi purpureo striato Pluk. mant.  
p. 101. tab. 440. fig. 3.

umbriata Mallea

~~Fungus~~ ~~gestr.~~ de seminebus cent. 3. Tab. 35. fig. 5.

non v. d.

*Helleborine flore majore purpureo etc.* Monf. hist. 3. p. 480. sect 12.

tab 11.

Habebat herb. mus

habes

Helleborus mons. hort. tab. 11. fig. 16

Habitat herb. Musci

habes

*Helleborus virginiana bulbosa* flore atro rubente Pluken. alm. 102  
Habitat in Virginia herb. Kauf.

habeo

alternis

Calceolus Cypripedium foliis caulinis ovato-lanceolatis, petalis acuminatis  
Calceolus mananur ~~God. pempt. 100 fig. 1. 2.~~

Helleborine flore rotundo f. calceolus ~~Hauk. Pin. 107~~

Calceolus foliis ovato lanceolatis ~~Gmel. pb 1. p. 2. tab. 1~~

Cypripedium calceolus radicibus fibrosis, foliis ovato-lanceolatis cau-  
linis ~~lin. 1. V. pag. 277. 8l. Dan.~~

vestiment ad  
Casi: parviflorum  
salus

~~Helleborine calceolus mananur dicto, caule folioso, flor. luteo~~  
~~re ~~H. mant. 101. tab. 10. fig. 2.~~~~

Helleborine virginiana f. calceolus flore luteo majore ~~Monf. herb. 3.~~  
p. 400. f. 12. v. 11. fig. 15.

~~Calceolus mananur canadensis Corn. canad. 204.~~  
~~Helleborine flore majore purpurea Monf. herb. 3. p. 400. f. 12. tab. 11.~~

vestiment ad  
bile salus

~~fig. 17.~~  
A. Calceolus minor, flore varo ~~Amm. ruth. 133. t. 22.~~

Calceolus foliis ovatis binis ~~Gmel. pb 1. p. 5~~

Habitat in Europa, Asia, America septentrionalibus &

in herb.

in alpinis altari-

Radicibus fibrosis. Caulis peralis glaber ubi omnia. Folia tria vel quatuor  
in caule ovata acuminata, amplexicaulia, nervis 7. elev. a horibus basi  
acuta. Pedunculis terminalibus pluribus interdum <sup>in floribus</sup> ~~quibus~~  
alternis foliorum ellipticum versus apicem flor. nutans

petala lanceolata attenuata

Cypripedium calceolus labio superiore ovali concavo subtus corno luteo  
canalicula<sup>ta</sup>, inferiore petalis brevioribus compressis Salub. in Art.  
Sicil. Len. 1. p. 26. Tab. 2. fig. 1



Spectabile C. corolla labii superiore ovali basi retuso concavo subtus carina  
obtusius inferiore petalis longiore grosso Sal. in h. Tr. l. p. 70. v. 3. f. 3.

\* Cactu bot. mag. pag. 276

Helleborus calceolus D. R. manana, flore gemello candido, venis  
purpureis Anato Platin. mant. p. 101. v. 410. f. 3. presuma

Cypripedium album

Hort. Kew. 3. p. 303.

Habitat in sylvis pennsylvanicis

Sal. bony huc quoque vocat Hall. etc. Monf. 3. f. 12. v. 11

Calceolus Cypripedium corolla labio superiore ovali concavo subtus canna late  
canaliculata, inferiore petalis brevior compresso Galub. in L. S. p. 76.

Tab. 2. fig. 1. 7

parvi florum P. corolla labio superiore sagittiformi basi deflexo subtus  
canna anguste canaliculata, inferiore petalis breviora compresso  
Ralsbury in L. F. 1. p. 70 Tab. 2. fig. 2.

Helleborus caliculis dentatis, manana, caule folioso, flore luteo  
minore Plukh. mant. p. 101. t. 4 10. fig. 2. presuma

Habitat in Virginia

caule Cyp. foliis oblongis radicalibus Hort. kew. 3. p. 303 Curtis botan mag.  
p. 192.

Helleborine caliculus dicta, manana, foliis brevibus e radice ex  
adverso prodeuntibus, flore purpureo Pluk. man. 101. t. 410. fig. 1.

Habitat in America sept. & nova scotia

Radix fibrosa

Cypripedium humile. Corolla labio superiore rhomboides acuminato  
latentibus deflexo subtus canna angustissima obtusa, inferos petalis  
longiore antice fuso. Salub. in let. sci. lin. 1. p. 74. tab. 3. 4

Caliculus flore maximo rubente Cat. bot. Car. app. p. 3. t. 3

~~Helleborine flore maxime purpureo Mon. bot. 3. p. 400.~~

~~f. 12 + 11 fig. 17~~

bulbosum *Cypripedium folio <sup>ovato</sup> subrotundo radicali* Lin. J. V. pag. 217

*Cypripedium folio subrotundo fl. Capp. 319. t. 12. fig. 5.*

*Scapulus scapo unifloro Gmel. sub. t. p. 7. t. 2. fig. 5.*

*Cypripedium bulbosum sub bulbos subrotundo, folio subrotundo  
radicali* Lin. J. V. p. 217.

Habitat in Lapponia, Russia, Sibiria &

Radix bulbosa una cum pari scapi anni precedentis persistens  
in alterum annum ut in *Ophryde* Coefelii. Solum peticola  
tum ac ut in *scapulo*, nervi quinque absque nervis interstitiis  
et nervis. glabra uti tota planta pollicari. Petiolus  
longitudine folii. Scapus digitalis squamis duabus tubulosis  
oblique truncatis membranaceis pollicaribus inferne & ceteris  
superne nudus ap. versus apicem foliolum coloratum  
angustum lineare. flori erectus. petala 5 lanceolata  
attenuata incarnata. nectarii labium latum medio pilosum

Labium nectarii laevifolii descriptum

*Japonicum* *Cynripedium* foliis caulinis subrotundis suboppositis Thunb.  
*Cynripedium japonicum* foliis caulinis suboppositis nervosis Thunb.  
fl. japon. 30 Lin. b. V. p. 217

Habitatio Japonia

Caulis teres, villosus, pedalis. Vicia in medio caule duo, am-  
plexicaulis, acuta, pinnatis-undulata, nervosa, glabra  
palmam lata. Infra florem foliolum sessile, solitarium, oblon-  
gum, acutum integrum, via pollicare. Flos solitarius, mag-  
nitudine floris *Calceoli* Thunb.

non vid.