

Elaeodendron

orientale L.

argan Elodaeus. ruminis spenofis; foliis cratis obtepi Retz off. p. 26*

Habitat in Mau regno Maroccano.

cuneifolium Elaeodendron glaucum oblongo cuneatis integerrimus

Habitat in Domingo? Leb. Ausf.

Rami Veretos; superne tetragoni, dictata planta glabri. Dolia breviter petiolata,
obovata vel cuneiformia, basi acuta, apice ^{obtuse} emarginata, nitida, pallidior.
Pedunculi axillares; folio breviorer; tenuissimi; partiales racemosi, dichotomi

Prubentia

Habitat her. Gus.

Prami oppositi, uti reliquo parte glabri, inferne verete, superne compressi. Solum petiolata, opposita, bi-3 pinnata, oblonga, basi acuta, apice obtusa, glandula fū scia terminata, crenato-serrata, serrulati fū glandula fū fer, margine crassiora, supra nitidula. Pedunculi axillares, folio brevioris. partales racemosi, dichotomi. Cneter magnitudine nucis Coryli, oblong^{us}, basi acutus.

Folia varie forma secundum statem, sic in individuo
mo, unius acuto, remote denticulata, alia duplo lateora coetenum exoden
in foliis.

Est species Elegans

coriaceum *Elaeodendron* ^{nermis} foliis obvatis crenatis, pedunculis de hotores
Rubentia

Habitat in Apula Francia et Hispania

Rami recti sparsi glabri superne quadrilateroni folia
petiolata alterna opposita plerunque obvata
intervalem oblongo lanceolata 2-3 polliciana caes
mucrone minute pectio, utrangle glabra, coriacea subtilis
venosa, obtusa. Pedunculi attollentes foliis breviter
detinentes

Facies Pelastri an Almanni sit et *Elaeodendron* argon

Elaeodendron a Java foliis obvato-oblongi face et elastri habet nisi vel huper
generis

Conspicuita ramis levioris foliis pallidis obvatis fructu magnitudine sem. Concord
vel huper vel Pelasti generis

orientale Elodendron Gauguin Act. helv. et. g. p. tab. fig. 2.

Habitat.

non vidi

Sphenopum & *Sphenopum*, foliis perennantibus Lin. I. V.

~~Lycii similes fructus indicis Com. Hort. t. p. 164 f. 83.~~

~~Lycium maderae planum byzantii & gracile inibus fructu horridum Bluhn.
alm. t. 35. t. 202. fig. 2.~~

~~Rubra - molli Rhod Mal. sup. 77. t. 39. Rajahst. 1634.~~

~~Habitat in Malabaria & Barboria~~

~~Fruitus 4 vel 5 pedali ramulis aculeatis. folia oblongo rotunda crassa
densa vix crenata, nitida atrovirens, subtruncata dilatatoria nervosa,
flores subcrenides graveolentes. Bals. primum virides dein
rubro, tandem nigro, accido Wettm.~~

S. Sphenopum meum ex Hort. Parf. an Lin: Rami ^{erecti} glabri. folia
petiolata aggregatas, ^{in quinquefida} Lycii, unguicularia basi acuta obtusa, gla-
berrima, inde per rimos pupura vix venosa. Petiolari brevis. Fru-
ta ad latius gemmo rugosus subulatus, purpureus; foliis dimidiis brevior
flores in axillis foliorum subpletari parvi

Elaeodendron argan. folios ^{lignos} crenato Lanceo cuneiformes integri
m. florum spissibus

~~in fructu Bluhn: fructu oppositae in mea ali. foliis crenatis foliis
alternis alium ex 4nd orient habeo opposite decussatis sphenopum foliis~~

bui cymofis non vero in omniibus fere Liderotylei forte ~~ne~~ non ha
non ad Liderotylei genio pertinent tales Calastor. qd' amandare