

# Epigaea

1. repens L. foliis cordato-ovatis integris planis appressis, corollis cylindricis
2. cordifolia L. foliis cordatis subrotundis serratis convexis hispida rigidis, corollis ovatis

cordifolia Eryssa

Genethera Sphagnicola Richard.

Habitat in Canibus herb. Fusf.

Caulis fruticosus, repentes, ramosi: Ramis foliis, hirsutis. Foliois vix semipollicariis, ovato-cordato, rigida, arenis, acuta, serratis, serraturis serratis, subtus venosa, pallidiora, utraque foliis nigris raris adspersa. Racemi terminalis sessilis; foliois vix longiores. Calycis foliola ovato-lanceolata, glabra.

repens Eryosa Lin. Am. Acad. 3. p. 17 \* S. V. p. 407.

*Pyrrola affinis repens* fruticosa, foliis rigidis scabritie exasperatis, flore pentapetaloido, Litu  
loso Pluk. alm. 309. t. 107. fig. 1. Raj. suppl. 596.

*Arbutus foliis ovatis integris, petiolis laxis longitudine foliorum* Gron. Viny. 49.  
Memeçillum Mitch. gen. 13.

Habitat in Virginia, Canada punctis 5

<sup>suffruticosa</sup> ~~humilis~~ <sup>basifera</sup> ~~repens~~ <sup>ramosus</sup>. Ramis tenues, <sup>nisi</sup> ~~pollicares~~, <sup>vel</sup> ~~pollicares~~, <sup>et</sup> ~~pollicares~~. Folia rariore planta mag  
na <sup>vel</sup> ~~pollicaria~~ <sup>et</sup> ~~ultra~~, <sup>ovata</sup> ~~integerrima~~ <sup>subternata</sup> ~~acuta~~, <sup>basifera</sup> ~~non~~ <sup>venoso</sup> ~~reticu~~  
lata, costa subtus pubescenti. Flores terminalis 40 <sup>in</sup> ~~racimo brevissimo~~ <sup>pedunculat.</sup> Bractes ovato  
~~condens colorata~~ ~~enugitudine foliorum~~ ~~calycis~~ ~~adnatis~~. Calyx glaber coloratus  
exterior triphyllus foliolis ovatis acutis velosis <sup>omnis</sup> ~~exterior~~ brevior

repens E. foliis cordato-ovatis integris planis appens; corollis cylindricis  
Sw. prodr. pag. 74.

~~Bractes calyx exteriora lanceo acuminat cum a flore remoto sunt patius bractes~~

Cordifolia L. foliis cordatis subrotundis serratis convexis hispido rigidis,  
corollis ovatis In. prod. 73.  
Habitat in Guadeloupe 5