

Euphorbia

x Truncosa aculeata

1. antiquorum E. aculeata subnuda triangulans aciculata: ramis patentibus
aculeata
2. canariensis E. aculeata nuda squadrangulans: aculeis geminatis
3. heptagona E. aculeata nuda septemangulans: spinis solitariis subulatis flori-
feris
4. mammellanis E. aculeata nuda: angulis tuberosi spinis interfractis
5. carciiformis E. aculeata nuda multangulans: spinis solitariis subulatis
6. alibiinarum E. aculeata nuda multangulans: aculeis geminatis
7. triaculeata E. aculeata nuda, caule tereti sulcato, aculeis ternis
8. nereifolia E. aculeata: angulis obliquis tuberculato.

aleppica E. umbella quinquefida: dichotoma, involuellis ovato-lanceolatis
mucronatis, foliis inferioribus petalici len. L. V

Tithymalus cyrenicus Alp. exot. 65. 7. 64.

Tithymalus foliis inferioribus capillatissimis: superioribus myrto similibus
Morr. hist. 3. p. 330.

Habitat in Syria, Aleppo 2

maculata? Euphorbia
caulis pilosus
convenit satis cum her. folia antedum pubes pilis raris
plenuque vero glabris utroque. flores plures axillares
calycis pilosissimi.
Depressa Dr. v. Rohr n. 115

Euphorbia N. Engl. 197

menina Venicalli fl. Spec. pl.

arbuscula *Euphorbia* foliis oblongis integerrimis subserratisque, pedunculis axillaribus ^{terminalibus}
~~seminatis~~
~~petiolatis~~, caule erecto.

Habitatio India Orientali Herb: Biblioth. Christianav.

Tota planta Caulis suffruticosus, semipedalis, statim a basi ramosus, foliis
filiiformes nodosae, articulatae. # folia brevissima petiolata, vix semipollicia

purpurascens ser na, acuta, basi altero latere breviora, ~~trinervis~~ ^{obsoleta} trinervis, areolaria
vix glaucescentia # Pedunculi terminales ramulorum; foliola duo sub

plenique integerrima vix ad altum obscure latius extrosus feratura 23 obscura
singulis flore eoque longitudinis, semillimi foliis caulinis ut magis versis
apicem serrata, florem ante evolutionem involucentes. Petala semi-
obscure. Capsula bicondru minor, glabra, laev.

Simpler Euphorbia foliis ovato linearibus remote serrulatis, pedunculis axillaribus

Solitariis

Habitat in India Orientali Herb. Biblioth. Christian. 0

Tota glabra. Caulis erectus, simplicissimus, raro ramo uno vel altero
tereti, ad articulos nodosus, purpurascens, lavis. Folia subpetiolata,
remota, sesquialternis, linearia, basi rotundata, angusta, utrinque
serrata, altero latere frequentius et fere usque ad basin altero, parvis
et ut nisi ad medium, obtusa, mucronata, venis enervi. Pedunculi
brevissimi, uniflori, Patula puberulata, alba. Capsula mucronata, glabra.

rubra Euphorbia foliis cuneatis emarginatis imbricatis umbellis trifidis, corollis pentapetalis
Ravan: c. 1. pag. 21. tab. 34. fig. 1.
Habitat juxta Amoyes C.

Salicifolia Euphorbia

Hungaria ^{in altis} ~~in~~ ^{bipedales in 1580 m}
Caules plures, simplicissimi ^{superne in ramis simplicibus}
~~et ramosi floriferi~~ ^o ope lentis pilis raris brevissimis adpressis
Si folia sessilia ^{sesquipedicari} primum patentia dein
reflexa lanceolata, basi attenuata, ² vixtus ¹ curvata, integerrima
nervosa supra ope lentis costa et margine ^{per} ope lentis
^{subpilis} pubescento subpilosus, subtus frequentius vixtus ^{sub-glabrescentibus}
Umbella terminalis multifida bifida, ^{prostratis} laterales bifidae ^{rad}
In nodis ⁴ ¹⁰ phyllon foliola lanceolata involuella
cordata ^{integerrima} ^{acuminata} Petala 4 bicornia vixtus flavescentia

en ~~de~~ Hoffne
Godefruct

Gaertner Josephus De fructibus et seminibus plant

~~H Verhandlung von het Bataviasche g~~

~~obscuritatem~~ ~~Edenbr~~
~~Petrus Penna et Libet~~

~~Hofner~~ ~~medu w tetrad~~

~~Adversaria~~

~~in Turpender~~ ~~Sole~~ ~~mentha~~ ~~botan~~

~~Calculus~~ ~~ad~~ ~~Sancti~~ ~~Bar~~

~~Herbich~~ ~~in~~ ~~Bath.~~ ~~1779~~ ~~de~~

~~gand~~ ~~herbal~~

~~muller~~

~~Glacivert~~

~~Hannovers~~ ~~magaz~~
~~Journal de Phys.~~

~~ut~~ ~~de~~

~~abr.~~ ~~Munting~~ ~~noan~~ ~~keurige~~ ~~beethen.~~

~~Verhandl. von het gewest schap te Vless.~~

~~Pluk~~ ~~de~~ ~~de~~ ~~de~~

~~Meyer Phys~~ ~~de~~ ~~de~~

~~Branch~~ ~~prodr.~~ ~~de~~ ~~de~~

~~Op~~ ~~de~~ ~~de~~ ~~de~~ ~~de~~

~~anale~~ ~~de~~ ~~de~~

~~met~~ ~~de~~ ~~de~~

~~Tru~~ ~~de~~ ~~de~~ ~~de~~

~~Commerc~~ ~~de~~ ~~de~~

~~Haller~~ ~~de~~ ~~de~~

~~Historia~~ ~~de~~ ~~de~~

~~Rombard~~ ~~de~~

~~de~~ ~~de~~ ~~de~~

~~Hat~~ ~~de~~

~~de~~ ~~de~~ ~~de~~

~~at~~ ~~de~~ ~~de~~ ~~de~~

~~de~~

~~Born~~ ~~de~~

~~de~~

~~Mayer~~ ~~de~~ ~~de~~ ~~de~~

~~de~~ ~~de~~

~~de~~ ~~de~~ ~~de~~

1. *antiquorum* L. aculeata subnuda triangulari articulata. ramis patentibus

Hort. Cliff. 1764 Am. acad. 3. p. 106 x

Schadica calli Rhed. mat. 2. p. 11. v. 42.

Euphorbia antiquorum venim Com. hort. 1. p. 23. v. 12.

β. *Tithymalus airoides* triangulari et quadrangulari articulata
et spinosa; ramis compressis Com. prol. 5. 5. v. 5.

Euphorbia trigonum et tetragonum spinosa ramis compressis
Afr. act. 1720. p. 305. n. 2.

Habitat in Cap. i. spec. 5

Caulis angulis tribus marginis repandis fere articulatus. Rami
horizontaliter angulis superioribus horizontalibus, margine
repandis; angulis inferioribus. Aculei duo oppositi, et supra hoc
foliolum minimum ovato-lanceolatum, deciduum haec
folia videntur a dentibus caulis interdum simplicibus, collis una
cuneata hest. fus.

aculeata E. cauli fistulosi adnatis subrotundis D. r. fl. a. p. n. 94
Habitat in Arabia

non vid.

canariensis ^{aculeata nuda subquadrangulans: aculei geminati Hort. Cliff}
1967 ^{acul. 3 p. 1074} ^{Lin. Mant. 394}

Tithymalus airoides fruticosus, canariensis aphyllus quadrangulans,
spinis geminis aculeis abronchitibus armatis Comm. hort. 2.
p. 207. t. 104.

Tithymalus airoides lat. folius, f. Euphorbia canariensis quadrilatera
et quinquelatera ceni effigie Planch. aln. 370. t. 320 fig. 2
Habitat in Canariis

Caulis terribus induratus, interdum quinquangulans, aut angulis
compresso-planis. spinis duobus oppositis armatus. folia nulla
dentata.

flores non vidi.

Tribuloides Euph. humillima daphylla aculeata, quadrangulari, spinis
geminatis patentibus La Mark En.
Habitat in Canariis &

non vid.

3 heptagona L. aculeata nuda septem angulans: spinis solitariis subulatis floriferis
Hort Cliff. 1764 Am. acad. 3. p. 1094

Euphorbia cuneata, spinis longis simplicibus Bridl. Succ. 2. p. 4.

7. 13.
Euphorbia heptagonum, spinis longissimis in apice frugiferis Boeck.
Leydb. 1. p. 250 & 250

Habitat ad Cap. b. spec. 5

Caulis hinc inde ramosus. aculei palli, rigidi solitarii, longitudi-
naliter ex angulis singulis dorso ciliatum prodeunt, Flores ex
apice aculei interdum prodeunt sed solitarii, in

4 mammillari E. aculeata nuda: angulis tuberosi spinis interstinctis Lin. Sp. Pl. pag. 647*

Tithymalus aizooides africanus, validissimus spinis ex tuberculorum internodiis provenientibus Comm. prol. 59. t. 9.

Euphorbum polygonum, aculeis longioribus ex tuberculorum internodiis procedentibus G. a. 1720. p. 386.

Habitat in Africa

differt ab Heptagona angulis duplicatis torulosis; spinis solitariis inter singula tubercula longitudinaliter digesta Lin.

He non vidi

5 Cerei formis l. aculeata nuda multangularis: spinis solitariis subulatis l.
f. v.

Tithymalus africanus, spinosus cerei effigie Mont. bot. 3. p. 245. Plukn
alm. 370. t. 231. fig. 1.

Euphorbia aphyllum angulosa, flosum comadenissima Burm. afr.
14. t. 9. fig. 3.

Habitat ad Cap. b. spec. 5.
flos in apice copiosissimi

6 officinarum *E. aculeata* nuda multangularis: aculeis geminatis Hort. Cleff.

1764. Am. acad. 3. p. 1074. L. V.

Euphorbium Bauh. pin 387.

Euphorbium cerei effigie, caulibus crassioribus spinis validioribus armatum

Com. hort. 1. p. 21. 8. 11. Tab. thes. 1. p. 24. 8. 14. fig. 2.

Euphorbium polygonum spinosum, cerei effigie Sp. act. 1720.
p. 385. 6. 10.

Habitat in Africa calidioribus.

Caulis erectus, teres, squamis simplex, sulcis octodetum aut pluribus longitudinalibus et aratus. Aculei ubique duo. Folia nulla, et eorum loco tuberculum acuminatum, supra et juxta aculeum par, et cum adhuc tenelli sunt aculei, eandem fere ac folia ferunt faciem sed solum contabescit aculei vero indurantes hinc.

abyssinica E. nuda multangulari, aculeo quinque Bruci: Triv. in Egypto S. ap. 1770
42 u.

non vidi

Fruct. of

E. multangula subpedali, caespitibus pedunculatis Forst. fl. Oryz.
nuz 94.

non di.

neriifolia E. aculeata: angulis oblique tuberculatis Hort. Cliff. 1964 W.
Am. acad. 3. p. 1094
The *Stygnalus airoides* atropicensis *spinosus*; caudice angulari, *neriifolio*
Com. prol. 22. 56 t. 6. Brad. succ. 3. p. 10. t. 20.
Tithymalus indicus spinosus, neriifolio Com. hort. 1. p. 25. t. 13.
Ela Calli Rheed. mat. 2. p. 83. t. 43.
Ligularia Rumph. amb. 4. p. 88. t. 40

Habitata India

Caulis teretis scaberrimus, quinquefarius nam Tuberculis prominulis an-
gularibus, angulis his spinescentibus lente adscendentibus, in sum-
mo caulis. Folia lanceolata, ad quorum basin duplex alu-
leus, foliis deciduis persistunt aculei cum cicatrice et
callo prominulo. L.

fimbriata h. angulata angulis aculeatis ad bos. subnoveci, fimbria Verm
natura fimbriata bruleatis / cep. del. fl. inf. 3. p. 27 47

non vid.

linanifolia E. umbella ~~multifida~~ dichotoma, involuelli subcordati obtusi mucronati, foliis unifloris ~~ramis nullis~~

Euphorbia

triaculeata Euphorbia aculeata nuda, caule tereti sulcato; aculeis ternis
 Euphorbia triaculeata; caule tereti, septem sulcato; floribus ternis septi-
 bus ex ala aculeorum Cosch. descr. pag. 94 ubi descriptio habetur
aculeus inter medius pollicaris, laterales quadruplo minores,
 recurvi

Lonicera Euph. aculeata femina, caule tereti squamato, basi pennis foliato
nis mucronato, foliis lanceolatis cuneatis La Mark. Enc.
Tithymalus africanus et. Plukh. Alm. 370. Tab. 230. fig. 5.
Habitat ad Q. b. per