

Genippa

1. laterna G.

Genippa americana Desf. Ryan. folia elliptica, laminolata, basi sensu attenuata

eripula *Genippa*
glomerata *Mussenda* foliis ternis oppositisque elliptico subobovatis, floribus
terminalibus congestis subsessilibus

Anonyma *Merian.* *Surinam.* *men.* 7. 43.

Suria *Eriopula* *Lin.* *Suppl.* p. 209.

Genippa americana *meriana*; omnibus partibus hirsuta, foliis obla-
go-obovatis, floribus in summitate congestis, fructu depresso globoso
hirsuta *Ait.* *Bot.* *hist. nat. Paris.* 1. p. 107.

Mussenda glomerata; foliis ovatis acutis, floribus terminalibus, glo-
meratis *La Mart.* *Dact.* *bot.* 4. p. 393.

Habitat in *Surinamo*, in *Guiana* *Richard*, *La Mart.*, *Trioli* *Marmola*
De Boossee Boom dicta. *Fructus* *sepidus* $\frac{1}{2}$

Rami *superne* *triquetri*, *juniores* *angulati* *subpeltati*, *epili* *adpressi* *hirsu-
ti* *folia* *prexipua* *petiolata*, *pedalis*, *tertia* *minore*, *elliptica*, *basin* *versus*
angustata, *juniora* *vellosa*, *stata* *glabra*, *costa* *subtus* *hirsuta*, *supra* *obscure*
venosa, *subtus* *costata*, *pulchre* *venosa* *reticulata*, *acuminata*. *Pedunculi* *brevissi-
mi*, *uniflori*. *Calyx* *multicostatus*, *denticulatus* *subulatus*. *Corolla* *extus* *hirsutissima*
basin *apice* *que* *genicis*, *non* *pollicari*.

~~rhomboidale
setangulare~~ Solanum venenosum caule ramisque aculeatis, foliis oblongo-ovatis
integris angulatoque, floribus tetrandris

Habitat ad littora maris Martineque Islet

Caulis herbaceus vel suffruticosus, ramosus, teres, uti tota planta tomento
denso flavo vestitur. Aculei sparsi in caule et ramis, unus vel alter
subtus in costa folii, recurvi, glabri. Stipula petiolata, geminata, sub
sessilis, basi angustata; altero minore, absque angulis, tripollicare;
altero tripollicare utrinque angulis duobus remotis absque si
mis, obtusa, crassa, scabriuscula. Pedunculi supra foliacei: pe
dicelli 3-5: Flores parvi. Calyx inermis. Bacca magnitudine semi
ni Conarii.

ni fallor aderat quoque in herb. Mus.

Catena / Genespa.

Habitat in Guinea.

americana Genipa Lin. S. V. pag. 227.

~~Genipaba Mangr. braj. 92. Pl. Bras. 150.~~

~~Genipa fructu ovato Plum. spec. 20. Pl. 136.~~

~~Habitat in America Meridionali~~

vellofa Genyspa foliis oblongis vellosis subtus incanis, floribus terminalibus aggregatis
pedunculatis

Habitat ad Senegal herb. Guisidi et Dupui

Rami teretes, oppositi, si abruis ^{no} sculi, superne vellosi. folia ^{no} apicem ramorum, opposi-
ta, conferta, bipollicaria, lato-oblonga, acuta, basi versus partem angustata, utrinque
vellofa, supra nervosa, subtus subnervosa, incana, profertim in nervis et costa. Squama latis
stipularum oblonga, geminae, obtuse, glabra, substriata, caduca. [#] Stipula ovata, attenua-
ta. Pedunculi in apice ramorum aggregati, plures, breves. [#] Petiolus unguiculata
res-incanai. Polyis lacinae breves, subulatae. Corolla glabra, sesqui pollicaria.
Tubus superne vellosulus. Anthera inter lacinas, alba, lacinis brevior. Stylus
corolla duplo longior. Stigma bilobum

Petiolus unguiculares incani

Canocolata Genippa

Habitat in Cayenna herb Turf. 5

Rami teretes, glabri, cæcaneati, oppositi, superne nodosi, folia versus apices
ramorum, breviter petiolata, 2-3 pollicaria, canocolata, attenuata, tenuissime venosa, ne-
tida, integerrima. Sepala ovata, glabra. Flores terminales, plures, sessiles, calyx
quinque dentatus, minutus. Corolla angustulans, extra torrensosa.

ab Duli forte non diversa

oblongifolia Garden Genappa

Genappa oblongifolia; foliis oblongo-^{ob}ovatis subsessilibus, racemis terminalibus & R. Penn. 2. pag. 67. Tab. 220. fig. a

Habitat in Andium nemoribus in eis calidis &
Arbor triquadriangulis. Truncus erectus. Rami teretes; e stipulis densis
Dens ciliolato-annulato. Folia opposita, conferta, breviter petio-
lata, tripalmata et ultra, undulata, oblongo-obovata, obtusa
integerrima, supra nitida, subopacea, venosa, nervo et venis
transversalibus leviter pubescentibus, marginibus revolutis. Petioli
an gardenii brevissimi. Stipula interfoliacea, semianplectens; petioli longior;
Species membranacea, ovata, acutisima, decidua. Flores terminaliter
longi, breviter pedunculati, basi bracteis duabus lanceolatis.
Calyx tubulosus, leviter 5-quefidus, persistens. Corolla lutea in
fundibuliformis hypocrateriformi. Tubus herbaceus. Anthera ova
fo lanceolata, incumbens. Germen oblongum. Stigma oblongum
acuminatum, stratum, longitudine fere corollae. Bacca ovata,
magnitudine mali persici, biloculari, basi calycis coronata.
Semina numerosa, subrotunda per 2 generi in singulis loculis
compressa

Edulis Genespa glaberrima, foliis oblongis, floribus in ^{summitate} ~~summitate~~ congestis
plerumque quadrifidis, fructu globoso lavi acti sic hem. Parf. f. pag

107.

Habitat in America meridionali in herb. Rich.
flores in apice sive plura sive riles hermaphroditi; in alio ramis soli
tanti fertiles Rich.

Cousoarea Aubl. fucus gnyssa Richard

^{Pohna}
Pseudogenespa hujus generis, Rich. genespa inflata dicta flores uterque
et interdum sive fidi confer. cum Cousoarea Aubl.

~~name~~

menana Genappa omnibus partibus hirsuta, foliis oblongo^{ob}ovatis, floribus
in summitate congestis fructu depresso-globoso hirsuto Act. for. port. pag
107.
Habitat in America h. b. Aub

Uxi de marmelade uxioli Du Rora Linnæi sanat familibus 5. 6. 7.
interdum polygamum. Non rarum in subtracis fructu baccato quod na
men. ~~se~~ american sanat
~~non~~

buffalena Genappa. foliis papilionatis ovatis glabris, floribus paucis lous-fl.

Cochinch. pag. 149.

Habitat in Cochinchina

Arbor 9 pedalis. aciculi oppositi longi, recti. Folia catenarii glabra, plana, dura. flori solitarii albi. Calicis calycis rotundato. Bacca exsucca. Corolla rotata. Stigma ovatum bifidum

non vidi

eficulenta Genappa folio fasciculato ovato piloso, flentibus congestis Lour
fe. corh. pag. 144.

Habitat in Cochinchina 5

Arbor, parva 6 pedali. Folia utriusque parva fasciculata opposita.
aculei longi recti oppositi. flores laterales albo-vires. Corolla rotata
et fida. laciniis oblongis acuminatis. Stigma oblongum emarginatum.
Bacca teris aequali corobata 1. Corulata.

flava Gemppa! foliis lato-lanceolatis, flore Verminali solitario linc-fl.
Coch. pag. 149.

Habitat Cantona. 5

Index, spudalis. aculsi spasi pauci recti, folia opposita, integerrima, glabra.
flore lutei. Corolla rotata, crassa subpelop. Stigma subrotundum mag-
num angulatum sespale.