

Gladiolus

90 Gladiolus. Cor. 6-partita, laciniis limbi inaequalibus, tubo subcurvato. Stam. ascendente

antholyza spicata
prope dichotomiam

Scapus erectus & teres

Rachis flexuosa

gladiolus involutus La Roche

4x2

gladiolus gladiola in la m...

Vogel gladiolus cristatus pl. rar. deias

27. 28. 4.

gl. inflatus
permacabris

De S. Buzmann

cum Haec que collata

gladiolus pulcherrimus
la m... forte n...
thum

* Scapo monostachyo

montanus 1. Gladiolus corolla ringentis tubo & spathae obtusae aequali, laciniis pra
 thulatis, foliis ensiformibus nervosis

gladiolus montanus; foliis ensiformibus nervosis glabris, floribus, spicatis,
 corolla ringente. lin. suppl. 95.

gladiolus montanus; corolla ringentis tubo & spathae obtusa aequalibus, foliis
 ellipticis nervosis Thunb. Diet. Glad. n. 1. t. 1.

Habitat in planitie montis tubularis Cap. 6. spec. 3

Gladiolus folia radicalia, subpedalia, vix semiunguem lata, vel angustiora, inferne
 sensim attenuata, trinervia, glabra. Scapus foliis parum longior, ~~erectus~~
 subflexuosus, ~~terres.~~ Flores distantes. Spatha limbo obtusissimo
 scenter. Corolla unguiculans: tubo & spathis brevior. laciniis ^{superior} ~~superior~~
~~Canaliculata, apice ovata; laterales, patentes, divaricatae, re~~

Gladiolus par
 riflorus; foliis
 linear-ensifor
 mibus; caule pim
 plia & semis; flori
 bus spicatis Paige
 obs. 4. p. 2. t. 70

obovata; ~~obovata~~ ~~long~~ ~~labiam~~ ~~tripartitam~~, ~~aequali~~ ~~laciniis~~ ~~aequalibus~~,
~~obovata~~: laciniis ~~spathulatis~~; superior fornicata, canaliculata, maie
 la purpurea; laterales patentes, divaricatae; inferiores tres aequales, usque

luteus Glad ~~folus lineatis~~ angustis longispinis, floribus spicati subpediculis
curvis, tubo curvo ~~protha~~ brevior La Mark. T. Enc. pag. 117.
Habitat in Madagascar

non vid.

Spicatus 2. Gladiolus spica imbricata disticha, scapo glabro, foliis linearibus

+

Gladiolus caule simplicissimo, ^{obis} floribus linearibus floribus spicatis

Boyer Lugd. 19.

Gladiolus spicatus; foliis linearibus, caule simplicissimo, floribus spicatis. Syst. veg. 87.

Gotia alopecuroidea; caule simplici, foliis linearibus, floribus imbricatis Lin suppl. 92

Gladiolus spicatus; corollis erectis, spica imbricata disticha, foliis linearibus Thunb. dia Glad. n. 13.

Habitat in Cap. b. spici 7

Scapus caulis spithamaci; erectus, 4 res. Folia quatuor, longe vaginantis, inferiora sensim breviora, angustissima, glabra. Flores cinister 12 ^{coerulei} duodecimparium, in spica pollicari. Thunb. l. c.

Differt a Gl. alopecuroide: floribus majoribus, spica breviori, scapo simplici. Thunb. l. c.

Spathaceus gladiolus polybrachyus, spathis membranaceis aristatis, foliis ensiformibus plicatis hirsutis Thurb. Desf: p. 22.

gladiolus spathaceus caule ramoso, floribus spicato-umbriatis, spathis membranaceis aristatis, foliis plicatis lina. Suppl. pag. 96. S. V. p

L

Habitat in Cap. b. p. 24

Scapus vaginatus, spithaceus. Folia nervosa, scapi longitudine. Flores albi.
Spathis multo longiore. Spica umbriata, digitales, plurimo. Spatha
sepe laevior. Similis fl. tubifloro et plicato. Thurb. l. c.

fistulosus 3 Gladolus spica disticha, foliis teretibus fistulosis Thunb. bot. schoen. t. p. 16.

^{7. 16.}
Gladolus turulosus, foliis cylindricis nervosis glaberrimis, florum spica disticha

Burm. prodr. Gladolus fistulosus; spica disticha; foliis teretibus, fistulosis Thunb. bot. schoen. t. p. 16.

2.

schoen. t. p. 16.

Habitat ad Cap. b. spec.

Folia crassitie pennae Cygnae & ultra, bi-tripollicaria, basi fixa ad us-
caria, ^{nervosa} striata, apice obtusa, furca, obtusa, mucronata. # Scapus vaginis

vaginae inter foliorum Textus. Spica palmaris & ultra, pyramidata, Flores copiosi
grae, parvae conferti, parvae apice ante expletionem florum teratiuscula.

Flores copiosi, conferti, parvi. Spathae ovatae, compressae, ^{denticulatae} dorso fuscae, latera-
bis membranaceo hyalinae, mucronatae. Corolla spathe parum longior

Spica exata gl. alopeuroidis, at tamen ab illa quam ab omnibus
reliquis distincta si ^{us} fit, foliis fistulosis, ad alterum apex modo
operculi deciduo,

+

caulescens aff. glad. spicati

Arctostaphylos foliis linearis lanceolatis villosis plicatis, tubo spathe equali Hort

Herb. 1. p. 63

a. corolla saturate coerulea, tubo et basi laciniarum e purpureo nigri

b. corolla pallide purpureo-pente tubo coeruleo

+ . . . Habitat in Cap. 6. p. 61

Garq. G. D. volu. nervosus: foliis ensiformibus plicato-nervi villosis, vate
multis pluribus alternis, tubo spathe brevior La Mart. Encegl

2. pag 715

Scapus erectus pedalis angulatus villosus, griseus, cetera vix
polystachyus. Folia ensiformia, basi breviora et nerva villosa
utrinque attenuata plicata. Rami floriferi albescenti, distincti
bracteae duae subulatae, breves ad basin pediculi rami. Spatha folium
exterioris linearis oblongum, obtusum mucronatum, nervosum villosum
longitudine tubi: interioris lanceolata attenuata tubum quan-
ter flores secundi tubo spathe brevior.

imbricatus 4 Gladiolus corollae imbricati species imbricata secunda, folia saepe
glabra, foliis pensiformibus +
Gladiolus imbricatus, foliis pensiformibus, floribus imbricatis Lin.
Spec. pl. 52. Georg. itar. 1. p. 196.

Habitat in Russia citeriore 4

Flores parvi versus unum latus imbricati Lin. l. c.
an species Antholyzæ? Thunberg.

Biflorus gladiolus corollae erecta limbo campanulato, scapo bifloro, foliis linearibus
Thunb. v. s. p.

Habitat in cap. bono p. 7

Scapus totus vaginis foliorum vinctus, erectus pedalis. Folia circiter tria
longe vaginantia, glabra, in summo brevissimo, supremo longitudine
scapi. Flores terminales, bini, coloris vincti Thunb. l. l.

umbriatus
spicatus
nerviiflorus

5 Gladiolus spica imbricata, scapo villosa.

Gladiolus spicatus; caule simplici villosa, floribus imbricato-
spicatis Lamark. illustr. 1. p. 120.

Habitat in Cap. b. spici. ex herbario Lamarkii

Differt a sp. spicato hic. caule villosa; spica undique imbricata;
floribus menoribus.

nigens

~~Gladiolus~~

~~Gladiolus nigens; polystachyus foliis plicatis hirsutis, corolla nigente,
spathis hirsutis Thunb. prodr. 106. Act. Soc. h. n. hagn.~~

~~Habitat in Cap. b. sp. 2.~~

~~Bulbus tenuis magnitudine avellano, tunicatus, alte insidens.~~

~~Folia digitalia usque perithamoe, scapum subequantia, subpetiolata,
ensiformia. Scapus brevior vel longior, palmari vel subpedalis.
raro simplex, saepissime polystachyus, flexuosus, superne villosus,
compressus. Spatha longitudine tubi, ovata, convoluta, pl. apice
membranacea et obtusa, toto valde hirsuta. Corolla caerulea, plus
minus albedo vel aduratiore, vel flavescentes: limbi laciniis divan-
cato-rungentes Thunb. Diss. C. 1.~~

flexuosus 6 Gladiolus corolla nigra tubo filiformi spatha lanceolata ^{longiore} ~~longiore~~ [#] ~~longiore~~

~~no flexuosus, foliis linearibus~~

#laciniis ⁶ ~~6~~ Gladiolus flexuosus; foliis linearibus, caule simpliciter flexuoso, floribus speciosis
nigra tubo longo Lin. Suppl. 96

Gladiolus flexuosus; corolla nigra tubo spatha acuta longiore, scapo
flexuoso Thunb. Diss. n. 2. t. 1.

Habitat in Cap. b. Spei 7

~~Spatha~~ ^{capitulum} pedalis, ~~caulis~~ ~~terres~~, versus bulbum valde attenuatus. Folia duo
vel tria, brevia, canaliculata, striata, vaginantha. Spica triflora, ovata.

Flores primum approximati, demum remotiores. Spatha lanceolata-oblon-
ga, attenuata, fusco ferruginea, tubo corollae parum brevior. Corolla
laciniis lanceolata

Differt ab insequente: scapo nigriore; flori foliis angustioribus:
locis spathis brevioribus, ferrugineis.

~~Gladiolus flexuosus; foliis linearibus, caule simpliciter flexuoso~~

flexuosus
Gladiolus

montanus gladiolus corolla argenti tubo spatha obtuse equantibus, foliis ellipticis nervo-
sis Thunb. Diss. p.

Synonymum riparium Planch. phyt. 224. fig. 8.

Gladiolus montanus foliis ensiformibus nervosis glabris, floribus spicatis, corolla argen-
te Lin. Suppl. pag. 95. * N. V. p.

Habitat in planitie montis tabularis Cap. b. spec. 4

Capus teres, flexuosus erectus, viridescens glaber, pedalis, solum, radicalia linea-
ria, trinervia, glabra, latitudine ^{vel penultimum basi attenuata} lineae, longitudine fere saepe spica
oblonga, floribus remotiusculis. Corollis spathisque obtusi. Bractea propria
bivalvis, ovata. Corolla tubus spathae brevior. Corolla galea tripartita, ma-
cula purpurea: lacina media fornicata, canaliculata, linearis, apice
obovato-lacinio laterales patentes, divaricatae, recurvatae, obovato
ungue longiorae. Labium tripartitum aequale: laciniis obovatis, ungue
longo, purpureis. Stamina sub galea lacina media Thunb. et Lin. l. c.

Signo cuius notato in W. d. invenit

in Spec. plantar. Synonyma huiusmodi sub gladiolo alata invenimur

ex Blaupus 7 Gladiolus corolla virgatis Tubo ^{phli formi} Spathis, ^{laciniis lanceolatis obtusis} ~~laciniis lanceolatis longioribus~~, & capo

flexuoso, foliis linear-ensiformibus
Gladiolus ex Blaupus foliis ensiformibus trinerviis tubo corollae fili
formi Spathis longioribus Thunb. prodr. 104. Act. Soci. h. n. hagn. vol.

6. t. cum in ore

Habitat in Cap. b. spaci

Scapus ~~caulis~~ ^{basi anivatione adscandens, demum} epithamosus, usque pedalis, ~~simplex, erectus, flexuosus,~~
ubi tota planta glaber. folia duo radicalia, ~~capo breviora, duo~~
caulina, ^{et infra} ~~margine et costa~~ ^{at altera} flavescentia. Flores duo-sex, erecti, albi
Spatha bipolliaris, linear-lanceolata, acutior vel obtusior.
convoluta, subtilissime striata. Corolla Tubus ^{in quinque et ultra} ~~in quinque et ultra~~ pollicaris, arcu
tus filiformis & ~~lamina lanceolata, obtusa~~

Dignoscitur a reliquis mihi notis longitudine Spatharum & Tubi
a praecedente differt foliis longioribus, latioribus: Spathis viridibus,
nec fuscis.

brevifolius & Glad. corollae nungentis tubo spathis ovato-lanceolatis aequantibus
trivinculatis, scapo aphyllis. Glad
Glad. Tristi saicth Thunb. L. n. 8 i. d. h.

Glad. brevifolius; corolla subringente; foliis ultra vaginas omnium
brevissimis; spica secunda, erecta 4. p. 156. G. rar.
2. t. 249.

Glad. brevifolius; monostachyus; aphyllus; corolla curva incarnata,
spathis cuspidatis levibus longitudine tubi Thunb. prodr. 105.
act. Soc. h. n. hof. vol. 6.

Habitat in Cap. b. Spei 4

~~1. Glad.~~ folia nulla sub florescentia, eorum loco vaginæ duæ

conlem ante vestitionem
suprema in
folium unguic
late acutum
minus attenuatum
desinent

quatuor, inferiores breviores, superiores longiores, striatas, apice obli-
que truncato, apice fusco. Scapus pedalis, ^{bipedalis} ~~erectus~~ ^{erectus} ~~indivisus~~, apice

carneus & flexuosus. Olexo spica palmatis, ^{tr} ~~quædam~~ ^{tr} ~~decem~~ ^{tr} ~~plora;~~

Olexo plerumque secundi, int. interdum distichi, incarnati, carnei.

Spatha bivalves, ovata [#] ~~car~~ [#] ~~unquidulares~~ [#] ~~sal~~ [#] ~~parum~~ [#] ~~ultra~~, [#] ~~ovato-~~
lanceolata, attenuata, longitudine tubi. Corolla tubus sub tubus
curvatus: limbi lacinia ~~longior~~ ^{longior} ~~superiores~~ angustiores

tubo quales

Setifolius Gladiolus polytachyus, foliis linearisetaceis Thunb. Desf. p.

Gladiolus setifolius, caule ramoso, corolla ringente, foliis infimo filiformisetaceo Lin. Suppl. pag. 96. N. V. p.

Habitat in cap. bonopei 4.

+ *Scapus rarisime simplex, flexuoso-erectus, glaber, palmatis. ramis teretibus, erectis
Folia circiter 3. semper breviora, infimo longissimo, scapum equante
Flores alterni albi, ringentes, tubo vix spathe longior Thunb. L. c.*

*Plantam a Dahlis accipi et a Thunbergio lecta quo forte setifolius Thunb
quamvis descriptis haud optime quadrat*

*Scapus semipedalis inferne teres superne flexuosus angulatus,
vix raris simplex. Folia spatheae convoluta teretia filiformia
glabra scapo breviora superiora semper breviora. Flores, alterni
distanti distanti. Spatha lineares 6 pollicari et obtusa
glabra striata. Corolla tubus capillaris 3/4 vel 4 pollicari
ns. laciniis limbi lanicolata*

non vidi nisi eadem cum meo galato

elongatus ~~9~~ Gladiolus. corollae nigrae tubo spathae longitudine spathae, lobis ~~4~~
~~9~~ Gladiolus tristis ^{corollae nigrae, virescenti albidis,} ~~var. 10. 11.~~ Thunb. Glad. no. 10. 11. plurimis Thunb. Glad. n. 10. 11.
~~9~~ Gladiolus elongatus; foliis quatuor linearibus bisulcis, corolla vire
 scente, spathae laevibus longitudine tubi Thunb. prod. 10. 11. aut.
Bot. h. n. havn. vol. 6
Habitat in Cap. 6. Spei 2

Totus glaber. Folia radicalia quatuor, scapum aequantia, uti tota planta glabra, su
perne bisulca, inferne sex sulca; duo spathae formis, sub pollicianis supra
medium blani, lanica, acuminata. Scapus bipedalis vel paulo ultra,
simplex, erectus, superne fleuosus. Flores septem vel octo,
erecti. Spatha tubo corollae vel longioris, lanicata. Corolla
six semipollicianis, virescens. Thunb. l. c.

~~9~~ Gladiolus tristis var. collected 4. p. 15 in var. tab. 244

+ 0

~~Gladiolus tristis var. 10. 11.~~

~~fimbriatus~~

t. 244 lacinae lanicae
 total obtusa

tenellus 10 Gladiolus corollae ringentis tubo spathae aequante, laciniis 4

lanceolatae
obtusae +

gladiolus tristis, humilis & luteus Thunb. gladiol. no 8. f. h.
Gladiolus tenellus; corolla subringente, foliis linearibus convolutis
glabri, caule flexuoso bifloro Jacq. collect. 3. p. 255. f. c. var. 2.
t. 248.

Gladiolus tenellus; monostachyus foliis linearibus basi longe, corolla
curva lutea, spathae strictae tubum aequantibus Thunb. prod. 105.

Art. bot. h. n. kafri. vol. 6.

3 Jacq. collect. 4. p. 169. f. c. var. 2 t. 3. f. 1.

Habitat in Cap. b. sp. 4

Scapus

Solum radicale unicum, longitudine ^{scapi} caulis, duo caulina, sensim
breviora. omnia ^{erectiuscula} linearia, acutiuscula. ~~caulis~~ ^{biflorus} semipedalis, subfloratus

2us, ad folia geniculatus, supra folium summum destinationatus,
gracilis, debilis. & l. resuncialis. Spatha diphylla, bival
vis; valvulis linearibus subaequalibus, linearibus lanceolatis
erectis, concavis. Corolla orbiculata subringens, linea longi
tudinali bordice violacea. Limbi laciniis duo patulis, ad san
tem punctato Jacq. l. c.

+ 9

M³

ve lanceolatae obtusae

vate

Caulis semipedalis.

β. Folia longitudine scapi, infera breviora, sequentia lanceolata, linearia, & tri-
tuba. Spatha flore dimidio brevior. Corolla laciniis lanceolatis rotundatis
limbo luteo. Fructus 4. l. l.

gracilis ~~et~~ Gladiolus corollae subringentis tubo spathae aequante, laciniis
concordatis

Gladiolus tristis, punctatus Thunb. Glad. n. 8. 9
Gladiolus gracilis; corolla subringente, foliis linearibus glabris,
caule superne geniculato, foliis alternis vestito subbifloro, foliis
altiora Jacq. collect. 4. p. 159. Ic. rar. 2. t. 246.

Habitat ad Cap. b. speir
Totus glaber. ^{scapus} Caulis ^{vel bi pedalis} pedalis, inferne foliosus, gracilis, debilis, ~~erectus~~
superne geniculatus. Folia caule breviora, longe vaginantia, supra
vaginam brevia, canaliculata, striata. Flores plerumque duo, raro unicus,
bivinculati, parum cernui. Spatha bivalvis, valvulis longitudine pau-
cis corollae, lanceolatis, com. acutis, concavis. Corolla tubus cylindricus
albus; faust incurva: limbi, laciniis pallide infundibuli formis,
pallide violaceis: laciniis ~~concavis~~, acuta, tria superiora duplo
latiora, linea saturatiora in dorso; duo inferiora striis punctisque
variegatis; intermedia linea unida picta. Jacq. l. c.

An Gladiolus punctatus, foliis linearibus, spathae acuta, petalis punctatis

+ 10 in sejoribus longioribus et acutioribus lamarck illud. 1. p. 117. huc pertinet

Gladiolus gracilis; monostachyus, foliis
linearibus multistriatis, corolla orate
coccolae, spathae longioribus tubi. Thunb.
profr. 105. t. 9. 802. h. n. 105. v. 6

floribundus G

non var

gladiolus floribundus Jacq. coll. 4. p. 162.

Habitat in Cap. b. sp. 4.

Solia plura ensiformia acuta, striata, disticha, altero margine fuso amplexi-
caulia, erecta, sempetalia. Scapus teres flexuosus supra sepe genicula-
tus, simplex foliorum sex longitudine totus a basi floriger. Flores
sexles, erecti, distantes, trunciales. Spatha bivalvis, convoluta, valvula
viridi, bipollinari interne membranacea, tenui duplo brevior. Tubus
albidus subcylindricus, longitudine spathe. Calicis limbi lanceolata
pollede octobulata linea utraque media purpurea. Calicis stigma
triquere latifolium

ter. spec. gladiol.
non unicus
+ inter polyp. 7.

non

Carinatus 12 ^{sub} ~~Gladiolus~~ ^{corollae} ~~corollae~~ ^{ringens} ~~antheris~~ ^{sub} tubo spathae limboque brevior, stigmatibus
indivisis complicatis, foliis utriusque carinatis +

Gladiolus carinatus; foliis linearibus utrinque carinatis glabris, tubo
spathae limboque brevior, stigmatibus indivisis complicatis. Hort. Kew

^{1817. 64} Gladiolus punctatus; corolla ringens, foliis linearibus; scapo subquin-
quefloro, floribus erectis distantibus, folio infimo vaginatis puncti-
tato Farqu. collect. 4. p. 160 ⁴ var. 2. 1. 247.

Habitat ad Cap. b. Spei 7.

Scapus ~~caulis~~ sesquipedalis, erectus, teres, superne foliosus. Solum in bulbo
na folio infimo vaginas spathae, ^{scapus} ~~caulam~~ ante vaginatis; Superius
^{scapo} ~~caulis~~ altiora, uti vagina striata, angusta. Flores ^{sub} ~~qui~~ ^{bi-}
unciales. Spatha bivalvis: valvulis lanceolatis, subaequalibus, ^{inter-}
virescentibus.

Corolla tubus cylindricus semipollicaris, cylindricus: lacina limbi
laciniis tres superiores obovato-oblonga; inferiores lanceolata, ^{inter-}
media

^{et sumis} ~~Corolla~~ tubus albidus; limbi laciniis superiores pallide violaceae, inferiorem
infra mediam longitudine superiorum infra medium flavidam;
laterales; laterales breviores, medio flavo Hort. Kew. 6
Solum in bulbo non florente sesquipedale, ^{angustiore} ~~angustis~~ ^{sumis} ~~sumis~~
Vagina infima punctis badius variegata Farqu. 6. u.

Gladiolus ringens; foliis linearibus, caulis, floribus, ^{ringentibus}, corollae sumis and inferioribus. 679

Cisteus gladiolus

+

Habitat

herb. sup.

Caulis pedalis, teres. Colea linearis - ensiformis, caule altero, subtile pime
stnata. Spica ^{lat} ramifera, secundis. Flores approximati. Spathe oblon
ga, acuta, Corolla tubus longitudine spathe. Lacinis unguiculari sa
panem ultra.

lineatus¹³ Gladiolus corollae subnigrescentibus, laciniis ovato-oblongis obtusis dorso tri-
striatis, foliis linearibus lanceolatis +
Fraxinea squalida s. striata; foliis striatis, laminis ovato-oblongis integri-
mis basi concoloribus. Hort. kew. t. p. 61.

Gladiolus lineatus; corollae limbo tubo quadruplo longiore, incurvo patente,
laciniis ellipticis extus linearibus tribus parallelis, suprema majora,
exterioribus rotatis Salob. prov. 40. Cust. may. 4 87.

Habitat in Cap. b. spe. 2

Solia acuminata, striata, plana, vix spithamea. Scapus foliis
duplo longior, teris 8 locis remoti. Spatha bivulvis: valvulae membrana-
reae, apice fibrae vix gemunciales. Corolla pallide lutea,
venis obscuris Hort. kew. l. c.

Scapus tripedalis. Flores quatuor - per Cincas tres extus in medio lacini-
arum Cust. l. c.

Laciniae duae superiores laterales apice emarginatae secundum
figuram Curtisii. Spathae potius similis Fraxinum quam Gladiolus
lunum

Cunonia A. corollae subpapilionaceae: labii lobis duobus externis latioribus adfer-
dentibus - Mel. ec. 113. Lin. Spec. pl. pag. 544

Cunonia floribus sespilibus, spathis maximis Benth. cur. 211.

fig. 1.

Habitat in Persia cap. b. spec.

Folia utroque attenuata Labrum superius obtusum
emarginatum inferius 5 partitum. laciniis tribus intermediis
minutis lateralibus latis Labro superiore demum brevioribus
Spica longa difflua. Spatha lanceolata
glabra striata

Cunonia foliis ex inferioribus utroque ~~attenuatis~~
Superioribus oblongis diffluis Thunb. prod. pag. 7.

an potius e cap. b. spec.

le no 14 Gladialex: corollis secundis cernuis, spathis tubo longioribus, foliis
~~trigulis~~ Gladialex nitida inodora Thunb. Glad. no 8. D.
Gladialex laevis; foliis linearibus bivalve longioribus, corolla curva lutescente, spathis
laevibus tubo longioribus Thunb. prod. 104. Act. Soc. h. n. hann. vol.

Hb. Capus

Caulis pedalis, ^{simplex}, ^{erectus}, ^{apice}
^{flexuosus}.

Habitat in Cap. b. Spei 2

^{Caulis} Folia radicalia duo, inferum ^{longius}, alterum brevius, apice
attenuata, glabra, uti reliqua planta, nitida, Liapor pedalis, ^{simplex},
^{erectus} caulium unicium, ^{in scapo} supra medium, apice valde attenuatum #
Caulis pedalis, ^{simplex}. Flores ^{bini tres} (belluandi cernui, lutescenti albi). Spa-
tha ^{valvula} bivalvis, ^{in aquales} longitudine tubo, convoluta; exterior corollam fere
aequans, inferior longitudine fere tubi, convoluta, obsesse lanico
lata, nervosa. Corollae lacina exterior duplex purpurea; lac-
inibi lacini interior purpureo striatis Thunb. b. c.

brevifolius G.

Glaucolus brevifolius Gaertn. Coll. 4 p. 156. 42. var. 2.

Habitat in Cap. S. Jaci 4

Totus glaber. Caulis pedalis, erectus flexuosus; basi glaber vaginata
solca radicali nulla. caulium quatuor, vaginata, spatula, bre
vissima acuta, cauli adpressi, spatula formis. Spica terminalis
simplex, septemflora, diffusa erecta, secunda. Spatha bivalves
lanceolata, striata erecta, subaequalis exteriori fuscescenti interiori
viridis. Corolla petala, rubra incurvata. Tubus gracilis albus limbi
calicis superiores latissime lanceolata obtusula: inferiores angustiores
breviores. Gaertn.

puberulis tubo & pappis longiore,
tristis. Gladiolus corollae campanulatae laciniis lanceolatis ovato-lanceolatis
planis, foliis linearibus bisulcatis

unicolor. Gladiolus foliis sulcatis linearibus; floribus alternis; petalis acutiori-
bus Mill. Dist. ed. 157. t. 235. f. 1.

Gladiolus tristis; foliis linearibus - cruciatis, corollis campanulatis

versicolor
versicolor
Lini 8 pec. pl. 53. Jacq. collect. 4. p. 157. f. var. t. 243
β. Gladiolus, floribus Gladiolus; floribus ex radice et flavo variega-
tis, suprema laciniis brevissimis Breyn. ic. var. 20. t. 7. f. 5

Gladiolus et bifolius et biflorus; foliis: quod ranguli Sw. Eret.
t. 39

Gladiolus tristis Jacq. Suppl. 20. f. var. t. 245.

Gladiolus tristis Purt. mag. 8. t. 272.

Habitat in Cap. 6. 8 pec.

sp. 19. Gladiolus versicolor. foliis linearibus cruciatis; floribus maximis, versicoloribus Sw. t. 19

Habitat in Cap. 6. 8 pec. 24

a longe vaginantis, unicum, subulatum,
folium radicale & caelo altius, totum, angulis quatuor
exstantibus planis cruciatis, reliqua breviora: vaginae & viciae.
scapus varicosus, erectus, superne hirtuosus. Flo. interdum unicus,

paucis
4ae pe plures, multae, odoratae simus. Spatha trivalvis: valvulae lanceolatae, concavae,
marginis tenui membranaceo; exteriori longitudine fere tubi. Corollae lobis lanceola-
tae, suprema latior erecta, reliquae angustiores paucis reflexae.
In multiflora folia rite ensiformia evadunt. L. C.

albidus 16 Gladiolus ^{tribus} corollae subingentis laciniis lanceolatis, ^{superioribus} undu-
latis, ^{tribus} inferioribus apice conniventibus bimaiculatis. †

gladiolus albidus; corolla subingente; laciniis acutis, ^{tribus} superiori-
bus undulatis, ^{tribus} inferioribus apice conniventibus, foliis ensiformibus
subobliquis Jacq. collect. suppl. 21. t. v. 256.

Habitat ad Cap. b. Spei 4

^{superis}
Bulbus subrotundus, magnitudine avellanae. ^{caulis} viridialis, crassi-
tie pennae columbinae, ^{erectus} flexuosus. folia plura, infimum
semipedale, superiora sensim breviora, erecta, firma, striatula
spatha bi-dores tres-quadro, ^{trispallione} ~~altero~~, inodori, triuncialis
spatha bivalvis; sesuncialis: valvulae lanceolatae, aluminatae,
convolutae; interior brevior. Corolla ^{alba} ~~alba~~ ^{tribus} longitudine pen-
spathae, subcylindricis, incurvis, albis; limbus campanulatus
laciniis lanceolatis, ^{laciniis} acutis; ^{superiores} tres ~~superiores~~ longiores spatha
duplo longiores, undulatae, inferioribus apice conniventibus et
perperascentibus, supra medium maculas duas roseas Jacq.

l. c.
an gladiolus bimaiculatus Lamour. det. bot foliis perangustis,
petalis superioribus brevioribus patenti, reflexis: trium inferum
lateralibus maculatis angustioribus
an huc pertinet bimaiculata Lamour.
superioribus ~~superioribus~~ duae laciniis modo in figura macul.

Tubulofus 9.

Gladulus tubulofus Jacq. coll. 4. p. 153 in. rar

Habitat in Cap. b. p. 2

Planta elegans. Aliter referens. Tota glabra. Folia radicalia 3 vel 4. exi
formi-linearia, integerrima, obliqua, late sesquipedalia. Scapus latera
lis, teres foliis longior, erectus, parum flexuosus, foliis paucis vaginan
tibis instructus, saepe saepe spica. Cava quinquiflora secundaria. Flores
cornu bicornei. Pertha trivalvis valvulis foveis purpurea
dentibus acutis extenore ovata internodiis linearibus lanceolatis corolla
quadriplo brevioribus. Corolla saturate corinea tubo angusto incurvo
laciniis ovatis aequalibus. Carina stigmatis bifida. Saigu

in 20
wid.
non unv.

tubul p

non vidi

Cuspidatus 17 Gladiolus corollis subnigentibus secundis carnis, laciniis lanceolatis
undulatis cuspidatis, tribus inferioribus macula ocellata.

Gladiolus cuspidatus; corolla subnigente; laciniis lanceolato-cuspidatis,
undulatis; foliis linearisensiformibus, glabris; floribus secundis, carnis

Jacq. collect. suppl. 17. Fr. rar. t. 257. Redouté lilie t. 36

Habitat ad Cap. b. spec. 4

Scapus

Bulbus subrotundus, avellana multo major. Caulis sesquipedalis
crassitie plurae columbinae, ~~tristis~~ ^{strigulosus}, flexuosus. Folia radicalia

4 Longitudine tria, duo caulina brevia, Flores secundi, plerumque, quatuor, & esse
fere caulis, ter diutantes, quodnunciales, patulae. Spatha bivolvris, binucialis
longe vaginam
si et ultra, convoluta; exterior integerrima, interior brevior, bi
dentata. Corollae tubus sensim ampliat, incurvulus, extus sordid
is sine pallens: limbi lacinae ^{laciniae} lanceolatae, cuspidatae, magis
vel minus undulatae, saepe obliquae, patentes sinuae, tres super
iores, duplo longiores latiores, parum longiores, utrinque
sordidissime et pallide carnea, de in medio dorsi purpurascen
tes, inferiores stramineae, basi macula purpureo rubra ocellata.
Jacq. l. c.

punctatus 9

Gladolus punctatus Gaey. coll. 4. p. 160 et var.

Habitat ad Cay. b. p. 1. v.

+
Solum in bulbo non florente linearis sesquipedale, angustissimum, in plan-
ta florente alterna, basi vaginaria, striata, lanceolato-linearis, plana
vix sesquipedalia. Tolum inferum spatuliforme maculis punctis baduis
variegatum. Scapus sesquipedalis teres erectus. Flores racemosi, alterni
distantes, biantheri, sex quini. Spatha bivalvis: valvulis lanceolatis
subaequalibus, venatis. Corolla tubus cylindricus. ~~Corolla~~ Spatha con-
grua, compressa. Lacinia limbi lanceolata, superioris tres latiores
inferiores patentiore parte media flava Gaey.

et varietas *G. punctatus*

trimaculatus 10 Gladiolus corolla carnea tubo & pathis longiora, ~~tres~~ laciniis tribus

inferioribus macula cordata purpurea.

Gladiolus floribus patentibus externe carnea, interne candidis, & cella
mafulisque purpureis insignitis, suprema lacina proxima Bryn.
rar.

Gladiolus trimaculatus; foliis linear-lanceolatis, tubo curvo limbo
vix longiore, petalis tribus macula cordiformi inscriptis Lamarck

Diat. bot. ? p. 927

Gladiolus cordatus, monostachyus foliis ensiformibus nervosis,
corolla curva incarnata: macula cordata, & pathis longiora tubo
brevioribus Thunb. prodr. 105. act. Soc. h. n. hafn.

Habitat in Cap. b. Spei &

Totus glaber. Folia plura, usque ad quadruplum, infimum scapo longius, reliqua pen-
sum breviora, apice attenuata, marginata costis tribus, quarum media
major, glabrescentibus. Scapus simplex, & rectus pedalis, ex

apice flexuosus, glaber. Flores duo - quinque, ^{bipolliares} campanulati, carnei.
Pathis oblonga subpolliares, oblonga, acuta, nervosa, tubo pau-
lo breviora. Corolla albida incarnata: limbi ~~tri~~ ^{trid} tubo ~~curvato~~:

limbi laciniis & equipollentibus, curvatis: limbi laciniis infimo macula
cordata purpurea. Thunb. l. c.

Gladiolus cordatus

14 Aug. 27

#

~~prodr. 105~~

Tenellus Gladolus

gladulus tenellus Gay. coll. 3. p. 255. 4200

Habitat ad Cap. b. p. ei

Folium rad. caule unicem linearis basi vagrant acutum,
consolatum erectiusculum longitudine caulis. Duo caulium
alterna inferne vagrantia familia radice scapula
v. rami. Caulis sempetalis, gracili debili subflexuoso ad
folia geniculatus, supra folium summum dilatatur, biflo-
rus. Flores perianthali. Spatha diphyllo folioli linear-lan-
ceolatis, acutis concavis, erectis, subaequalibus. Corolla octo
lucida subnigra linea longitudinali sordida ~~non~~ violacea
lacini duo patula ad faucem punctata Gay.

non radi.

hastatus 19 Gladiolus ^{campanulato} corolla tubo spathae breviora, laciniis ovatis obtusis.
tribus inferioribus macula hastata +

gladiolus tristis ^{hastatus} Thunb. glad. n. corollae albo-incarnatae macula hastata Thunb. glad. n. 9. 0

Gladiolus angustus ^{tristis} aequ. collect. 4. p. 154 G. rar.

Gladiolus hastatus monostachyus foliis linearibus bisulcatis, corolla campanulata: macula hastata, spathae brevibus tubo longioribus. Thunb. prodr. 105. act. bot. hort. n. hafn.

Habitat in Cap. 6. spec.

Folia radicalia duo, glabra scapo paulo breviora, basi latiora striis pluribus, apice linearibus bisulcatis, caulinum versus apicem simile, bipollicare. Scapus ^{pedalis?} ~~simplex~~ erectus, apice flexuosus. glaber. Strobili cincti tres. Spathae convolutae, longitudine dimidia corollae, ovatae, obtusae, laevae nervosae. Corolla albo-incarnata: limbi lacinae ovatae, inferiores macula hastata.

recurvus 20 Gladolus corollae laciniis lanceolatis planae reflexae recurvatae

+
Gladolus foliis linearibus sulcatis, florum uno versu dispositis, tubo floris
longiore Mill. Fl. ic. 157. t. 235. fig. 2.

Gladolus recurvus; foliis ensiformibus, petalis subaequalibus lanceolatis
recurvatis Lin. mant. 20.

Habitat in Cap. b. spec. 4

Differt ab gl. undulato petalis non undulatis minusque acuminatis;
tubo corollae non filiformi, lato aut pendulo; ceterum valde affinis
lin. l. c.

Scapus testis fere virginatus, palmatis, curvatus. Folia pro caps breviora,
linearibus. ~~Floris~~ Floris quatuor, duos rubri, inter albi
ex Thunbergii eadem ac Linnaei? secundum Thunbergii nimis affinis
est gl. falcato, secundum Linnaeum vero affinis gl. undulato, qui
multum distat a falcato.

Gladolus recurvus

~~##~~
grandis 16 ~~Gladiolus cordatae limbis~~ ~~cap. per~~ ~~litho~~, ~~cap.~~ ~~cap.~~ ~~cap.~~ ~~cap.~~
~~longitudinis~~ ~~cap.~~

~~##~~
~~gladiolus~~ ~~grandis~~; ~~foliis ensiformi~~ - ~~attenuato~~ ~~multistriato~~,
~~corolla curva purpureo~~ - ~~strata~~, ~~tubo simplici amplia~~
~~to~~ ~~Thunb. prod. 105. Act. Soc. h. n. havn. vol. 6.~~

Gladiolus versicolor, ~~habitata in Cap. b. spei v~~
foliis linearibus ~~crucis~~

~~duo~~; ~~floribus~~ ~~ma~~ ~~tubum~~ ~~radicale~~ ~~unicum~~, ~~longitudine~~ ~~blapi~~, ~~apice~~ ~~valde~~ ~~athe~~
~~stimo~~, ~~versicolori~~ ~~nudum~~, ~~erectum~~, ~~multinervosum~~; ~~et~~ ~~tota~~ ~~planta~~ ~~glabrata~~;
~~bus~~ ~~Andr. repos. t. 19?~~ ~~cuculino~~ ~~sub~~ ~~floribus~~ ~~bractei~~ ~~formia~~, ~~corollam~~ ~~sub~~ ~~qua~~
~~tra~~, ~~crista~~ ~~duo~~, ~~obsera~~, ~~convoluta~~, ~~ensiformia~~ - ~~scapo~~

grandis 21 ~~Gladiolus~~ ~~pedalis~~ ~~vel~~ ~~ultra~~, ~~simplex~~, ~~flexuosus~~ ~~caetera~~, ~~terre~~. ~~flori~~
~~bus~~, ~~cernui~~, ~~magni~~. ~~spatha~~ ~~intra~~ ~~bracteam~~, ~~similis~~, ~~sed~~
~~brevis~~. ~~corolla~~ ~~tubus~~ ~~curvatus~~, ~~sensim~~ ~~amplatus~~ ~~in~~ ~~tubum~~
~~campanulatum~~, ~~albida~~ ~~flavescentem~~: ~~limbi~~ ~~laciniis~~
~~tres~~ ~~superiores~~ ~~intus~~ ~~linea~~ ~~duplex~~ ~~purpurea~~, ~~extus~~ ~~purpureo~~
~~multistriatis~~; ~~inferiores~~ ~~intus~~ ~~bulbo~~ ~~medio~~ ~~flavescente~~, ~~apice~~
~~et~~ ~~purpureo~~ ~~cente~~. ~~Thunb. l. c.~~

+ 15

~~Andr. Glad. versicolor~~ ~~Andr. videtur esse varietas~~
~~hujus~~

trimaiculatus Glad. foliis linearibus canaliculatis, tubo limbo limbo
Cingulo petalis tubis macula cordiformi impressis la
Mark J. C. pag. 116 - Tab. 32. fig. 3.
Habitat in C. b. p.
eadem videtur ac angustus Gay.

2000

inflatus 22 *Glaucolus* ^{in flato-} corollae campanulatae tubo & spathis ovatis ^{nervosis} aequante, foliis
^{tristis +}
Glaucolus tristis violaceus Thunb. *glad. n. d. n.*
Glaucolus inflatus; folio lineari ^{tristis}, corolla campanulata violacea,
Spathis ovatis nervosis tubum aequantibus Thunb. prodr. 185. Act. Soc. Ich.
n. havn. vol. 6.

Habitat in Cap. b. & pai

Solia radicalia duo, inferum & cap. longius; alterum brevius,
base latior, parum attenuata, apice linearia, multistriata; utri-
toto planta globa; Stapi & equipollicare. Flores bini vel tres qua-
tuor, & equipollicares, subsecundi curvi, inflato-campanulati,
inornati vel purpurei: tubus brevi. Spatha longitudine tubi,
valde lato, ovato, convolato, acuto, nervoso. Thunb. l. c.

~~Venellus~~

~~13 "Glaber" Venellus Gagn. coll. 4. p. 169~~

~~Habitat in Cap. b. p. p.~~

~~Torus glaber. Folia supra breviter sequentia lanceolato-linearia~~

~~acuta convoluta. Anthera longitudine scapi, testis, semipedalis
flexuosi, biflori. Intra diphylla floris dimidio brevior: folioli
lanceolatis acuti. Ovulo flavo. Carina lanceolata saturate
luteo~~

+
venellus
+ tenellus

non vid.

hyalinus 23 Gladiolus corollae subringentis laciniis lanceolatis, foliis vaginis
que glabris
Gladiolus strictus Jacq. collect. 4. p. 170.

Gladiolus hyalinus; corolla subringenta; foliis caulinis aeternis,
ensiformi-linearibus, glabris, caule multo brevior; caule et floribus
erectis Jacq. ic. rar. 2. t. 242.

Habitat in Cop. b. p. 24

Scapus pedalis, dactylis, triphyllus. Folia tria in firmum radicale,
stricta, subtrinaria. Flores circiter tres. Spathae lineari-lanceola-
tae, obtusiuscula, viridae: valvula exterior semunialis: interior
brevior, apice bifida. Corolla binialis: tubus cylindricus, cae-
uniculis, coeruleis; fauce pallida, cum maculis bord de purpu-
rascentibus: limbi laciniis lanceolatis, acutis, subpallidissae;
supremo triplo latiore albida, superne flavescente, cum striis
punctisque purpurascens, reliquo concolora et borda flavescens:
bas, cum linea perpendiculari media, striis punctisque pur-
puris Jacq. l. c.

germanica 4 foliis ensiformibus glabri falcatis brevioribus scapo multifloro
Thurb. Fr. n. 9.

4 ni vulgaris germanica s. sylvestris Bauh. pin 30

4 ni germanica corollis barbatis, caule foliis longiore multifloro,
floribus inferioribus pedunculatis Lin Spei: pl. pag. 554

4 ni germanica
Thurb. hist.
3107

Habitat in Capto Germania edisi

scapus inferne simplex, multiflorus, foliis longior, bipedales
Folia reflexo-falcata, nervosa, pollicem lata. Flores coerulei
petala minima integerrima. Thurb

4 ni vulgaris
violacei Bauh
hist 709
yros cuba
math di
quid in flore albo
Bauh. 719
Besler.

~~Caulis fistulosus. petala majora extus coerulea venis saturatioribus
caulis
bui medio tubuli albidis emargina extus inferne flavescientia venis
violacei dorso virentibus basi reflexis, majora extus, lanceolata
cuneata parva saturatiora venis violaceis basi convergenti
concolorati~~

~~herbena dependens~~

~~Pseudacorus~~ ~~foliis ensiformibus, petalis alternis stigmate minoribus Thunb~~

~~fr. n. 10.~~

~~ac. in falsum Cam. ep. 6.~~

~~Acorus adullerinus Baub. pin. 34. Theatr. 634.~~

gladiolus
Cabe

~~In pseudacorus corollis imberibus. petalis ^{interioribus} longioribus stigma
te minoribus; foliis ensiformibus Lin. Spec. pl. pag. 567~~

math.

~~Habitat in Europa ad ripas paludum, Persiarum &~~

~~Scapus tripedalis multiflorus. Folia ^{cubitalia} radicalia et caulin-
alternantia, plana, pollicem lata, scapum aequantia
Flores pedunculati alterni, plures. Spatha lanceolata
tripollicaris, pollicem lata petala omnia flava~~

~~acorus officinarum Tuct 13~~

~~math.~~

~~acorus Brasero fer~~

~~E. Chamoini latifolia minor variegata Plin. pan. 2. 25. Chamoini
latifol.~~

~~D. Chamoini latif. minor pallida et alba Pan 225~~

~~Chamoini lat. m. 4. Plin. hist 225 Ey. O. 1.~~

~~mun. hist. 5~~

Striatus Gladulus corollis ringentibus secundis, laciniis alternis crispis, foliis lanceolatis ensiformibus undulatis ciliatis +

Gladulus Striatus; corolla divaricata, ringente; laciniis tribus alternis crispis; foliis lanceolato-ensiformibus, crispis, glabris, ciliatis Laegu
collect. Suppl. 28. 4c. rar. 2. t. 260

Habitat in Cap. b. Spei 7

Bulbus subrotundus, avellanae magnitudinis. Scopus foliosus, pedalis, inferne aphyllus, atroviolaceus. folia semipedalia vel minor, glauca. Spica subperiflora, interdum horizontalis. Corolla 2 esqui-pollicaris. Spatha diphylla, superne scariosa, germine duplo longior; valvulae subovatae; acuminatae anterior acuminata, interior bifida, Corollae tubus angustus, spatha duplo longior, pallide et bordide visens. Limbus profunde fissus; laciniis alternis acutiusculis, sat flavescens, tribus mucronatis, pallide violaceis cum fascia longitudinali media rubra, plana Laegu. l. c.

galeatus 26 *Gladiolus* corollae laciniis nigris laciniis tres inferiores ~~et~~
~~spathe latae erectae spathe latae~~, foliis ensiformibus obtusis
Synnichium viperatum ^{alm} Plukm. phyt. 1748. t. 24. f. 10

Gladiolus flore singulari ^{minuto}, inferioribus laciniis luteo, vindi-
bus Weyn. ic. rar. 25 t. 12. f. 2. Seb. Thes. 1. t. 7. f. 3.

Gladiolus alatus; foliis ensiformibus, petalis lateralibus latis, si-
mis ^{amoen} Lin. Amoen. ac. 6. p. 83.

~~*gladiolus* alatus, Thyrid. Diss. n. 15. var. 2~~
Gladiolus galeatus; corollae divaricate nigrae, laciniis summa
erecta, foliis ensiformibus glabris Faigu. coll. 4. p. 167. Ic. rar. 2. 7.

250.

Habitat in Cap. b. 8. p. 24

~~Totus glaber. Caulis erectus, spathe ^{compressus} ~~mois~~, erectus, panem ~~flexuosus~~, ~~caulina~~ folio-
rum et spathe ~~decurvante~~. Solia ~~una~~ ^{compressa} ~~vel~~ quatuor in
caule eoque breviora, angusta. Flores~~

Totus glaber. Caulis ^{semipedalis} ~~semipedalis~~, flexuosus, erectus, ~~terres~~. Solia
radicalia plura, erecta caule breviora. Flores plures, distantes, panem
nutantes, inodori. Spathe ~~scario~~ ^{scario} bivalves; longitudine faulibus co-
rolle: valvula acuminata, ut plurimum bifida. Corolla est ~~violacea~~
-verte

borderi ^{de} pallet: Limbi lacinia superior magis violacea, saepe ad latera flavens; tres
inferiores interne circa medium aut ex integro aut latitum ad latera extremum
saturate flavescent. Gaesq. l. c.

^{lociniis & spatulatis}
^{secundis}
 Communis 26 *Gladiolus corollis subargentibus, & pathis tubo longioribus, foliis*
^{ensiformibus nervosis}
Hesperion, gladiolus & patha seu gladiolus vegetalis Ruell. de nat. Hist.

781

Gladiolus Dod. ~~causa~~ pempt. 209. *Fiv. monop.*

Gladiolus narbonensis flore incarnato et flore purpureo Best.
 hort. cyst. Ost. ord. 4 fol. 16

Gladiolus communis, foliis ensiformibus distantibus Lin.
 Syst. veg. 87 Roth. fl. germ.

B. *Gladiolus utrinque floriger* italicus binis florum or
 dinibus cinctum. Tab. 2. 99

Gladiolus utrinque floriger Dod. pempt. 209

Gladiolus italicus flore rubro Best. hort. cyst. Ost. Ord. 4. fol. 12

Habitat in Europa Australi et Africa boreali &

folia & scapo multo breviora, pollicem fere lata. Scapus bipedalis
 erectus, teres & simplicissimus, ultra medium foliis vaginatus. Spatha
 lanceolatae, ~~bivalves~~ ^{flexuosa}. Spathae trivalves; Calvula
 lanceolatae; exterior pollicaris & ultra, atque tubo longior, attenua
 ta, interior brevior, longitudine tubi

Index Auctorum

Act. ac. nat. cur.

Acta physico-medica Academiae Caesareae Naturae
Curiosorum Norimb. 1727-1754. 4to

Act. ac. mogunt.

Acta Academiae Electoralis Moguntinae Scientiarum et
Artium, quae Erfordiae est. Erfordiae 1757 et sequ. Tom. 1. 8vo reliqui
in 4to.

Acta ac. neap.

Atti della Reale Accademia delle Scienze e Belle-Lettere
di Napoli, dalla fondazione sino all'anno 1787. Napoli
1788. 4to

Act. acad. Berol.

Histoire de l'Académie Royal des Sciences et Belles Lettres
de Berlin. Berlin 1746. et sequ. 4to

Act. bonon.

De Bononiensi Scientiarum et Artium Instituto atque
Academia Commentarii Bonon. 1748 et sequ. 4to

Act. Hafn.

Skrifter som udi det Kiøbenhavnske Selskab af Lørdagene
og Videnskabers Elskers ere fremlagte og op læste
Kiøbenhavn 1748 et sequ. 4to

28
carneus 27

^{carneus}
Gladiolus corollis pubescentibus distichis, laciniis oblongis,
foliis ensiformibus marginatis

gladiolus carneus; caule simplicissimo, petalis superioribus divergen-
tibus; foliis ensiformibus La Roche Desp. n. 3. t. 4.

gladiolus carneus; corolla pubescente, foliis ensiformibus striatis
glabris, capro flexuoso erecto, floribus carnis Gay. collect. v. p

370

gladiolus ventricosus; foliis ensiformibus nervosis glabris, limbo
ventricoso difformi, styli laciniis dilatato membranaceis & patulatis

Lamarck. illustr. Dict. bot. 2. p. 727. illustr. 1. p. 118

4 quadri-ortoflorus

Habitat ad Cap. b. & p. Ex herbario Lamarckii et Royeni

^{capis}
totus glaber. Caulis pedalis, ~~semplax~~ flexuosus, ~~erectus~~. Folia
radicalia semipedalia, duo vel tres in caule, erecta, ad rige

4 incarnati, tripollis
caesi, vel parum
ultra

dula, acuta, ~~florata~~, ^{marginis} nerva crassior ⁱⁿ nervo crassiore margi
nata. Flores ^{sesquialter} alterni, distichi, distans ² ~~tripathis~~ ^{tripathis} bivalvis, extenor

30 sic bipollis canis, interior longior, lanceolata, attenuata, viridis.
Corolla tubus & pathis vix longior, curvatus: limbi laciniis lanceolatis;

25

verte

lacina superior latior. Stigmata extrorsum dilatata

29
equitansi 20 Gladiolus corolla campanulatae tubo brevissimo, laciniis ^{erectis} oblonga ovatis
— Stapo compresso, foliis ellipticis ensiformibus multinerviis marginatis
Gladiolus equitansi; foliis ensiformibus marginatis multinerviis, stapo
compresso, corolla campanulata tubo brevissimo Thunb. prodr. Act.
Bot. h. n. hafn. vol. 6 t. cum figura

Habitat in Cap. b. spci 4

folia circiter quatuor, stapo breviora; equitanti in femore ^{un} plusquam
pollicem latam, subfalcatum, patens, equitanti multinerviis; nervis
et margine crasso flavescenti. Scapus pedalis, simplex, erectus, flexuosus;
graber, uti tota planta. Flores purpurei. Spatha corolla longior es,
oblonga, acuta, convoluta, pallida, margine et carina purpureo.
Corolla tubi brevissimi. Similis glad. alato, sed longe major et
simplex Thunb. l. c.

Corolla Aspinia
oblonga, ovatis

Dignoscitur a reliquis foliis latissimis:

#13 in di folius 29 Gladiolus corollis campanulatis tubo spathae longiora, laciniis oblongo
^{subtrianguliformi oblongis, capsulis} ^{foliis ensiformibus}

Gladiolus maxi-
mus promont. b.
Spici flore rubro
incarnato Swert.
flor. t. 66
Corollis carnis
faciq. ii. rar. 2.
t. 235

Gladiolus pyramidalis; spica pyramidalis longissima, basi laxa sub
ramosa, floribus amplis liliaceis natis; stylis tripartito-bifi-
dis Lamark. dict. bot. 2. p. 726. Illustr. 1. p. 110

Gladiolus in di folius; corollis infundibuliformibus, limbi laciniis equa-
libus, foliis ensiformibus distichis rectis faciq. collect. 4 p. 151.
fac. rar. 2. t. 234

#14 Habitat in Cap. b. spici. Ex herbario Lamarkii 2
vel minor, inaequalia
Tubus radicalis, circiter septem bipedalis, pollicem ^{max} lata, utrinque abe-
nuata Scapus tripedalis, crassitie penna cygnes, vaginis non nullis lan-
ceolatis vestitus, erectus, teres, lavis. Spica sesquipedalis, multiflora,
parum flexuosa. Spatha bisulca; valvulae semipollicares, lanceolae
squales, semis lanceolatae, adnatae, superne flavescentes cum
rubescine. Corolla fere tripollicaris, pallide carnea: tubus longitudine spathae
fere flavescente: limbi laciniis subtrianguliformi, alternis mucronatis

4 disticha

an gladiolus pyramidalis Burm. differt ab ante cedente: 3 Capis altior
foliis ^{angustis} longioribus, angustioribus, minime nervosa spiculis minime raris
nervosa: spathae breves multo angustiores: floribus plurimis

glabrus gladiolus polytachyus, corolla laciniis lateralibus latissimis,
caulis foliis linearibus ensiformibus basi compressis
Thunb. Desf. par. 16. n. 18. Jacq. coll. 4. p. 165 Schum. ii: pl. rar. 6. tab. 12
75. v. v. gladiolus flore singulari mucato, inferioribus laciniis luteo
viridibus B. Reyn. var. p. 25. t. 12. fig. 2.

Habitat in cap. b. p. i.
Scapus flexuosus erectus striatus simplex vel ramosus, prolon-
gatus usque pedali. Folia marginata foliaceo a basi spatham
decurrenda compresso subulata. Folia alba vel rubra vixantia
ensiformia striata glabra superioribus sensim brevioribus
rachis flexuosa, toxa. Flori alterni vel secundi. Spatha val-
vula exterior cancellata compressa tubo duplo longior obli-
qua canna et margine purpureo. Spatha interior breviori tubo
corolla longiori variis floribus incarnatis striatis et violaceis

Cardinalis 31

Gladiolus Corolla ⁸ campanulatis secundis, tubo spathae brevior, laciniis lanceolatis, foliis ensiformibus marginatis

gladiolus cardinalis; corolla erecta limbo campanulato, floribus secundis, scapo multifloro, foliis ensiformibus multinerviis Cust. Mag. 4. t. 135

gladiolus cardinalis; foliis ensiformibus; caule multifloro, floribus secundis, corolla laciniis inferioribus macula alba rhomboida distinctis Voh. ic. pl. rar. t. 27.

Habitat in Cap. h. spec. 4.

Caulis bi-tripedalis, erectus

Folia unquam vel ultra lata, scapo breviora, patentia, laeva, margine parum incrassata, striato nervoso-striata; ^{scapi} ~~laeva~~ superne, vaginantia brevia, tri-quadrupollicaria; vasa glauca adspersa. Scapus bi-tripedalis

erectus, teres, vasa glauca adspersa. Spica solitaria vel gemina, parum recurva, sex-septemflora. Flores secundis, sancti, erecto-patentes.

Spatha bivalvis: valvula exterior sesquipollicaris, lanceolata, attenuata, concava, extus convexa, vasa glaucescente adspersa; interior ab exteriora inclusa, brevior, angustior, apice bifida. Corolla tubo spathae longior, sursum sensim latior coliciner. limbi laciniis superior

apice reflexo-patens
maior et longior quam reliquis, erecta, oblongo-ovata, obtusa; duo laterales, lato-lanceolata,
reflexa, coccinea, ut superior immaculata; tres inferioribus lateralibus micantibus
angustioribus, planioribus, lanceolata, obtusa, macula laevi coccinea, a medio ad basin
albida in lineam albidam tenuissimam desinens. filamenta coccinea. Stigmata
compressa - clavata, coccinea, albido-villosa

Planta diversa a *gl. indifolia*

32
roseo, albus. ³² St. gladiolus corollis infundibuliformibus, limbi laciniis ellipticis aequa-

^{lex, foliis linearis-ensiformibus}
gladiolus roseo albus; corollis infundibuliformibus; limbi laciniis aequalibus; fauce tereti; foliis anguste ensiformibus, obliquis; spatha valvula interiori bifida Gaerth. hort. schwenb. 1. p. 7. t. 13.

Habitat ad Cap. b. spei. 7

Radix tunicata ^{foliis} ~~caulis simplex~~ Scapus foliis brevibus vestitus. Folia anguste ensiformia, pedali
scapo foliis brevibus vestitus, longitudine foliorum. Flores distincti sessiles in
catervis. Spathae lanceolatae, fuscas acuminatae, fuscas: valvula inte-
rior bifida. Periclyae tubus calycis parum longior: limbus patentissimus. Lacini
inae ellipticae, aequales.

bicolor *Polypodium polytrichum*, corolla nigente, spathe lacero aristatis, foliis copiformi-
bus glabris Thunb. Disf. p. 16.

habbris *Habitat in collibus ad Cap. b. p. 4*

reticulatus
ovatus

Caput angulatus fructus, foliis utraque profundo excavatus, basi foliis vaginatus, gla-
ber, flexuosus erectus, spithamou. Folia alternatim, vaginantis, obversa, obtusa
cum acumine, pinnato, capitulo breviora. Specie duo; altera uniflora, tenuior,

7. 21

+

3/4 sequi

Collis 1. 5

4 c. var. 2

altera triflora crassior, longitudine dimidia fere. Rachis angulata, flexuosa
glabra. Spatha membranacea, apice lacero-partita, laciniis acutis, venosa
basi grisea, apice ferruginea. Corolla tubus si bifloris, sensim appressatus
superne inflatus, spathe 2plo longior flavus: limbilaciniis flavo: suprema
major, ovata concava, apice conulea: laterales inferiores lanceolata un-
voluto, ultimo apice conules: infima brevior parum a reliquis separata, ex-
tra laterales collocata: omnes basi linea duplici purpurea in ore tubi
affinis 7. 21 bulbifera Thunb. l. c.

Watsonius 3337

lanceolatis acutis

Gladidolus corollis subcampanulatis tubo cernuis, fo laciniis oblongis,
spathis acuminatis foliis linearibus ensiformibus marginatis
Antholyza revoluta; staminibus adscendentibus, corollis revolutis fo
liis lanceolatis nervosis Burm. prodr.

Gladidolus Watsonius; corollis cernuis, tubo duplii, limbi laciniis oblon-
gis, foliis linearibus glabris Thunb. Glad. n. 10 Curt. magaz. 450

Gladidolus Watsonius; corollis infundibuliformibus; limbi laciniis sub-
aequalibus linearibus margine revolutis Jacq. Collect. 2. p. 257. is
var. 2. t. 233.

Gladidolus procox; foliis ensiformibus, apicibus tortis, linearibus cras-
satis; corollis subcampanulatis, rubris; petalis acuminatis Andr. repos.

t. 30

B. corollis majoribus variegatis Curt. magaz. t. 569.

Habitat ad latera montium Cap. t. spec. 2

Solia tria, scapo breviori, longe vaginantis, angustis pinnis, stenuato,
trinervis cum nervis marginalibus, utrinque bifurcata. Scapo peda-
lis et ultra, erectus, teres. Flores tres, quinque subsessili, purpurei

Corolla tubus persequipollianis, geniculatus curvatus Thunb. l.
Variat 3, petalis brevioribus et longioribus convolutis acas longe acumina-
tis

N. p. 105

Handwritten notes at the bottom, including 'Hand 104' and other illegible text.

albus Sch. culmo subtriangulo foliis, floribus fasciculatis, foliis glaucis Len. S. V.

Fr. Dan. Tab. 320.

Cyperus palustris hirsutus minor, paniculis albis Mon. hist. 3. J. 1749
fig. 39.

Gramen lucule accedens glabrum, in paludibus proveniens, panicula
lata. Pluk. alm. 170. t. 34. fig. 11.

Gramen cyperoides palustre Cuius antherum Scheuchz. gran. 503.

Habitat in Europa borealis paludibus hirsutus 7

Culmi plures ascendentes, ere vel erecti. Spiculae. folia angustis
pina 2-3 in culmo inferiori breviora superiora altitudine fere cul-
mi pedunculati et ala suprema et apice. fasciculi florum duo
in culmo alter ex stella suprema alter ex apice spiculis 4
vel 5 que subsessiles breviter sub fasciculo inferum longior interdu-
m fasciculi longitudine spicula pallide nivescentes lanceolata
squama lanceolata

spicula oblate flavo femineae.

laciniis ^{ternis} ~~ternis~~ ^{saucis} ~~saucis~~ ^{veretibus} ~~veretibus~~ ^{obtusis} ~~obtusis~~ ⁴ ~~4~~
 laciniis ^{oblongis} ~~oblongis~~ ^{saucis} ~~saucis~~ ^{foliis} ~~foliis~~
 H ^{spathis} ~~spathis~~ ^{lanis} ~~lanis ^{ensiformibus} ~~ensiformibus~~ ^{spathis} ~~spathis~~ ^{acuminatis} ~~acuminatis~~ ^{foliis} ~~foliis~~ ^{ensiformibus} ~~ensiformibus~~ +
 laciniis ^{attenuatis} ~~attenuatis~~ ^{gladiolus} ~~gladiolus~~ ^{laciniis} ~~laciniis~~ ^{corollis} ~~corollis~~ ^{infundibuliformibus} ~~infundibuliformibus~~ ^{limbi} ~~limbi~~ ^{laciniis} ~~laciniis~~ ^{aequalibus} ~~aequalibus~~ ^{saucis} ~~saucis~~ ^{veretibus} ~~veretibus~~ ^{foliis} ~~foliis~~ ^{anguste} ~~anguste~~ ^{ensiformibus} ~~ensiformibus~~ ^{obliquis} ~~obliquis ^{saucis} ~~saucis~~ ^{collet.} ~~collet. ^{4.} ~~4.
 p. 171. f. rar. t. 2. t. 232.~~~~~~~~

Habitat ad Cap. b. Spei 7

~~Totus glaber.~~ ^{Solia} ~~plura~~ ^{plura} ^{angusta} ~~angusta~~ ^{pedalia} ~~pedalia~~ ^{angusta} ~~angusta~~ ^{erectiuscula} ~~erectiuscula~~ ^{nervo} ~~nervo~~ ^{medio} ~~medio ^{utrinque} ~~utrinque ^{protuberante} ~~protuberante. ^{Scapus} ~~scapus~~ ^{foliis} ~~foliis~~ ^{paucis} ~~paucis ^{altior} ~~altior ^{foliorum} ~~foliorum ^{vagina} ~~vagina~~ ^{vestitus} ~~vestitus. ^{Stores} ~~stores~~ ^{cinctor} ~~cinctor ^{quatuor} ~~quatuor ^{sessiles} ~~sessiles~~ ^{distantes} ~~distantes. ^{Spatha} ~~spatha~~ ^{diphyl-} ~~diphyl-
^{la} ~~la~~ ^{valvulis} ~~valvulis ^{anguste} ~~anguste ^{lanceolatis} ~~lanceolatis ^{acuminatis} ~~acuminatis~~ ^{basi} ~~basi~~ ^{veretibus} ~~veretibus~~ ^{superna} ~~superna~~ ^{sphaeclatis} ~~sphaeclatis~~ ^{et} ~~et ^{fuscescentibus} ~~fuscescentibus~~. ^{Corolla} ~~corolla~~ ^{colore} ~~colore ^{lacio} ~~lacio ^{florentina} ~~florentina: ^{tubum} ~~tubum~~ ^{longitudine} ~~longitudine ^{fere} ~~fere ^{spatha} ~~spatha; ^{saucis} ~~saucis~~ ^{supra} ~~supra ^{tubum} ~~tubum~~ ^{incurvatis} ~~incurvatis~~: ^{limbus} ~~limbus~~ ^{campanula-} ~~campanula-
^{tus} ~~tus~~; ^{laciniis} ~~laciniis~~ ^{immaculatis} ~~immaculatis~~. ^{Stigmatibus} ~~stigmatibus~~ ^{laciniis} ~~laciniis~~ ^{bifidis} ~~bifidis~~. ^{saucis} ~~saucis~~ ^{l. c.} ~~l. c.~~

sesifolium - 9

gladiolus sesifolius Jacq. coll. 4. p. 164 n. var
gladiolus anceps Lin. (non Thunb.)

Habitat ad Cap. b. p. 4

Gladiolo anceps affinis, singulari folio sesif. Caulis exilitur anceps flexuosus, glaber, erectus, foliosus, spithameus. Folia alterna, amplexicaulia, ovata, inferora acuta, superiora obtusa, nervosa, subtus hirtula, margine crassa, lateribus reflexa, semuncialis. vel vixus folia conformia margine albis interiori sesif amplexicaulia quo sesifera in foliis superioribus ad marginem folii oppositum usque pertingit, unde hic nervus evadit folii medius dem ac diviso horizontaliter expanduntur non tantum sed etiam reflexuatis. flores solitarii exiti sesifis univales spathe bivalvis brevissima, membranacea alba: valvula externa ovato-acuminata, dorso medio viridi. Corolla tubus filiformis, prope incurvatus albis, superne violaceus. lacinus limbi lanceolato-peltato pallide violaceo, supra purpureo rubro basi albis limbo medio ad purpurea. Jacq.

affinis anceps

non vidi

Merianus 35

H. prothia oblongis
obtusiusculis

~~Gladiolus~~ ~~oblongis~~ ~~obtusiusculis~~ ~~capitulis~~
Meriana flore rubello Trew. chrol. t. 40

Antholyza meriana; corollis infundibuliformibus; foliis ensiformibus
Lin. spec. pl. 54. Syst. veg. 2. Mar. 87 80

Watsonia meriana Mill. Dict. 2. t. 276.

Gladiolus merianus; corollis cernuis subcylindricis, tubo duplici,
limbi laciniis ovatis, foliis ensiformibus glabris. Thunb. diss. n. 12

Gladiolus merianus; corollis infundibuliformibus, limbi laciniis
squalibus, ^{usq.} facie compressiuscula, foliis angusta ensiformibus
obliquis Faiqu. coll. 4. p. 150 Faiqu. Fe. rar. 2. t. 230

Habitat ad Cap. b. sp. 1

parviflora β. Faiqu. gladioli Meriani varietas Faiqu. collect. 5. p. 17. Fe. rar. 2 t. 231.

Habitat ad Cap. b. sp. 2

Solia radicalia saepe quatuor, scapo breviora, parum striata, margi-
nata. Scapus sesquipedalis, simplex, inferne foliis vaginatus, ses-
quipedalis, flexuosus, teres, striatus. Stem. quatuor - octo
_{vert.}

36
merianellus ~~35~~ *Gladiolus* corollis infundibuliformibus cernuis secundis, laciniis limbi
~~substantibus~~ <sup>foliis lineari-
oblongis, vaginis pilosis</sup>

gladiolus *Watsonia* humilis; foliis linear-filiformibus; tubo floris longis-
simis Mill. ic 190. t. 297

Antholyza merianella; corollis infundibuliformibus, foliis linearibus.
Lx. syst. veg. 89.

Gladiolus Merianellus; corollis cernuis, tubo duplici, limbi laciniis ovatis, foliis ensiformibus pilosis. Thunb. ^{glad} Diss. n. 11

Habitat in Cap. b. spec. 7

Solius, & caepo breviora, tres- quatuor, angusta, nervoso-striata, longe vaginantia, interdum pilosa, interdum glabra: vagina sulcata, semper pilosa: Scapus basi quipedalis, simpliciter erectus, superne flexuosus.

Floris. Flores tres- quatuor, secundis, cernuis. Spatha bivalvis, lanceolata, striata acuta, striata. Corollae tubus inferne tenuis, supra spatulas ampliatas & limbi laciniis oblongis

Falcatus gladiolus corolla limbo patenti, foliis lanceolato falcato Thurb. dig. p.

gladiolus falcatus caule simplici, foliis falcatis, floribus alternis, stylo trifido

L. Suppl. p. 96. t. 1. v.

Habitat in Cap. bono p. 17.

x

Capus compressus, strictus, flexuosus, glaber, palmatis. Solum 1-2. obvege
amplexi caule, ovato, lanceolatum, obtusifolium, integrum, ana
tum, glabrum, pollicare. Solum spathaceum in medio scapo. spathe
virides obtusa, tubo multo brevior. Corolla conica limbo pa
tenti Thurb. l. c.

altheoides 37.

tubulosus 36

Gladiolus corollae tubulosae reflexae, laciniis ovatis erectis, foliis ~~mucronatis~~
gissatis ensiformibus marginatis

Antholyza altheoides; corollae declinatae monopetalae vix dentatae genitalia
inclusentibus, foliis ensiformibus Burm. prodr. fl. cap.

Gladiolus Tubulosus

Habitat ad Cap. b. Spai. Ex herbario Burmanni et Royeni

vel althei

Tota glabra, Folia Alceae. Folia radicalia, tria - quatuor, sesqui-
pedalia, unguem lata, striata, margine serris incusata. Scapus
sepe bipedalis, simplicis simus, erectus, teres, vaginatus. Spica

spithamea, quinque - duodecim flora. ~~Rachis spathistata~~ Spatha alterna,

4 vix semipollicaris

subimbricata, disticha, trivalvis: valvula ~~exterior~~ ^{exterior} complicata, obovata,
striata, ~~interiores~~ ^{interiores} ~~lanceolata~~ ^{lanceolata} ~~gordide purpureascent;~~ ^{gordide purpureascent;} interiores lineari-

lanceolata. Corolla ~~patula~~ ^{patula} tubulosa, ~~crassitie~~ ^{crassitie} reflexa ~~crassitie~~ ^{crassitie} per-

no sesquipollicaris, ~~crassitie~~ ^{crassitie} diametro penno anserino, reflexa

tubulosa, reflexa, saturate coccinea: laciniis breves, ovatis, erectis

aequales. Stigmatibus laciniis bifido.

Jacq. coll. 4. p. 157

an potius Witsenii speculo

gracilis?

Gladiolus gracilis Jacq. coll. 4. p. 159 u. var.

Habitat in Cap. b. p. 4.

Totus glaber. Folia linearia striata longe superantia, supra vaginam
brevis canaliculata. Scapus inferne foliosus debilis, gracilis erectus.
Superne gemmulata, pedalis. Flores plerumque duo raro unus, binarius.
panem cernuus spatia vixit bivalvis: valvulis lanceolatis, acutis conica
in longitudine laevi corollae. Tubus cylindricus albus. Sicut in curva
infundibuliformis pallide violacea. Limbus pallide violaceus
calicinis lanceolatis acutis tria superiora duplo latiora linea saturatione
in dorso. duo inferiora tria puritatisque variegatis intermedia linea unice
pecta. Jacq.

an varietas *Gladiolus* Jacq.

mibi ignotus

Spec. Distinct. Jacq. w. ed.

viridis 30 Gleditsia corollae limbo patenti - reflexo, foliis ensiformibus
 glabris, stipite triquetro, angulis membranaceis Hort. Kew. 3. p. 481. +
Habitat in Cap. b. p. 2.
Singularis corollae viridibus, eadem striis obsolete purpureis
 Hort. Kew. l. c.

auct. a 46 foliis ensiformibus nervosis, bracteis indigenis, floribus racematis
Hort. Kew. 1. p. 57. +
Habitat in Cap. b. p. 2.

aulica 49 ~~fr. in foliis ensiformibus nervosis, floribus racemosis, bracteis integris~~
~~Hort. kew. t. p. 52. t~~
~~Habitat in Cap. b. Spec. 4~~

aulica fr. ~~fr. in foliis ensiformibus~~ ~~Hort. kew. t. p. 52. t~~
~~Habitat in Cap. b. Spec. 4~~

38 ~~39~~

crispus ~~37~~

laciniis linearibus obli-

gladiolis corollis in fundibuliformibus secundis, foliis lanceolatis ensi-
formibus undulato-crispis

G. 97.

~~gladiolis~~ crispus, foliis lanceolatis crenatis undulatis, floribus secundis,
spicis duabus, tubo filiformi longo. Lin. Suppl. 94.

gladiolis crispus; corolla erecta limbo campanulato, floribus secundis,
foliis crispis. Thunb. Diss. n. 7. t. 1. f. 2.

gladiolis crispus; corolla tubo gracili longioris, laciniis rotundatis,
foliis ensiformibus striatis glabris, ad oras crispis et barbatis Paigu
collect. Suppl. 27. Ic. rar. 2. t. 267.

Habitat ad Cap. b. sp. 4

^{JEANUS}
Caeni ~~Spithamensis, erectus~~, flexuosus, teres, glaber. folia quatuor-quin-
que, tripollicaria, striata, glabra. Flores quatuor-quinque. Spatha brevissima
obtusiuscula, denticulata, rubra. Corolla tubo sesquipollicaria, albida inae-
nata.

laciniis linearibus oblongis

Carneus gladiolus

Genus gladiolus carneus de la Roche desj. p. 30. t. 4. Jacq. coll. 2. pag. 370

Habitat in cap. b. p. 6

Totus glaber. Caulis pedalis simplex foliosus, flexuosus erectus.
Folia ensiformia acuta striata, rigidula erecta alterna disticha
Semi-pedalis ^{magis} castilaginea. Spatha oblonga-lanceolata
vires. Spica terminalis, 4 flora lassa ~~spica~~ flori incarnata.
4 pollicares tubus cuius vix spatium longior limbus inaequalis
lacini superior lateris omnes lanceolati Jacq.

39 40 49

biflorus ^{3^a} Opheodolus corolla erecta limbo campanulato, ^{scapo} ~~caule~~ bifloro, foliis linearibus

Thunb. Diss. n. 5 +

Habitat ad fretum magellanicum

Caulis ped. folia circiter tria, supremo longitudine scapi, infimo brevis summo longe vaginantis, glabra. Scapus pedalis, erectus, totus velut vaginis foliorum. Flores terminales, coloris tristici Thunb. l. c.

indifolius G

glad colus indifolius Taiqu. coll. 4. p. 156. et var

Habitat in Cap. b. p. 7

+
Idnea var
et generis
habes

~~Tolia radicalia caeteris 7. Diffusa capiformis, acuminata, lineata
nervo medio utroque protuberante usque pedalia vel minora in qua
luc. Scapus 4. ens. procerus crispis calami inferne vaginatus,
tripedalis. Spica 12 florum, erecta, teretes flexuosa tandem pedalis
spatha bivalvis valvulis lanceolatis acuminatis superne flavescenti-
bus cum rubedine. Corolla 3 unciali tubo longitudine spathae
laciniis oblongis lanceolatis acutis. Laciniis stigmatibus filifor-
mibus bifido Taiqu.~~

150112

4040

falcatus - Gladiolus corolla limbo patenti, foliis lanceolato falcato Thunb. Diss. n. 4

Gladiolus falcatus; caule simplici, foliis falcatis, floribus alternis, stylo trifido Lin. Suppl. 96.

Habitat ad Cap. b. Spei ?

folium unicum vel duo, pollicaria, obverse amplexicaule, ovato-lanceolatum, obtusiusculum, striatum, glabrum, spathaceum in medio scopi. Scapus palmaris, compressus, flexuosus, striatus. Spathae virides, obtusae, tubo multo breviores. Corolla coerulea: Limb Thunb. l. c.

ramosus & caule ramoso, foliis lanceolatis limbo patenti Thunb. Diss. n. 4
Habitat ad Cap. b. Spei

aletroides - *A. spica* disticha floribus tubulosis reflexis

~~Habitat in cap. b. spici~~

~~Antholyza aletroides corollis declinatis monopetalis vix
dentatis genitalia includentibus; foliis ensiformibus~~

~~Bur m. prodr. fl. cap.~~

Glad tubulosus

4 aequ.

coll. 4.

li. var.

~~Habit. in cap. b. spici~~

~~Folia plana pedalia. Scapus bipedalis simplicis pinn
glaber. Spatha altera spica furthamou. Spatha
alternis striata glabra subumbriata disticha acuta
oblique truncata emita. Corolla sesqui pollicari
tubus unguiculari spatia a basi angustior. Tubus
extrem spatiam pollicis parum longior craspti pen
no anfrina dependens lacinis brevis ovata subaequalis~~

an potius glad glus

41.
subulatus # Gladiclus corollae sublinguatae tubosae ^{obtusae} nervosis longiore, laciniis oblongis subaequalibus, foliis subulatis terdistiusculis.

Habitat in Cap. b. spec. Bulow. 7.

Scapus semipedalis, teres, tri- & tetraphyllus, vaginae foliorum testis.

Nervia, subtilis
me striata

Folia altitudine scapi, subfiliiformia, acutis ^{sinuatis}: vaginae anticae spicae sesquipollicari, distiche subimbricata, dimiter duodecim flora. Spathae vix unguiculares, obtusae, multinerviae, ferrugineae margine minime membranaceae. Corolla ^{semipollicari} ~~subulata~~ ^{subulata}

^{violacea} ~~testis~~; semipoll. ^{tubosae} ^{albidae} ^{spicatae} purpureae lineatae: laciniis acutis
superne curvatus An Gladiclus ^{spicatae} ^{alopeuroides} ^{secundum Thunberg} ^{lin?} sed illa similis # et est
Glad. ^{alopeuroideae} ^{cum quo planta} ^{descripta} nequidem
minimam affinitatem habet. An Linnaei ^{spicatae}

Gladiclus bracteolatus; foliis convolutis filiformi subulatis, floribus spicatis, bracteis alternis multinerviis spathis includentibus Lamarck. Musc.
Bot. bot. 2. p. 225. illustr. 1. p. 117.

42
rubeus 42 Gladiolus corollae tubo spathis acutis nervosis longiore, laciniis linearibus
oblongis subaequalibus, foliis linearibus nervosis.
Habitat in Cap. 6. & p. 7

Solia & Capos fera altiora, angusta, quinque nervia. Scapus, rachis
Et spathae fuscae purpureascentes. ^{Superna}

Similis antecedenti. Diversus: foliis planis nervosis, nec laciniatis
& tubulatis enerviis. & Capo altiora: Spathis acutis nec obtusis
Simis

~~Setosus C. Spica linearis oblonga, involucriis setosis, fetis inermibus; intinori
bus basi villosis, vallis ciliatis, glumis lobatis Sw. prodr. pag. 26.
Habitat in India occidentali~~

~~Culmus sesquipedalis genuiculatus ramisque ramo uno vel altero
inferne vaginis, uti folia, pilosae, ore barbata. Spica
4-5 pollicari ascendente
facie Benchni ciliani~~

Panicum alopecurus La Mart.

~~48. 42. 44. 41.~~ *Gladium* ~~H~~ ^H ensi formibus plicatis
viresco-cyanea ~~41.~~ ^{41.} foliis oblongo-lanceolatis scapo declinata longioribus, spatha
villosa, subaequalibus, corollae limbo patenti primo

Corollae limbo patenti primo, lacini
nisi oblongis, foliis oblongo-lanceolatis, villosis plicatis
scapo ~~longioribus~~ longioribus, spatha univalvi, corollae limbo patenti primo Gauche
viresco +

Collect. 3. ~~41. var. p. 260. 41. var. 2. t. 205. Curt. mag. t. 410.~~
~~41. var. p. 260. 41. var. 2. t. 205. Curt. mag. t. 410.~~
Habitat in Cap. v. p. 41. t. 16.

Tota^{us} exceptis corollae villosa. Sola spatham aca sel hemipedalea,
interdum parum latiora, interdum angustiora, basi attenuata,
trinerviata. Scapus foliis saepe duplo brevior, superiora interdum
foliis setaceo brevi, quodam quinq. floris^{us} spathae univalvi.
valvulae lineari, acutae; exterior reliquis longior. Corollae
tubus pallide rubens: limbi lacini subaequales, lacini
longae, alternae mucronatae, saturate cyaneae, basi rubrae,
Gauche. l. c.

apice parum flexuo
sus

Gauche

Tanta est huius affinitas aequae ac *Glaucis puniceae*, purpureae
villosae ^{41. var. p. 260. 41. var. 2. t. 205. Curt. mag. t. 410.} cum *Glaucis plicata*, bambulino et reliquis iis proximis
ut potius varietates ejusdem speciei esse appareant, quam speciei di-

Curt. 410

39 forte ut non nullae ad obliq. ad *Glaucis* transferenda propter corollae inaequalitatem
et nervi ab altero basi marginem basi exori.

^{us}
pyramidata ^{us} ~~caule pedunculato, foliis linearibus striatis distichis patenti-~~
~~bus: superioribus ciliatis, inferioribus brevioribus spatulatis La Mart~~
~~T. Eni: pag. 108~~

β humilior

Habitat in Gopula Franco β in Cap. I. p. 1

Distincti generis, nam in dictis gladiolis corolla non laciniata non magis
inaequalis, nec tubus magis curvatus quam in novindis ^{us} ~~us~~, si itaque
haec erunt ^{us} ~~us~~ generis, gladiolus plicatus, uti ^{et} reliquae illae trans
ferendae ~~us~~ ~~us~~ sunt

non vid.

44 44 44 44

~~Sulphureus~~ ^{villosus} ~~Gladidius~~ ^{Gladidius} corollis campanulatis secundis, laciniis lanceolatis undulatis, & caps ~~dentatis~~ foliis nervoso-plectatis, scapo declinat +
gladidius sulphureus; corolla campanulata; foliis ensiformibus, plicatis, hirsutis; spatha trivalvi; valvulis omnibus acutis, integerrimis Jacq.

collet. 3. p. 258. in var. f. t. 239
Habitat in Cap. b. spec 7

^{ensiformis}
Folia quadri-quinque pollicaria, erecta, ^{acuta}, plicatula, nervosa, villosa, mollia: ^{scapo pedalis} vaginis arduis, villosis, infima purpurea & cetera. Scapus pedalis, declinatus, ^{terres}, villosus. Flores cincti
septem, ^{bipollicares} alternis, secundis, sessile. Spatha trivalvis, lanceolata, acuta, villosa: valvula externa reliquis duplo longior. Corolla tubus ^{incurvus}, rotaceus: limbus ^{exterior} campanulatus; laciniis ^{longe lanceolatis}, obtusis, pallide sulphureis, alternis, & opius undulatis. Jacq. l. c.

H. Spatha brevior
Sub aequalibus

+ 38. Jacq. coll. 3. p. 258
Tuba ensiformis longitudo tubi

~~Senecioides g.~~

~~gladiolus senecoides Jacq. coll. 4. p. 160 G. rar.~~

~~Habitat ad Cap. b. sui. 4~~

~~Stolonum radicele unicum, caulina alterna, inferiora duo interdum opposita
bulbus ampiformis striata 2-3 pollidaria altero fuso latere breviter vaginanti
obovato glauco vix glabra. Caulis semipedalis gracilis, glaber inferne ramosus
+ debilis flexuosus multiflorus. Flores alterni sessiles secundi spathe trival-
ves; valvulae linearibus, acutis conniventibus longiori bunciali intima
bifida. Tubus corollae pleiformis bunciali albus inferne incurvatus
Cilium eleganter purpureo laciniis lanceolatis Tubus inferioribus parum
minoribus basi macula ^{quo} flavescente macula utrinque sanguinea Jacq.~~

michi ignotus

* * Scapo polystachyo
- glabi

alopecuroides 46. Gladiolus spica imbricata disticha, foliis linearibus
formibus.

Gladiolo affinis, floribus minimis caeruleis dense spicatiis Rain
herb. 3. p. 560.

Gladiolus alopecuroides; foliis linearibus, spica disticha
imbricata. lin. am. acad. 11. p. 301. Syst. veg. 87.

Gladiolus alopecuroides, polystachyus, spica imbricata
disticha, foliis ensiformibus nervosis Thunb. gladiol. n. 14

¶ in plantaginea; foliis linearibus strictis, spica disticha
imbricata, & pappi marginis & Hort. kew. 1. p. 59.

Gladiolus alopecuroides; foliis linearibus nervosis, spica
subpolitana imbricata disticha, & pappi marginis & caris
per Lamarck illust. 1. p. 110. t. 32. f. 4.

Habitat in Cap. b. spec. 4

Bullelli plurimi. Scapus & pappi maculosa pedalis,

flexuosus. Solia duo vel tria in scape eoque breviora, latitudine vagis,
longe vaginantia. Spicae solitariae, interdum duae vel tres, digitales, erectae.
Siliquis copiosis, parvis. Spathae ovatae, compressae: valvula exterior
parum longior, dorso fusco, marginibus membranaceis hyalinis.
Variat floribus albis et caeruleis.

setifolus 46 Gladiolus corollis nigentibus, foliis lineari setaceis +

Gladiolus setifolius; caula ramoso, corolla nigente, foliis infimo filiformi-
setaceis. Lin. Suppl. 96.

~~Gladiolus setifolius; polydactylus, foliis lineari-setaceis Thunb. Desc.~~
n. 19 glad.

Habitat in Cap. b. spec. 2

Folia civitate ^{trio}, sensum breviora, infimo longissimo, scapum exante.
Scapus ^{palmaris} raris semisimplex, flexuosus ^{us} erectus; glaber: Rami teretes,
erecti. Flowers Corolla tubus spathis vir longior, Thunb. l. c.

flavus 9-folius lanceolato-ensiformibus planis, faucibus labii superiori
trilaminato: lamina unguiformibus perpendicularibus, bracteis acu-
minatis. Hort. Kew. 1. p. 654

Habitat in Pap. b. p. 7

Differt a 9. securigera bracteis acuminatis, corolla tota intense
flava, et foliis paulo paulo latioribus Hort. Kew

alatus 47 Gladiolus ^{collis ringentibus, laciniis spathu-latis, duabus lateralibus}
^{corollae tubo spathae}
^{majoribus, seapo subalato}
Gladiolus alatus; corolla divaricata ringente, lacinia summa falli-
 ta, foliis ensiformi-linearibus glabris, seapo vamoso. Facq. coll. 4.

9 p. 165. tz. rar. 2. t. 259

~~Gladiolus~~ ^{seapo} pythamou vel subpedalis, ^{simplex vel ramo auctus,} ^{superior} flexuosus, striatus, carina
^{inferne}
spatharum decurrente subalato; folia plurima ruia, unicum vel
duo in caule, infima caule sepe altiora, superiora brevia, angusta, tri-
quinque striata. Stores secundi, distantes. Spatha valvula exterior
tubo corollae duplo longior, lanceolata, compressa, obtusae, carina
et margine purpurea. Corollae tubus pallidus: limbis, laciniis cin-
rascens; lacinae superior lateralis superiores saturatius coloratae
~~Chalchabul~~ lineatae; inferiores basin versus flavae.
 # Videtur differre colore lacinarum superiorum coeruleo

inferne tere

odoratis simi

et subtilis pura
lineatae

9 Gladiolus alatus; foliis ensiformibus costatis; petalis lateralibus latis
sinis and. r. repos. t. o.
Habitat in Cop. 6. Spei

Sulphureus Gladiolus foliis ensiformibus saporque acriter tenato villosis

Gladiolus sulphureus Jacq. coll. 3. pag. 258. u. v.

Habitat in C. b. p. e.

Folium ensiforme acuta erecta 4-5 polliciana utraque
villosa mollis, plicatula nervosa: vagina apiciter villosa
infimo purpurascens. Caput 3-4 pedalis villosus dentici-
natus. Flores sepales alterni secundum: unctus 7. erecti bipetal-
licares. Spatha 3 valves lanceolata acuta villosa extrinseca
reliqui duplo longiore: Tubus nervosus violaceus: limbus
campanulatus lacinus longe lanceolatus, obtusi pallide
sulphureus subsqualis albione sapor undulato Jacq.

not
non videtur

permeabilis 48 Gladulus corolla ringens, laciniis spatulatis longe unguiculatis, duabus lateralibus latioribus, & capo tereti
gladulus permeabilis; caule ramoso, floribus imbricatis secundis,
foliis subulato-quadrangulis De la Roche Diss. n. 1. t. 2.
Habitat in Cap. b. spei. Ex herbario Royani &
Scapus bi-tripedalis, & simplex vel ramosus. folia radicalia

angustissima Duo vel tria, longitudine fere & capis; scapi sensim breviora vax
angustissima cyinaeformes, attenuata, nervo utrinque attenuato marginata.
flores plures, secundi, pollicares. Spathae bivalves: valvulae
lanicollatae, membranaeae, obtusiusculae, tubo breviores.
limbi fauce inflata, ventricosa, inter laciniarum unguibus
haerent, patulo permeabilis

Gladulus permeabilis Thunbergii prodr. 105. non idem?
descriptio laud. & h. comae in Act. Soc. hist. n. par. vol. 6 laud
& h. comae

India H. fructuosa laudanda in Suppl. p. 105.

alpinum E. culmi nudi triquetris, spica pappo breviora Len. t. V.
fl. Dan. tab. 620

Juncus alpinus bombycinus Bauh. prodr. 6. pen. 12
Hexagostis juncea alpina, capitulo parvo. tomento varioso Scheuchz.
gram. 385. t. 7. fig. 4.

Habitat in Leropo alpinis, locisque appenninis &
culmi filiformes struthamisi apophylli basi vaginati terminati
& spica ^{non} oblonga imbricata bractea subulata longitudine spicis
aequalta. squama lanceolata ferruginea

Habitat in Muntana ex ab alpibus Divesim. Culmi triquetri. Spica pappo
brevior

gramineus ^{spicis subtrifloris, ovatis}
49 *Gladiolus* ~~corollae laciniis~~ ^{pubae} ~~quolibet aristatis~~, ~~scapo paniculato~~,
~~ramo flexuosior~~

Gladiolus phalangoides ~~ethiopicum Planch.~~ 187
t. 122. f. 9.

Asphodelus foliis planis, caule ramoso, foliis acutissimis Mill. in Bot. 36.

Gladiolus gramineus; petalis lanceolatis petalaeo-acuminatis Lin. Suppl.

95

Gladiolus

~~grain~~

gladiolus gramineus; polystachyus, scapo laxo, spicis capillaribus flexuo-
sis, foliis ensiformibus glabris Thunb. Bot. n. 26.

Gladiolus gramineus; corolla tubo brevis sumo, laciniis subequali-
bus aristatis, scapo paniculato Farqu. collect. 2. p. 303. f. 2. r. 2. t. 236.

Habitat in Cap. b. 8/100

#3 Bractea duae
petalae ad basin
pedicellorum. & corollae
distantes.

~~Edo. totus glaber.~~ folia tria-
dex, unicum in scapo coeque breviora, ten-
uinem lato, tenuiter nervosa. Scapus pedalis et ultra, vix crassitie
pennis columbino, superne flexuosus, ~~erectus~~. Pedunculi divaricati

capella filiformes; pedicelli capillares, bi & triflori, ~~minuti albi~~.
Spatha ovata, acuminata, virescens, margine albo membranacea
corolla ~~virescens~~; tubo vix ullo: limbi laciniis ~~ovatis~~, ~~aristatis~~,
concoloris, subequalis, ~~erectus~~ carina virescente, in tuo linea ~~nervosa~~ ^{ca}

S³
sete ~~4/100~~ ~~totum~~

Y
menana A corollis infundibuli formibus, foliis exsiformibus L. V.

Watsonia Mill. 9. p. 276.

menana flore rubello Thunb. Eret. 440

† menanus gladiolus menanus corollis cernuis subcylindricis, tubo duplici:
tambi laciniis ovatis, foliis exsiformibus glabris Thunb.

+
70. var. Gl. n. 12.

Habitat ad Cap. 6. p. 12.
Scapus simplex teres striatus, inferne foliis vaginatus, erectus, flexus
filiis, pithamou. Folia plura, usque 4, obverso margine vaginon
tibi, obliqua, parum striata, nervo medio utrinque elevato, infra
gra marginata, scapo breviora. Flores 4-8 alterni cernui purpurei
Corolla tubo basi filiformi, prope amplicatus geniculis reflexis, inde in
flecto-cigano cylindricis, sequi pollicem laciniis latis ovatis
obtusissimo cum alaribus, fere in quinque, aequales, concavo
Thunb

70

Amst. L. 1790

ad Gladiol. referatur

involutus ⁵⁰ ^{et} ^{remotis distinctis,}
Gladulus corollae ringentibus tubo spathe brevior; laciniis lanceolatis
attenuatis lateralibus margine involutis

Gladulus caule ramoso foliis linearibus planis, spathe glabra acutiuscula Mill.
ic. 2. p. 150. t. 236. f. 1.

Gladulus involutus; caule ramoso, foliis linearibus spicis distinctis de la
Arch. Diss. n. 2. 13.

Gladulus bimaiculatus; foliis linearibus perangustis, petalis superiori-
bus brevioribus patenti reflexis; trium inferiorum lateralibus ma-
iculatis angustioribus Lamour. Dist. bot. 2. p. 727. illustr. 1. p. 116

Habitat in Cap. b. & p. 2

Scapus ^{ramis flexuosis;}
caulis sesqui pedalis; ^{erectus, teres;} articulatus, in ramos duos tres floriferos
divisus; glaber, ~~et tota planta~~. Folia striata angusta, striata; caulina sub
ramificationibus. Flores quinque - octo, distantes, distincti. Spathae bivalves;
valvulae lanceolatae pollicares, lanceolatae, laeves; ~~tenerae~~ attenuatae, tenerae
laeves, margine membranaceae. Corollae ultra pollicem longae; roseae: ~~tubo~~
spathe breviori: limbi lacinae tres superiores sa majores; ~~laterales~~ majores
involutae; ^{intermedia} inferiorum ~~intermedia~~ longior lance inferioras tres ~~patenter~~
intermedia latior lanceolata, laterales lineares, ~~non~~ inferna macula
purpurea

castalagenea 401a ~~foliis ensiformibus nervosis marginato-castalageneis, caule po
lystachyo, tubo spatii triplo longiore La Mart. T. Enc. pag. 112.~~
Habitat in Cap. b. p. r.

undulatus 5. ^{is saepe} ~~gladiolus cordatus subbracteatus tubo spathae longiore, laciniis subae-~~
~~qualibus undulatis.~~

~~gladiolus cordatus, undulatus Thunb. gladiolus~~
gladiolus undulatus; foliis ensiformibus, petalis subaequalibus laciniis
latis undulatis lin. Mant. 27.

gladiolus undulatus; corolla subbracteata; foliis ensiformibus; ^{p. 256.} caule flexuoso,
ramoso; laciniis petali undulatis Fourq. coll. B. F. C. var. 2
+ 257.

~~Habitat~~
gladiolus undulatus; polytachyus foliis ensiformibus, laciniis
limbi undulatis; tubo spathae longiore Thunb. prodr. 166. Act. Soc. h. n.

~~habn.~~
Habitat in Cap. b. sp. 4

Folia circiter duo, glabra, in scapo breviora, longe vaginantis, apice
attenuata, multinervia, glabra. Scapus ^{pedalis - bipedalis} rarius simplex, saepe
us superne ramosus, caulis flexuosus. Flores bini - ~~trini~~ ^{trini} spathae
convolutae, pollicares, tubo breviori, oblongae, acutae, glabrae.
Corollae tubus filiformis, bi-tripollicaris, rectus: limbi lacinae
lineari-caudatae, undulatae. Thunb. l. c.

+ Secundi

tubiflorus Gladiolus polytachyus, tubo corollae longissimo, spathe foliisque ellipticis
plicatis, hirsutis Thunb. Diss. p. 20.

~~capa paniculata La Roche Diss. pag. 261.~~

~~Gladiolus tubiflorus caule recto, tubo longissimo, spathe foliisque hirsutis Lin. Suppl.
p. 96. S.V. pag.~~

+ + Habitat in Cap. bono spec. 2 man.

seperne villosa-tomentosa

7. Affinitas

int. ex tubato
et tubifloro

Capus ~~capae~~ pollicaris, ^{pateman} foliis involutis, ^{seperne villosa-tomentosa} hirsutis. Folia ^{angusta} erecta, b. lineari-ensiformia,
utrinque attenuata nervata, erecta, scapo longiora. Bractea oblonga hirsuta
apice ^{attenuata} ferruginea, pollicaris. Flores alterni, inaequaliter: tubo filiformi
apice ampliato curvo Thunb. l. c.

+ gladiolus tubiflorus foliis lineari-lanceolatis villosis plicatis scapo longiori
bus, tubo longissimo spathe hirsuta Hort. Kew. 1. p. 62. *

angustus 52 Gladiolus corollae subringentibus secundis, laciniis ~~linearibus~~ ^{linearibus} candido
tubus in ferientibus inferna macula rhombica sanguinea, flaps antipeta

^{caule simplicis primo}
Gladiolus africanus, foliis gramineis, floribus carnis, macula rhom-
boides in scriptis Boekh. Lugd. b. 2. p. 127.

Gladiolus foliis linearibus Lin. hort. cliff. p. 20. t. 6

44 Gladiolus angustus; foliis linearibus, floribus distantibus, corollam
tubo limbo longiore Lin. Syst. veg. 8^o mill. ii. 1. p. 95. t. 142. f. 2.

Gladiolus angustus

Habitat in Cap. b. spai 7

Scapus pedalis vel sesquipedalis, aneps. Foliis caule vix latiora.

41 res - quatuor Flores secund. # Corollae ^{flaps antipeta} tubus ^{flaps antipeta} tripollicaris. limbi lacinae
caudatae, aiuteae, macula rhomboides, sanguinea superne in
laciniis inferioribus linea que ejusdem coloris ad faucem ducit
vens.

Gladiolus; floribus patentibus externe carnis, interne candidis
stalla maculis que purpureis insignitis, suprema lacinia maxima
Breyer. ii. var. 21. t. 7. f. 2.

Longiflorus Gladiolus polytachyus, tubo corollae longissimo, spathe obtusa, foliis
que caespitiformibus glabris Thunb. Desf. pag. 19.

Ixia paniculata La Roche Desf. pag. 26. Tab. 1.

Ixia longiflora polytachya, tubo corollae filiformibus longissimis nutantibus
Berz. Cap. pag. 7

fort e inter
of 9xw

Gladiolus longiflorus caule tereti; tubo longissimo, spathe folisque linearibus glabris
Lan. Suppl. pag. 96. L. V. pag.

4 or 5

Habitat in arenosis ad Cap. b. p. 24

coll. u.
var.

Scapus teres, interdum simplex, glaber, pedalis. Folia 3-4. nervosa, scapo breviora
Flores alterni, albo viciis, plurimi, approximati. Bractea membranacea Anato
brevissimo. Tubus corollae saepe bipollucaris apice ampliatum, cernuus erectus
Thunb. l. c. Lacinae lineari lanceolatae, tenores obtuse ciliatissimi

Longiflorus *I. foliis* ~~longiflorus~~ ~~longiflorus~~ ~~longiflorus~~

Longiflora *Ixia*. foliis caespitiformi linearibus spathe tubo filiformi longissimo

Hort. New. l. p. 50

in montis collina sitiente n. ex her. es. con nata in sy. Gaique
u. m.

blandus, B^o Gladiolus corollae ^{is inaequalibus} tubo compresso, laciniis ^{suprema} reflexa, tribus
~~inferis linearibus lanceolatis~~, & stigmatibus & subbilobis +
gladiolus blandus; foliis linearibus lanceolatis nervosis glabris, floribus spi-
catis; laciniis ^{suprema} reflexa, stigmatibus & subbilobis Hort. Kew. 2. p. 64.
Habitat ad Cap. 6. sp. 4.

Corolla tubo sesquiuncialis, angustius compressus, parum incurvus, palli-
di rubens: limbi laciniis ^{suprema} longitudine tubi, oblonga, acuta,
medio concava, ibique dilatata, supra medium reflexa et albo
incarnata; duo laterales ^{suprema} paulo breviores, oblongo-lanceola-
to, patentes, apice reflexo, incarnata; tres inferiores ^{superiores}
paulo breviores, linearibus lanceolatis, erecto-patentes, supra medium
parum reflexo ex albida incarnata, in medio macula rubra
transversa notata; intermedia ^{caeteris} alutior. Stigmata dilatata,
biloba, margine villosa. Hort. Kew. l. c.

Juncus 54 Gladiolus species secundis, foliis lato-lanceolatis subundulatis
Stigmatibus lufid Stigmatibus Stylo sexpartito Stigmatibus lufidis Stylo sexpartito #
Gladiolus juncus; foliis lato-lanceolatis, culmo ramoso, floribus secundis,
Stylo sexpartito lin. suppl. 94

~~Folia quatuor sex, basi breviora, subundulata, uti tota planta glabra.~~
~~Scapus palmaris, interdum simplex, filiformis~~

Gladiolus juncus; polystachyus, floribus secundis, foliis lanceolatis

Thunb. ^{glab.} sess. n. 10.

Habitat in Cap. b. spec 4

Gladiolus marino Folia quatuor sex, basi breviora, subundulata, uti tota planta
ratus; foliis ensi glabra. Scapus palmaris, interdum simplex, filiformis, flexuosus,
formibus nervis Stylis quatuor septem, erectis, coenulescentes. Bractea brevis bina
bis glabra manu l. l. Thunb.

losis, floribus disti
chis, Stylo sexpartito
Linnae dict. bot. 2
p. Willd. r. s. p. 110

marginatus 55
Glyptadiolus: spicis adpressis pluribus, ^{floribus cernuis} ~~tubo incurvato~~, foliis ensi-
formibus nervosis marginatis, & ligmentibus bifidis

Gladidiolus maximus ramo
Sui floribus ar-
gentis Plukn.
mand. 91. fide
herb. Plukn.
Gladidiolus marginatus; foliis cartilagineis marginatis glabris multinerviis,
& spica elongata, floribus alternis nudantibus Len. Suppl. 95
Gladidiolus marginatus; polystachyus, spica longis sumis, floribus alternis
cernuis; foliis ensiformibus multinerviis marginatis Thurb. Prod. n. 20.
Folia marginata; polystachya, foliis ensiformibus nervosis margine
incrassatis, spicis pluribus adpressis, tubo incurvo, & ligmentibus

Habitat ad Cap. b. spec. 4

^{Tuber glandis}
^{Tuber gla}
Folia pedalia, ^{semi} pollicem ~~lati~~ ^{erecta} vel ultra ^{erecta} ^{erecta} lata; nervi in medio et ad
margines ~~crassiores~~ ^{incrassati}, ~~flavescentes~~, disci tenuiores. Scapus inferne
foliis bipedatis, inferne foliis vaginatis. Flores cernui spica semipedalis.
ultra Flores cernui, ^{magni} remotiusculi. Spatha valident ^{an} ^{externis prolixiis,}
~~lacu colato in gutture attenuata, striata, corallo tubo perispermio longior.~~
^{longior}
Tamen colore flavo.

Axio cartilaginea; foliis ensiformibus nervosis ^{marginatis-} cartilagineis
Coula polystachya, tubo ~~spathis~~ ^{longior} ^{longior} Lamort. Dict. t. p. 340 illustr.
i. p. 112

crispus gladiolus corollae erectae limbo campanulato, floribus secundis, foliis crispis

Thunb. Diof. pag.

* gladiolus crispus foliis lanceolatis crenatis undulatis, floribus secundis, spicae
tubo filiformi longo Len. Suppl. pag. 94 * L.V.p

Habitat in montibus Cap. bono spici 2 unq alterove folio crenato lanceolato
scapulis teres flexuosis erectis glaber, spithameus. folia 4 anguliformia, glabra
scapo breviora. Flores 4-9 spathe brevis formis obtusis, margine scapo

400 of coll. 4
42. nr.
denticulato, rubro. Corolla albo incarnato, erecto tubo sesquipolli
cani Thunb. l. c.

ab 9x. ~~undulatis~~ crispis affert caule folios, corolla tubus glumis octes longior
petales lanceolatis, caule altiore len

^{int est} folium 2-3. in scapo superiore linearis vaginans magnitudine pla
nis

glumaceus 56. Gladulus spicis pluribus erectis, corollae tubo filiformi spathae brevioris, foliis ensiformibus margine incrassato

Gladulus marginatus purpureus Thunb. Glad no 20 d
Gladulus glumaceus; polystachyus foliis ensiformibus multinerviis marginatis, tubo filiformi spathae brevioris Thunb. prod. 1 P 6.

Act. Soc. h. n. hofn. vol. 6.

Habitat in Cap. b. spici 4

Totus glaber folia duo vel tria, scapo brevioris, attenuata, marginata & multinervosa; nervis crassis flavescentibus. Scapus bipedalis, erectus, teres, foliis vaginatus, apice ramosus. Spica spathae mox & ultra; spicae laterales breves. Spathae unguiculatae, membranaceae, convolutae, lanceae, acuminatae. Corolla campanulata, purpurascenti-coerulea: tubus filiformis, spathae brevior.

Thunb. l. c.

Spathae in setam attenuatae, striatae

testaceus 57 Gla diolu - spicis subsoliteris, tubo filiformi longitudine spathae,
4 subfoliatis corollis campanulatis, foliis lanceolato-ensiformibus
Antholyza spicata; floribus infundibuliformibus, spicatis, in
carinato-pallidis; foliis falcatis, brevissimis Andr. reposit 56
Habitat in Cap. b. spei 4

Secundum figuram affinis videtur Glad. in difolio et distorsus.
spicis geminis; foliis brevibus; spicis geminis; corollis minoribus.

luc. ensif. lac. ovatae acutae tubo filiformis longitud. spath.

Aethiopia A. corollis incurvatis: Labii quinquepartiti lobis duobus alternis
patalis majoribus lanceolatis *Her. Spei: pl. p. 547*
glad color *Aethiopicus*; flore coriineo *Corn. Canad. 79. 7. 79*
monf. her. 2. p. 421. f. 4. 7. 23. fig. 1.

Hyacinthus africanus; foliis colchici: floribus coriineis
Plukin. Tab. 195. fig. 2.

Habitat in *St. Cap. b. Spei 7*

Folia basi vaginaria, ensiformia disticha fol
licem fere lata. Corolla tubulosa coriinea Labio
superiore ^{lobis alternis brevioribus} lanceolata inferiore 5 partito lobis lan
ceolatis *Her. Spei* ^{lobis} *St. Cap. b. Spei 7*

Aethiopia H foliis ensiformibus utrinque attenuatis nervo
si, *Spei* a *St. Cap. b. Spei 7*

Dichotomus 58 Gladiolus scapo bis bifido, ramis bifloris, & corollis campanulatis, scapo
bis bifido, ramis bifloris, foliis linearibus +

Gladiolus dichotomus corolla erecto limbo campanulato, scapo bis dichoto-
mo quadrifloro Thunb. bot. Zeyl. n. 6.

Habitat in Cap. b. p. 2

Solum basi vaginans, scapo longius, glabrum; spathaceum folium sub
dichotomus, lineare acutum. Scapus pedalis, apice bifidus, toros, acutus
glaber, brachiis singulis bifloris. Flores secundis erectis. Thunb. l. c.

~~nerosus Glad. foliis angustiformibus plicato nervosis adlopi, racemis
plumbis alternis, tubo spatii breviora La Mart. T. Enc. pag
119
Habitat in Cap. b. p. 2~~

non vid.

59-60

bicolor 59 Aladiolus corolla virgata, spathe lacero-aristata, foliis ensiformibus
~~glabris~~ Thunb. Diss. n. 16. t. 2. Faerq. collat. Suppl. 25. 42. var. 2. f. 240

Habitat in collibus ad Cap. b. 8. p. 7

Bulbus ovatus, reticulatus. Folia scapo duplo breviora, alternatim vaginatis,
obversa, obtusata ^{cum} acumine, striata. Scapus 8 pithamos, basi
foliis vaginatus, ~~et~~ flexuosus, angulatus, striatus, sulcis utriusque
profundo excavatus. Spicae duae; altera uniflora, tenuior, altera
triflora crassior, longitudine dimidii scapi. Rachis angulata, ~~florosa~~
60. Spatha membranacea, apice lacero-partita, venosa, basi grisea,
apice ferruginea. Corolla tubus ampliatas, Spatha duplo longior,
filiformis, sensim ampliatas: limbi laciniis flavo; suprema major
ovata, concava, apice coerulea; laterales inferiores lanceolatae, con-
voluta, ultimo apice coerulea; in juna brevior, panem a reliquis se-
parata, extra laterales collocata: omnes basi linea duplici purpu-
rea in ore tubi Thunb. l. c.

Carinatus 9. foliis linearibus utrinque carinatis glabris, tubo spatulis em-
busque brevioribus, stigmatibus indivisi complicatis Hort. Kew. 1. p. 644

Habitat in Cap. b. p. 2

x

Caulis sesquipedalis teres. Folia caule longiora, flores subsessiles. Tubus
subcylindricus femuncula paulo longior albidus. Limbus subcampanu-
latus: subnervos. Lacinia superior obovato-oblonga, tubo duplo longior
pallide rosacea, superiora angustiora, intermedia longitudine superioris
nisi infra medium flavicanti. Lacinia brevior medioflava Hort. Kew.

60 60
Silencoides

exsertis

Gladior corollae tubo filiformi elongato, genitalibus longitudine
axillaribus, foliis linear-ansiformibus +
gladiolis silencoides; corolla tubo filiformi longissimo, floribus sessilibus
corolla Jacq. collect. 4. p. 160. f. c. var. 2. v. 270.

Habitat ad Cap. b. spec. 7

Scapus

Rubus obovatus. Caulis semipedalis, gracilis, glaber inferne ramosus;
debilis, flexuosus, multiflorus, glaber. Folium radicale unicum; caulina
inferiora duo, interdum opposita, bi-tripollicaria, breviter vaginantis
striata, glauco-viridia, glabra. Flores alterni, sessiles, secundi. Spatha
trivalves: valvulae sublineares, acuta, conniventes; longiori bin-
ciali, intima bifida. Corolla tubo biuncialis, albus inferne incur-
vatus: limbus eleganter purpureus; laciniis lanceolatis, tribus inferiori
bus parum minoribus, basi macula flavescens, et quae macula utraque san-
guinea Jacq. b. in duabus lateralibus Jacq. l. c.

rumela 9. foliis eniflor mebus glabris, fcapo unifloro petalis oblongis obtusis

Thurb. Desf. per Fr. n. 3.

1.2 Plaf. hist. 1. pag. 225.

Chamaeni latifolia minor Plaf. hist. 1. pag. 225.

~~Chamaeni minor, flore purpureo Bauh. pin. 33.~~

~~Ani sola
cum parva~~

~~Ani rumela corollis lobatis, caule foliis brevior uni-
floro lin. rei: pl. pag. 56. Gayn. f. l. aust. tab. 1.~~

~~Chamaeni flore purpureo caeruleo Bauh. pin. 33.~~

~~Chamaeni
latifolia minor~~

~~Chamaeni minor flore rubello Bauh. pin. 33.~~

~~Chamaeni flore pallido et albo Bauh. pin. 33.~~

~~Chamaeni latifolia
bo L. minor~~

~~Chamaeni minor flore variegato Bauh. pin. 324~~

~~Ani angustifolia maxima minor Bauh. pin. 33 ^{1/2} _{sub.}~~

~~Lobel.~~

~~Chamaeni flore luteo et pallide luteo Bauh. pin. 324~~

~~Lobel. ic.~~

~~Chamaeni saxatilis gallica Bauh. pin. 34. 324~~

~~Ani per ps.~~

~~Habitat in Austria, Pannonia collibus apricis y
scapis foliis vaginatis simplicibus, unifloris, palmatis. Folia
circa et sicut pinnata glabra scapum sequentia interdum
breviora~~

var. red. in Sept. V.

fissifolius † Gladiolus corollae tubo filiformi laciniis lanceolatis, & capso anaspiti
†, foliis ovatis undulatis, subtus subhirsitis +
gladiolus fissifolius; corolla tubo filiformi longissima, floribus sessilibus
aurillaribus, foliis hirsutis, in superiore margine fissis, superioribus
quidem ad oppositum marginem usque, in inferioribus autem ad apicem
indivisis Jacq. coll. 4. p. 164. v. var. t. 260

Habitat ad Cap. b. p. 4

Gladiolo anaspiti affinis, trigulari foliis fissis. ^{Scapus} Caulis erectus, spithameus,
aniceps, erectus, flexuosus, ~~foliolatus~~. Folia semundulata, amplexicaulia,
ovata, inferiora acuta, superiora obtusa, nervosa, subtus hirsuta,
margine crispa, lateribus reflexa. Vel venis folia ensiformia,
margine altero interiori fissis amplexicaulia, quo fissura in foliis
superioribus ad marginem folii oppositum sive exteriorem pertinet,
unde hic marginis erodit nervus folii medius, dum ora divi-
sa horizontaliter expanduntur non tantum, sed etiam reflectuntur.
Flores solitarii, erecti, sessiles, uncialis. Spatha bivalvis, brevissima,
membranacea, albida: valvula exterior ovato-acuminata, dorso

medio vindi. Corolla tubus paucum incurvatus, albus, superne violaceus: Limbi ca-
cinis lanceolata, subtus pallida violacea, supra purpureo-rubra, basi albida,
linea media atropis citro-purpurea Jacq. l. c.

62 ~~floribus~~ ~~longis~~ ~~imbricatis~~ ~~concolor~~
bracteatus 62 ~~Gladiolus~~ ~~strobilifer~~ ~~tubo~~ ~~filiiformi~~, laciniis ovatis obtusis, ~~scapo~~ ~~erectis~~
~~presp.~~ ~~foliis~~ ~~ensiformibus~~ ~~obtusis~~ +

Gladiolus bracteatus; polystachyus, scapo ^{apice} ~~erectis~~, corolla laciniis ovatis, spatibus ovatis imbricatis Thunb. prod. 106 Act. Soc. h. n. vol. 6. t. 4
^{cum figura}
Habitat in Cap. b. spec. 7

Totus glaber. ^{scapo} ~~Caulis~~ ~~spithamou~~, compressus, erectus, ramosus. Folcia
subramis sessilia, digitorum, ^{patentia} ~~ensiformia~~, obtusa, patentia vel subvix
vato-foliata, multinervosa. Flores plunni, ~~imbricati~~, ante anthesin
spathis ~~rectis~~. Spathis unguicularas, rhombos, ~~acuto~~, convolutis, integris.
Corolla alba: tubus filiiformis, erectus, apice geniculatus, dein amplia-
tur in limbum ~~rectum~~ ~~antem~~: limbi laciniis ~~ovatis~~, ~~obtusis~~ Thunb. l. c.

#. ovatae

floribus imbricatis, corollae laciniis ovatis, spatibus rhombos^{is} ovatis,
scapo compressis, foliis ensiformibus obtusis +

Fabrii 68 Gladiolus corollae tubo filiformi laciniisque filiformibus, scapo
non ancipiti, foliis spithisque carina integerrimis +
Gladiolus Fabricii; polystachyus scapo ancipiti, laciniis corollae fili-
formibus Thunb. prodr. 186. Act. Soc. h. n. Hafn. vol. 6

Habitat in Cap. b. Spei

~~Foliis~~ Caulis spithameus vel ultra, inferne teres, superne aneaps, ramosus.
Rami compressi, divaricati, erecti, virgati. Folia radicalia circumscissa
scapo breviora, ensiformia, acuta, multinervia; ^{scapi} caulibus sub ramis dimi-
cia, unguiculatis, angustiora. Flores plures, alterni erecti, albi - spathe
ovata linear longis, ovata, obtusiuscula. Corolla tubus pollicaris, fili-
formis. Lacini laciniis lanceatis Thunb. 6.

Thunbergius huc revocat Gladiolum Fabrii in La Roche, sed in illo angu-
li caulis dentati sunt, quod non in Thunbergii quantum a descrip-
tione conijcere licet, laciniis quoque ^{corolla} in tubo ancipiti non fili-
formes.

juniceus gladiolus polystachyus, floribus secundis, foliis lanceolatis Thunb. Disj. p.

gladiolus juniceus foliis lato-lanceolatis, culmo ramoso, floribus secundis, stylo sexpartito Lin. Suppl. pag. 944 * L. V. pag.

Habitat in C. bonapartei ?

+ Scapus interdum simplex, filiformis, flexuosus erectus, palmatis, laxis. Folia
4-6 lanceolata, subundulata, glabra, scapo breviora. Flores 4-7. sepalis erecti
conspicuis. Bracteis brevissimis. Th. l. c.

anceps ⁶ ~~4~~ ^{Scapo} Gladiolus corolla tubo liliformi, ^{caule} ~~caule~~ ancepsiti flexuoso, foliis
ensiformibus, ^{que carina extensorum} spatha ~~marginis~~ ^{exterioris} denticulata.

Atria Fabricii; caule dichotomo ancepsiti La Roche Diss. n. 5.

Gladiolus anceps; foliis ensiformibus undulatis, caule ramoso ancepsiti
divaricato Lin Suppl. 94. Gaerq. collect. 4. p. 163. fr. rar. ^{glad.}

Gladiolus anceps; scapo ancepsiti, spathis crispis Thunb. Diss. n. 17.
Lapeyrousa compressa ^{Poulet in act.} ~~caule~~ tolos. 3. p. 79. v. 6.

Habitat ad. Cap. b. spei 4

Scapo ^{scapo} Paulus palmaris et ultra, ~~anceps~~, angulis denticulatis: Rami simplices,
caule ^{bimiles}. Folia sesquipollicaria, amplexicaulis, conferta, acuta.

* Subpollicares, flores solitarii alterni. Spatha bivalvis: valvula exterior similis foliis
^{gen 3 in mino} ^{res} caulinis; obtusa, ^{marginis} ~~exterioris~~ ^{superne} crispa denticulata, apice
perurea. Corolla tubus sensim ampliatus; spathis longior, pallida

^{folia can} ^{ina carina} ^{decurrentis} ^{neato} folia canina carina decurrentis
coerulescens: limbi laciniis subaequalibus, ~~lineatis~~ lanceolato-pubis

Gladiolus denticulatus; foliis ensiformibus obtusis carina denticulata
decurrentibus, caule paniculato ancepsiti Lamour. illustr. 1. p. 110

Luteo-picea Anis corollis barbatis, caule unifloro, foliis longioribus subo in
protham inclusis La Mart. T. Eni: pag. 122.

Habitat in gallica germanica 4
Mant 134 Chamais latifolia minor y. Plus. hist. 227

~~quis perenni-
Hort. Exot. 2. 20. f. 6~~

65-66

refractus Gladialis ~~in fundibuli formibus~~ ^{in fundibuli formibus} ~~spithaei brevis semis~~ ^{spithaei brevis semis} ~~foliis linear ensiformibus~~ ^{foliis linear ensiformibus}

Gladialis refractus; corolla subnigra; foliis linear-lanceolatis ensiformibus, glabris; scapo ramoso; floribus & cavis, erectis, resupinatis

Hacque suppl 26. G. rar. 2. t. 241.

Habitat in Cap. b. spai. Ex herbario Royeri ?

Totunglae. ^{scapus} Caulis erectus pedalis, crassitie fili emporatici Yemionis, caulis, teres,
diphyllus monodistichus, glaber, laevis. folia Spithamaea vel mino-
ra, caulina breviora, angusta, leviter striata. Spicae remotae, tri-
septem flori. bracteis flavis. Olorea secunda. Spatha bivalves,
tubo fere quadruplo breviores; valvulae ^{aequaliter} subrotundae, membranaceae,
albescentes; ^{exterior} apice macula una ^{interior} (vel duae) fusco-purpurascens, ^{exterior} purpureo-
striatae. Corollae tubus croceus

purpureo-striatae

luteus **G** Gladiolus Corolla limbo campanulato ^{sub} spicis secundis, corolla lim-
bo campanulato tubo longitudinali ~~spathis~~, foliis brevioribus, foliis li-
nearibus angustis longissimis, floribus spicatis
^{negari} ^{angustioribus} ^{longioribus}
Gladiolus luteus, foliis linearibus angustis longissimis, floribus spicatis
subsecundis luteis, tubo curvo spatha breviora Lamark. Dict. bot.
2. p. 725. illustr. 1. p. 117.

Habitat in madagascar. Com. merson. Ex herbario Tussioi &

Talus glaber. ^{scapus} Caulis pedalis, teres, scandens, teres, glaber, simplex, interdum ramo
unico divisus. Solia caule altiora, basi ensiformia, striata.
Flores spicati, plurimi, approximati, lutei. Spatha oblonga, acuta.
Corolla tubi longitudinali spathae, curvatur: Limbus unguiculans.
vel parum ultra

ventricosi Glad. foliis cymiformibus nervosis glabris, limbo
ventricolo difformi, styli laciniis dilatato-membraceis
spathulatis La. Mart. T. Enc. pag. 110. x
an Glad. carneis Jacq. Coll. 2. c. var. 2.
Habitat in C. b. p. c.

subplenus, angustifolius et strictus La. Mart. pertinent abque dubia
ad species an Systemate recent.

67-68
secuniger

^{fauces}
Gladiolus corolla labia superioris trilaminato, laminis ungui-
formibus, spathis obtusis, foliis linearis-ensiformibus +
Gladiolus secuniger; foliis linearis-ensiformibus planis, fauce
labii superioris trilaminato: laminis unguiformibus perpendiculari-
bus, bracteis obtusis Hort. New. 1. p. 65. Pub. Lit. mag. 3 P. 3
Habitat in Cap. b. spci 2

Folia spithamea vel longiora; ~~glabra~~ Scapulis foliis paucis
brevior, interdum ramosus, teretiusculus. Spathis breves, ovatae
obtusae, interdum apice incisae, membranaceae. Corolla palli-
de fusca: tubus unci brevis: limbi lacinae oblongae, ovatae,
obtusae, interdum obsolete emarginatae, tubo breviores; supe-
riores tres in faucibus macula flava, margine rubicundo,
ubique aucta laminis tribus compressis, obtusis, perpen-
dicularibus, flavis, diametro sesquilineari Hort. New. l. c.

Xiphium 4. foliis ensiformibus canaliculato-fimbriatis scapo bifloro brevioribus
Thunb. fr. n. 17.

4ni bulbosa latifolia, caule donata Bauh. pin. 30

4ni *Xiphium* corollis imbricatis, foliis fimbriatis canaliculatis
caule brevioribus Lin. Spec. p. 50.

13 4ni bulbosa caeruleo-violacea Bauh. Pin. 40

Habitat in Hispania vagina foliorum vestis

Scapus sesquipedalis, crassitie pennis anfractu. folia angusta pa-
rim attenuata a basi ad apicem usque canaliculata. Spatha
superior ventricosa. floris caeruleis, petala emarginata.

Plus. hist. 1
210 et 849.

⁴ⁿⁱ caule donata Bauh

~~4ni bulbosa latifolia~~

4ni *Xiphoides* pedunculata 4ni anglicana major hort. Ey Ost. 0.4 f. 0

4ni bulbosa anglicana candido flore ib. 0.4. f. 7 ^{Fach.}

4ni anglicana major violaceo flore ib. fol. 0

4ni bulbosa dilute caerulea b. 10 Math. Bauh

4ni bulbosa anglicana flore caerulea not B

hui
omni
bus
speci-
bus

flavus ~~69~~ Gladiolus corollae faucis labii superioris trilaminatae, laminis unguiformibus, spathis acuminatis, foliis lanceolato-ensiformibus
gladiolus flavus; foliis lanceolato-ensiformibus planis, faucis labii superioris trilaminato: laminis unguiformibus perpendicularibus, bracteis acuminatis Hort. New 1. p. 65.

Habitat in Cap. b. spec. 2. c

Differt a Glad. secuniger: bracteis acuminatis, corolla tota intensa, flava et foliis paulo latioribus Hort. New 1. c.

Folia stricta, saepe longitudine Scopii, interdum breviora. Scapus septem-
 decem pollicaris, interdum simplex, saepe ramosus ut flores quatuor

lanceolati vel quaque #. Spathae membranaceae, bivalves. Laminae faucis tri-
catuae Lamarckii l. c.

9x in gladiolus; foliis linearibus, floribus aequalibus spicati, petalis tribus
 Corollae campanulatae p. medio squamula erecta medio carinata Lamarckii. Dict. bot. 3. p. 341. illustr. 1.

Galeatus G. foliis linear-eriformibus falcatis, spathe longitudine tubi.

Jacqu. coll. 4. p. 167 ? 4c

~~Gladulus galeatus Thunb. Def. n. 1.~~

~~Habitatio Cap. 6. spec.~~

Scapus erectus, pithamou, compressus & ^{simplex} equinatus. Folia 3-4 in spico
linearia angusta, reflexo falcata glabra spico breviora. Flores
4-6 alterni. Spatha lanceolata membranacea tubo corollae
nonn. brev. longior. Lacina corollae n. 5. lacina superior
lanceata duo laterales oblonga 4. et inferiores lineares

Jacquini
X galeata
diversa a
mea

1000.

an spec: Antholyza.

60
roses

Gladiolus corollae tubus longissimus, laciniis superiore ovata, ^{foliis} ~~lanceolatis~~ - en
siformibus +

Gladiolus roseus; corollae muticae; tubo gracili longissimo; foliis ensiformi-
linearibus, glabris; caule ramoso; spathae valvula exteriori integerrima,
interiore apice bifida Gay. collect. Suppl. 22. Fe. rar. 2. t. 261.

Habitat in Cap. b. spec. 2

Gladiolus roseus; foliis

Habitat in Cap. b. spec.

Tubus subrotundus, magnitudine avellanae. folia longe vaginorosa,
radicalia plura, pedalia. Scapus foliis parum longior, erectus, culos
vel supra ramos declinatus, flexuosus, teris, gracilis, ramosus: Bractei
basi folio brevi suffulti. Flores secundi, bini vel terni, binaria-
tes, inodori. Spatha bivalvis, & cariosa, concava, acuminata, & trami-
nea. Corollae tubus gracilis, incurvus pallide roseus: limbi

laciniis rotundatae, patentes biniae, intus pallide roscae; supra
ma ovata, duplo latior, medio dorso purpurea; laterales duplo us-
que roscae medio coloratior; in feroces ma ad basin macula subin-
teiformis, rubra cum area longitudinali flavo, dua alia ad singularem
marginem Gay. l. c.

affinis videtur longifloro folia parum latera laciniis superior oblonga
add. desc. Gay.

Vanum dixer
quo videtur

Longiflorus 70 Gladiolus ~~coll~~ corollae tubo longissimo, limbo limbi alternae
angustioribus foliis linearibus ensiformibus

- 4^{ta} paniculata; caule paniculato, tubo floris longissimo, arcuatis De la

Roche Dis p. n. 14.

Ystia longiflora; polystachya, tubis corollarum filiformibus longissimis
mutantibus Berg. Cap. 7.

Gladiolus longiflorus; caule tereti, tubo longissimo, spathis foliisque
linearibus glabris Lin. Suppl. 96.

Gladiolus longiflorus; polystachya, tubo corollae longissimo, spathis
obtusis foliisque ensiformibus glabris Thunb. gladiol. n. 22. Redouté pl. tit.
t. 32.

Gladiolus longiflorus Jacq. Suppl. 24. ii. var. 2. t. 263

- β. Gladiolus longiflorus; corollae mutuae; tubo gracili, longissimo; foliis
angustis, ensiformibus, glabris; caule ramoso; spatha valvula exte-
riore tridentata, interiore bidentata Jacq. Suppl. 23. ii. var. 2.
t. 262.

Habitat in Cap. b. p. n. 4

Folia tria vel quatuor, basi breviora, angusta, linearia longe vaginantia
trinervia praeter nervi marginales. Scapus pedalis et ultra, interdum simplex,
saepius superne in peduncululos remotos, patenter divisus.

municata & spica municata: squarrosi variis mucronatis Mant. 3024 S.V
Mant. 302. Schreb. gram. 2. p. 69. & 34.

Habitat in India orientali

Panicum squarrosi spica geminis horizontalibus; involucri
florum squarrosi, culmi deiciambentibus Retz. off. 4. pag. 15. et
Laf. 5. tab. 1.

Habitat in India orientali

Culmi ramosi inferne purpurascens. Folia et rami ad
genicula coarctata. Eorum vaginæ villosa, undulata bipolli
clena. Spica terminalis plerumque geminis. Rachis utraque
membranacea. Glor. glomerulis congestis adgenicula rathiori
confertis, seplibus, squarrosi e squamis rigidis, glabris, striatis
patulis, mucronatis, ovatis, lanceolatis subulatisque mutis

hoc quom hoc qua
minus inter se dicitur
spicis cordur quor
spicis cordur quor

Ambr. Linnæus in purpureo oblongo culmo, reliquis aequalibus. An spicæ dicitur
hoc extensorum tenentibus, induratis bus in densitate: Conditio dicitur magis curvato
p. 69. tab. 2. 6. quæ dicitur in dicitur tenentibus, induratis bus in densitate: Conditio dicitur magis curvato
tab. 2. 6. sic synonyma dicitur pedum minimum vero a glabris longiflorum
to quæ eadem sit ac auctorum. Superius dicitur in dicitur tenentibus, induratis bus in densitate: Conditio dicitur magis curvato
sub nomine tenentibus longifloræ frequentior occurrit glabris longifloris, sed ut dicitur
Linn. in Linnæo tenentibus, oblongo culmo, reliquis aequalibus. An spicæ dicitur magis curvato
in dicitur in dicitur tenentibus, induratis bus in densitate: Conditio dicitur magis curvato

712
floribundus. II Gladulus corollae subringentis tubo longitudine spatulae, lim-
bacinis ^{en-} limbi lanceolatis, foliis lanceolato-ensiformibus
gladiolus floribundus; corolla subringente, foliis ensiformibus striatis,
glabris, scapo spicato toto florifero Gaerq. collect. 4. p. 162. Gen. rar. 2. 1. 2514
Habitat ad Cap. b. Spei 4

Folia plura, semipedalia, erecta, disticha, acuta, striata. Scapus
teres, longitudine fere foliorum, flexuosus, teres, superne saepe
geniculatus. Flores triunciales, sessiles, distantes, erecti. Spatha
bivalvis, convoluta. Valvula exterior bipollicaris, interior longior
viridis. Corolla tubus, subcylindricus, longitudine spatulae, albidus.
Limbi lacinis lanceolatis, pallide ochroleucis, linea utrinque media
purpurea. Stigmatis lacinis apice latescentes. Gaerq. l. c.

anceps 9 ^{part 1682} ~~anceps~~, spathe crispis Thurb. Gl. n. 17. 4 Jacq coll. n. 47 4
Gladolus anceps foliis ensiformibus undulatis caule ramoso an-
tice divaricato Lin. Suppl. pag. 447

+ ~~Frax Fabricii La Roche Des. p. 10.~~

4 Jacq ~~Lagenouza compressa Act. Tolos. 3. p. 79. V. 6.~~
coll. 4 ~~Habitat ad Pap. b. p. 6.~~

~~4 var. Caput fistulosum, glabrum, anceps, angulis denticulatis, ramosus foliis
nervi alterni simplices caule simpliciter. Folia amplexi cuiuslibet ^{longi} polli-
cari ^{distalia} ~~superius~~ ~~minora~~ ~~spatheae~~ ~~compressa~~ ~~causa~~
na, ~~obtusata~~ ~~pollicari~~ ~~superius~~ ~~minora~~ ~~spatheae~~ ~~compressa~~ ~~causa~~
latere inferiore ~~crispato~~ ~~dentato~~ ~~apice~~ ~~et~~ ~~canne~~ ~~purpureo~~
tanti. Spathe foliola exteriora foliis caulibus ~~simile~~ ~~usque~~
sepe pollicari, superius minores, obtusa. Corolla tubus filii
strobili superius amplexatus spathe multo longior, pallide
coerulea: hinc lacina coarctata~~

diffusio generis e Thurb. Des. de Gladolis petendo