

Grindia

1. paniculata G. foliis sparsis linearibus, floribus terminalibus aggregatis sesquipedibus tubo villosis.
2. cuneifolia G. foliis linearibus subulatis oppositis, ramuli axillaris, floribus solitariis.
3. racemosa G. foliis linearibus glabris, floribus axillaribus terminalibusque geminis sesquipedibus, ramis virgatis.
4. tomentosa G. foliis sparsis ovato-oblongis glabris, margine scabris.
5. sericea G. foliis ovatis tomentosis, floralibus quaternis, caule hispido, florum corolla setis octo.
6. imbricata G. foliis oblongis quadrifariam imbricatis sericeis floribus terminalibus in axillis foliorum villosis.
7. oppositifolia G. foliis oppositis lanceolatis glabris, floribus villosis.
8. simplex G. foliis omnibus linearibus acutis, floribus terminalibus sesquipedibus glabris.
9. biflora G. foliis sparsis linearibus subulatis acutis avenis convexis villosis, floribus aggregatis sesquipedibus terminalibus tubo glabro foliis brevioribus.
10. capitata G. foliis sparsis lanceolatis glabris, floribus capitatis bracteis obvallatis pedunculo nudo.

11. radicata G. foliis subulatis triquetris acutis, capitulis terminalibus sessilibus radicatae,
bracteis lanceolatis.
12. filamentosa G. foliis elliptice ovatis glaberrimis approximatis, floribus capitatis, flo-
rentis capellantiis.
13. Daphnifolia G. delandrou, foliis oblongis planis integerrimis, capitulis lateralibus
terminalibusque pedunculatis involucriatis.

Species indeterminata in herb. Acad.

Scabra G. foliis lanceolatis acutis scabri, floribus capitatis
Thunb. prodr. fl. cap. pag 76.
Habitat in Cap. b. p. c.

non vidi

argentea G. foliis sparsi obovatis tomentoso-argenteis, floribus capitatis
Thunb. prodr. pag. 76.
Habitat in Cap. b. pra.

punctata India

Habitata cap. 6 spec. herb. La Martii

Folia inferiora ovata, superiora lanceolata, subtus punctis albidis piliferis
coctemur glabra. Flores terminales, incani, tres

1 oppositifolia ^{oblongo-} ~~Quercus~~ ~~foliis~~ oppositis lanceolatis glabris floribus villosis
Thymelea foliis planis acutis, coma et floribus purpureis Burma

Afric: 137. t. 49. fig. 3.

Pasjenna fol. longata foliis ovatis glabris acutis, floribus obtusis Hen-Am.

ae: p. 4. p. 312. Spec: pl. pag 513.

Habitat in C. bono spec. ^{safo-purpurascens}

~~Frutescens glaber ramis proli feris tenens muniti. folia opposita erecta~~
~~vel ^{partibus} unguiculata, acuta, glaberrima ^{obtusae} ~~avenia~~. ~~lovis~~ ~~terminales~~ +~~

~~diffusa tomentosa foliis non oppositi~~

3 ve e quatuor

oppositifolia foliis oppositis lanceolatis tomentosis, floribus Verminea
ebur Thunb. prodr: pag. 76

~~diffusa tomentosa~~

longata g. foliis oppositis ovatis glabris, floribus terminalibus sub-
petalis Thunb. prodr. pag. 76.
Habitat in cap. b. spai.

Frutex glaber ramis fusco-purpureis, proli feris mucatis
Folii erecta vel patenta unguiculata, acuta, glaberrima
obliue venosa. Flores terminales 4.

an Lin. opposita folia et Burm. figura huius pedicel cum folio
in huius glabra

^{imbriata}
6 ~~Vomentosa~~ *Guidia foliis oblongis quadrifariam imbricatis penicis floribus terminatis*
libus in axillis foliorum *Lin. Suppl. p. 225. P.V. p. 373.*

Habitat in *Q. bono spei* ^{obscure truncati}
Tota planta alba *tomentosa*. *Folia opposita, sessilia, inaequaliter* ^{longa} *marginata,*
confertiora imbricata, superora petula. Flores in apicibus ramo- ^{piloso tomentoso}
rum interjectis foliis.
flores in hac tomento tenuiore in penicis pilosone,

~~quomodo differt a *Guidia* *speciosa* *Lin.*~~

polystachya 9

Berg-Cap

^{anceps}
9 ~~Crucifera~~ *Grindia* foliis lacini subulatis oppositis, ramulis caetera anicipitibus floribus
solitariis glabris
Habitat in Cap. b. p. 3

~~Trites glaber ramis proxis apice saepe ac ramuli e linea decurrente anicipi-
tis. ~~Stipula~~ ^{loba} opposita sesipila unguiculata acuta, plana copiosa. Flores
laterales et ex apice ramulorum solitarii sesipiles. Tubo glabro longitudinali
foliorum ^{senim} superne amplius tubo ⁴⁰ lobatus, tenuis laciniis ⁴⁰ ovatis, acutis. Petala
ovata. anthera 4 or inferiores in tubo, 4 or superiores stigma capitatum~~

Facies Clavariae an pinnifoliae in sepalis

Mulla
cuneata

Q. foliis cuneatis ^{pedunculis longioribus} ^{incaeni mucronati,} vellosis subtus ^{capitulis}
lateralibus pedunculatis

Habitat in Madagascar 3 Superne

Rami dichotomi mucronati ^{lomentis} ^{incaeni.} Folia par-
ta linguiculata obtusissima mucronata sessilia Pedun-
culi in dichotomia et est illic supremi folio breviori-
bus supra in 2-3 part. flori minoris uti capitulum quam
in precedente

Daphne gnid. ~~Daphne~~ folia p. lra.

1 *Laphnaefolia* *Quercus* *decandra* foliis oblongis planis integerrimis capitulis lateralibus
terminalibusque pedunculato involuato, flomb

Habitat in Madagascaria Thaurin Subdichotomi

varietas ¹ Rami teretes, ^{ad verticillata} ciliati apice vellosi pubescenti. Folia alternata canescenti
glabra ^{ad verticillata} utrinque pubescenti ^{in canis} ovata.

capitulis ^{ad verticillata} glabra utrinque pubescenti ^{in canis} ovata. Pedunculi
pubescenti ^{ad verticillata} glabra utrinque pubescenti ^{in canis} ovata. #
by pollicaresi vel ^{ad verticillata} glabra utrinque pubescenti ^{in canis} ovata. #
vel ^{ad verticillata} glabra utrinque pubescenti ^{in canis} ovata. #

by pollicaresi vel ^{ad verticillata} glabra utrinque pubescenti ^{in canis} ovata. #
vel ^{ad verticillata} glabra utrinque pubescenti ^{in canis} ovata. #
vel ^{ad verticillata} glabra utrinque pubescenti ^{in canis} ovata. #

decandra, foliis oblongis cuneatis planis, capitulis lateralibus
involucris pedunculatis pedunculis brevioribus

varietas B. Can.

glabra est *Davis* madagascariensis La Mart. herb
hirsuta pubescens La Mart.

2 radiata Inidia foliis subulatis triquetris acutis, capitulis terminalibus sessilibus
radiatis, bracteis lanceolatis her Mart 67: #13, urm. prædr. 12

Habitat in Cap. bono spei;

Frutex scaber prolifer, folia parva, mucronata, glabra. Capitula ter-
minantia ramos, radiata bracteis lanceolatis, foliis latioribus. Flores
extus puberuli. Calycis limbus intus glaber, longitudine tubi. Petala qua-
tuor calycis minoris, pubesca. Stamina 4 extesta, 4 in faucibus. her

l. c.

Rame in ani fomentis, junca. Folia linearia, penninerviulana, glabra
Flores terminalis 4-5 in capitulis ciliatis, foliis involucri, sessilibus; semi-
unquivalves villosi herb. frus

non vidi

omittitur in Thurb. prodr

12 Filamentosa Gnida foliis ellipticis ovatis glaberrimis approximatis, floribus
capitatis filamentis capillatibus. En. Suppl. p. 222 224. + P.V. p. 373.
B — ^{et nunc. Hortic.} Habitat in capite bono preti

Erigit glaberrimus cicatricibus foliorum tuberculatis ramuli per
perascentes. ^{longe tuberosi et illati} Solia alterna ^{usque ulano et hauri} ^{multiflorum h. d. ellipticum} ^{integerrima} arenis capite
la florum solitaria. Involucrum glabrum. Corolla villosa: lim-
bo quadrifido. Laevis lanceolata. Filamenta capillacea:
quatuor longitudine corollae: quatuor ad faucem breviora
Sunt

Lactuca bursifolia La Mart. herb

~~Gnida biflora - penifida Suppl. p. 222 224. uterque et a penifida non speciem~~
herb. h. h. penifida

11 capitata *Gnidia* foliis parvis lanceolatis glabris, floribus capitatis bracteis obrallatis,
pedunculo nudo *Len. Suppl. p. 224. L.V. p. 378.*

Habitat in Cap. b. p. i.

Caulis fruticosus. rami teretibus erectis, purpureo-fertibus. Folia angustiora
remota, lanis. Flores aggregati terminalis, pedunculatus. Involucrum octo
phyllum lanceolatum, aequale, lobis amplexibus brevibus. Lobuli pedicem
lobis, pubescentes cani. Limbus brevis quinque partitus. Stamina 10. 8 eos
passeris similes. forte *Dais* *Len. L. C.*

Dais unifolia *Ca. Mart. herb*

modi

non

4 Tomentosa Gnidia foliis parvis ovato-oblongis glabris, margine scabris Lin. l. V.

pag 372. *

Gnidia pubescens Berg cap. 124.

Habitatio. Cap. b. Spei t.

Ramulorum apices lanati uti congenere. Folia suprema subpubescentia. Flores tomentosi, capitati, terminalis. Petala parva emar
cynata. Lin. ^{semiquinaria} ^{ventriculi} ^{conules}

Rami cortice fusco subpulso. Folia petiolo brevissimo erecta & denticulis
margine scabra, f. d. s. supra scabra s. p. scabriuscula nitida
subtus obpura nervos folia floralia magis lanata
ta

oculo armato
videndis f.

floribus tomentosis

villosi canis

umbicata

microphylla G. foliis linearibus glabris ramulis floribusque terminalibus
pilosiusculis
Habundat Cap. 6. ^{Dr Murmann} ~~preit~~ interna glabris ^{et superne}
Fuit cortex cinereus rami proliferi pilosiusculi ~~apice~~
iniani ^{apice} ~~marginato~~ ^{marginato}. folia ovata sessilia ^{grassa} glabra pubes
carinata acutiuscula facie foliorum ^{qu. raiemope.}
flores plures 6-7 aggregati unguiculati ~~et~~ ^{et} pilis rannonibus
receptaculum pilis albis

gnidua carinata foliis lanceolatis obtusis cinnatis glabris, floribus capiti
tatis Thunb. prodr. pag. 76

An Daphne capitata

3 racemosa Gnidi foliis ^{obtusis} lineariibus glabris, floribus axillaribus terminalibus
que sessilibus ^{sub} geminis ^{rami} ramis ^{erigatis}

Habitat in Cap. bono spei m. f. Dr. Dahl.

Similis G. simplicis foliis et facie at flores folio breviores laterales
terminalesque folia ^{fracta}

Rami mucronati glabri teneri. Folia seminaequivalenti proplea al
terna. flores extra villosi cerneri 2-3. axillares et terminales

Gnidi racemosa foliis obovato-lanceolatis latius glabris, floribus racemosis
axillaribus Thunb. prod. fl. cap. pag. 76

10 biflora ~~Gn.~~ ~~foliis~~ ~~gracis~~ ~~linearibus~~ ~~subulatis~~ ~~planis~~ ~~convergentibus~~
lobis. ~~tubo~~ ~~glabro~~, ~~foliis~~ ~~brevioribus~~
acutis ~~avenis~~, floribus aggregatis ~~sessilibus~~ ~~termina~~

Gn. ~~perfolia~~ foliis linearibus subulatis ~~superioribus~~ planis acutis, floribus binis
axillaribus. Len. Suppl. pag. 225* S.V. pag.

Habitat in Cap. bona ~~part~~ #

Caulis ~~ter~~ Rami ~~ter~~ nec tomentosi. Folio acuta non pungentia, subtus convexa, lacinia
patula. Limbus acutus. Tubo duplo longiore. Len

P. Rami ~~gracis~~ ~~ramulosis~~ ~~tetragonis~~ ~~apice~~ ~~ancipites~~ ~~purpurascens~~
foliis ~~sessilibus~~ ~~oppositis~~ ~~vel~~ ~~angustis~~ ~~linearibus~~ ~~patula~~. ~~flores~~
axillares ~~maiores~~ ~~et~~ ~~caeteros~~ ~~quam~~ ~~in~~ ~~reliquis~~ ~~tubis~~ ~~semper~~
ampliatus. Lacinia ~~linearibus~~ ~~patula~~ ~~vel~~ ~~lobis~~.

biflora Gn. foliis glabra ~~ramis~~ foliis linearibus subulatis planis
acutis, floribus binis axillaribus, ramis ~~superne~~ ~~ancipites~~
juniperifolia La Martini

Specimen herb. an potius simplex

penifolia Guedes

Simplex Gnida foliis omnibus linearibus acutis, floribus terminalibus sessilibus ^{glabris} Mart. 674
L. V. p. 372.

Gnida viridis Berg.

Thymelaea folio passerina Boeyn. Cent. 10. t. 6

Habitat in Cap. b. sp. 17

Caules semipedales inaequales, exasperati tuberculis a caespitibus ^{teretibus}. Folia
parva lineana, ac. erecta teria. Flores terminales, nudi ex apice rami
relloso, sessiles, lutei calyce, petalis et antheris. Folia floralia reliquis
non latiora. Petala oblonga, acuminata. Stamina 4 in faucibus & supra
cautem. Lin. l. c.

3 species
differt.

~~Rami proliferi remote mucronati. Folia ^{teris} glabra vix conseruata
laxa ~~laxa~~ apice et basi rami subangustata in ramulis confertis
tubus basi striatus, pubescens foliis longior latioribus oblongis
obtusis~~

pubulata La Mart

vel uniperifolia La Mart herb

2 *punifolia* Gnida foliis ^{terminalibus} sparsis linearibus floribus ^{aggregatis} sessilibus, tubo villoso
Rapunuculus foliis nervosis linearibus floribus argenteis non galeatis Burm afr.
112. tab. 41. fig. 3.

Valerianorides ^{Aethiopia} ^{Indiensi}

Gnida *punifolia*: foliis sparsis linearibus, floralibus verticillatis Mant 375

P. V. pag. 372. Berg. cap. 122.

Habitat in Cap. b. p. et proliferi exet. gemini vel terni

semunquiculana ^{vel paululum} ^{crassifolia} ^{subtus} ^{conserata} ^{marginata} ^{angustiloba} ^{floralia} ^{triplo} ^{latiora}
Rami vertes glabri, et apertis e cap. solonum. Folia ^{petiolo} ^{bravi} ^{frequentig}
flos ^{lacina} ^{tubis} ^{calycis} ^{villofis} ^{pollinaris} ^{lacinis} ^{obtus} ^{obtusis} ^{sementosis} ^{petalo}
lanceolata ^{glabra} ^{incana} ^{exte}