

Gomphocia

1. longata G. foliis lanceolatis obtusissimis integerrimis emarginatis, panicula terminali
2. angustifolia G. foliis lanceolatis extrorsum serratis, panicula terminali, petalis calyce longioribus
3. nitida G. foliis ovato-lanceolatis acuminatis serratis, paniculis terminalibus, calycibus corolla aequalibus
4. papotaputa G. petalis calycibus triplo majoribus, dupli receptaculo

Sandalli

Corcia huc pertinet secund. Muell.

7 8 9 10 11 12 1
guianensis Gomphia, foliis ellipticis nitidis coriaceis extrossum ternulatis, racemi
terminalibus compositis, calycibus corollam subquartibus

Gomphia guianensis; foliis oblongo-ovatis; subintegris, coriaceis;
panicula terminali Ad. B. h. n. P. 1. p. 168

~~Habitat Cayenne Richard 5~~

Othna guianensis; floribus pentalobis, desandri; foliis ovato-ob
longis, subserratis, panicula terminali, La Mart. Dict. Encycl.
4. p. 511.

Ourotea guianensis Clubl. fl. Guian. p. 397. T. 152

~~Habitat Cayenne Richard 5~~

Stipiculus 4-9 pedalis, glaber. Rami cinerascetes, varicosi, laxes.

Folia breviter petiolata, alterna, epithymosa sulcata, obtusa, subcoriacea, utrinque nitida, cavia, subtus tenui sive obscure venosa, extrossum obsolete serrata. Racemi erecti. Pedicelli bi-triflori. Calycis foliis

inferne ramo
uno alteriore

la linear-lanceolata. Petala calycis parum longiora, obovata, flava, unguicula breviter. anthera apice flavo. Semina 1-2. nigricantia, oblonga magnitudine pisci.

Topobea Melastoma

Habitat in Guiana Herb. Fus.

Rami crassitie pennis cœquea, tetragonis, quibus, apice pulverulentis. Folia per
foliata, opposita, ^{ovata} ^{oblonga} ^{acuminata} ^{vel ovata} ^{apice} ^{vegetis} ^{apice}
approximate nervosa, ^{stensa} ^{venosa}, in ^{eyemi} ^{ma}, pollicaria. Petoli:
les qui pollicares

unde & abruscul

supra oculo armato adspersa punctis numerosis semis albis

parositica

+

L. n

Squamis *Gomphia* foli⁴ lanceolatis oblongis nitid⁵is remotis serratis; racemis terminali⁶
bus subsimplicibus.⁷ ⁸
Habitat ad Terra Leonam. ¶

Rami teretes, glabri. Folia subsimplex, fere quadri-polluciana, lanceolata
oblonga, venis sine venosa, nitida, remote serrata, utrinque acuta. Racemus
tri-pollucianus, terminalis, erectus, basi ramo brevi florifero. pedicelli frequentes,
sparsi, saepe geminati, post caput floris persistentes ut in *Ochna squarrosa*. Squa-
mae in axillis et folia parva acuta.

Cinnamomi ⁷ ⁸ ⁹ ¹¹ ¹¹ ¹² ^{Sub} ¹³
folis Gomphia folus lato lanceolatis nitidis attenuatis remote ~~obtusis~~ serratis.
~~Paniculatis~~ lateralibus terminalibusque, calycibus corolla aequalibus.
Bohoroe Zeylan. Herm. herb.

Gopitabita Cinnamomi folis flexibus pinnatis Burm. Thesur. zeyl. pag. 123. tab. 56

Habitat in Zeylona herb. Hermanni 5

Folia lato lanceolata, nitida, tenuissime venosa, attenuata, exteriorum obscure
remoteque serrata, 3-4 pollicaria. Panicula lateralis et terminalis. Flores parvi.

Petalis longis in calycis. Calycis foliola ovata. Fructus obovatus, latere interiori ac-
uatis, base subcompressi.

Dwarfia omnino ab ocina squamosa

lanceolata *Gomphia* foliis lanceolatis utraque ⁹ attenuatis obscure serratis, racemis
¹⁰ ¹¹ ¹²
¹
Tex municipalibus compositis, petalis linearibus calyce aequalibus
Habitat in Brasilia Herb. Aussisi

Rami teretes, cortice cinereo, potentissimi, rami flexi, folia petiolata, tripollia
na, trinque attenuata, acuta, glaberrima, nitidissima, ex tropico obscure
serrata, basi versus integerrima, vix venosa. Racemi terminales ramorum,
compositi. Pedunculi purpurei: pedicelli solitarii, subtriflori. Flores parvi. Ca
lycis foliola lanceolata. Petala linearia, obtusa, calyce vix longiora. Sta
mina 10.

longifolia Gomphia ⁴foliis ⁵subcordato-oblongis ⁶elongatis ⁷integerrimis, racemis ⁸ser-

^{omnibus} ⁱⁿ ^{pleuribus} ~~Habitat in Guadeloupe Richard herb. Musc. & herb. Gussonei~~

Ramis teretibus, uti omnes reliqua partes glaberrimis, nitidissimis. Folia brevissimis
petiolata, pedalia, basi ovata, subcordata, oblonga, attenuata, tenuissime acuta

^{multis nervis} ~~ser~~, subvenia, coriacea, integerrima. Petioli unguiculares,
crassitie penna anserina, rugosa. Racemi terminales, quinque pollicares:
pedicelli sparsi, saepe geminati, unguiculares. Calycis linear-lanceo-
lato.

Othna longifolia; floris pent-petalis, foliis cordato-lanceolatis, em-
gus geminis, integris La Mart. Dict. Encycl. 4. p. 511

cardiosperma Gomphia.

Ochna cardiosperma: floribus terminalibus; lacinis bilobis; foliis ovato
pinnatis La Mart. Dict. Encycl. 4. p. 511.

Habitat Capeno la Mart. 5.

guianensis Gronchia

Act. fort. herb. nat.

Habitat Cayenne herb. Nutt. Ind.

Stem culm 4-5 pedalis, glaber. Folia alterna, ^{inota} breviter petiolata, spithamea
et ultra, elliptica, obtusa, obsolete venulosa, ~~vulgaris~~, membranacea, subtus
magis nitida, obscure serrata. Panicula terminalis, erecta. Calycis
foliola lanceolata. Petala calyce parva longiora, oblonga, unguiculis
brevi. Semina 1-4 nigricantia, oblonga, magnitudine P. si.

~~acuminata~~
~~nitida~~ Gomphia foliis ovato-lanceolatis acuminatis serratis, paniculis ter-
minatis, calycibus corollae aequalibus aequalibus
Oikna nitida: calycibus corollae aequalibus, dno p[er]ceptaculo super
imp[er]f[ect]is, foliis ellipticis serratis Sw. prodr. p. 67.

Habitat in Darén dedit Dr. Benth.

Brachi alterni, superne flexuosi, cortice fusco, teretes, glabri, utriusque
qua planta.

Folia brevissime petiolata, alterna, bis. tripollicaria, pollicis binos lata, conia-
cea, xx venosa, coriacea, serraturā mucronatis, basi integerrima.

Panicula terminalis, stricta, patens, palman vel ultra: Pedunculi
ut pedicelli alterni, pedi post casum florum e pedicellis perfectis
subus squarosa. Flores ante evolutionem ovati, obtusi

Calycis & lobula longitudine petalorum, lanceolata.

Petal flava.

Filamenta tubercula decem, parva: anthera longitudine corollae
subulata.

germen pentagonum: stylis longitudine antherarum. stigma
acutum.

decumbens
51-52.

Euphorbia umbellata quinquefida. Dichotoma involuelli ovatis cuspidatis
foliis linear-lanceolatis inferioribus obcordatis

Habitat inter segetes Tineti

Radix ^{annua} ~~perennis~~ ^{semperfolis}
Caulis herbaceus basi ramosus ramis decumbentibus teretibus simplicibus

foliis glabris ^{alternis} ~~approximatis~~ ^{glabris} ~~integerrimis~~,
Tolia copiosa, sparsa, sessilia: ~~inferiora~~ ^{inferiora} minor, cuneata, amarginata
mucronata, superiora ^{magis longe} sessili majora ^{magis usque usque} sessili basi attenuata ^{minora}

Pedunculi ex omnibus axillis ^{magis longe} ~~superioribus~~ bifidis. Umbella termi-
nalis quinque ^{septa} ~~septa~~ radiata bis bifida. Involuella ^{septa} ~~septa~~ penta vel
6 phylla ^{septa} ~~septa~~ foliolis obovatis mucronatis cuspidatis. Involuellis
ovatis, altero latere angustiora cuspidata. Flores solitarii parvi
Capsula glabra ^{septa} ~~septa~~ magnitudine comarum ^{septa} ~~septa~~

affinis esula et Taurinensi Allioni ~~du~~ qui differunt caule erecto
foliis utroque acutis

nomen triviale mutandum

Tabotopita g. petalis calycis triplo majoribus, drupis receptaculo immersis
L. prodr. p. 67.

Tabotopita Marigr. Braf. 101

Tabotopita pyramidato flore luteo, fructu rubro Pl. Amer

42. c. 153.

Ocena Tabotopita racemis terminalibus L. S. V. pag

Habitat in ~~India~~ ~~o~~ amer. meridionali &

arguta *Gomphia* foliis ⁷ lanceolato ellipticis ⁹ argute serratis ¹⁰ sparsis, ¹¹ racemis ¹²
minutibus ³ subcompositis, ⁴ calycibus ⁵ corolla ⁶ aequalibus

Habitat in Guinea Geit. 3

Rami teretes, laxes, cortice fusco purpurascente, uti reliqua partes
planta, glabri. Solcia breviter petiolata, distantia, spitha mea stri-
que acuta, frequentissime serrata, serraturis approximatis acutissimis ra-
in una aetate reliquis minor, basin versus spatio pollicari integerrima
senosa. Racemus palmatis multifloris, pedunculi palmatis sparsi, 6-6.
pedicelli saepe gemini unguiculari. Calycis foliola lineari lanceolata
semiunguicularia. Petala obovata, longitudine calycis

africana *Bygonia*

Habitat in herb. Tusj.

Folia pennata cum pari, triplica: foliola brevia petiolata, opposita, elliptica glabra, tripollicaria, nitida, glauca, subtus venosa, utrinque glabra, acumine brevi terminata, est, utrumque dente uno altero ve. Calyx glaber, semipollicaris: laciniis ovatis, acutis, coriaceis. Flores natescentes, rari amosi

~~Gomphia Schreb. Gen. pl. Gomphia~~ Gomphia Schreb. Gen. pl. pag 291
angustifolia ~~Alb. Schreb.~~ foliis lanceolatis, et ~~trorsum serratis~~, panicula
terminali, petalis calycis longioribus.

Habitat in India orientali. $\frac{1}{2}$ König $\frac{1}{2}$

Glaberrima saeva

Folia albena, sessilia, bipolliciana, a basi ad medium integer
nna, utraque acuta, membranacea, venis ^{subtilissimis} ~~multis~~ ^{per}
reticulata.

Panicula terminalis, par bipolliciana

Floris globosis, ante ²explicationem

lobis ²Calycis subrotunda.

Petala quinque

Stamina ¹⁶ decem.

nitida g. foliis lato-lanceolatis utraque acutis serrulatis, raris
subsimplexibus; calycibus corollam aequantibus

~~Ochna nitida foliis ovato lanceolatis acuminatis serratis, panicu-
lis terminalibus; calycibus corolla aequatis~~

Ochna nitida calycibus corollae aequalibus, prope receptaculo
superimpositis foliis ellipticis serratis Sw. prodr. p. 67.

Habitat in Cambes ^{alterni remoti}

Rami cortice cinereo ex hyllis ramuli glabri breves foliferi
vel floriferi apice purpurascens. Folio ad apicem ramo
suo membro naxea tenuissima venosa utraque nitida basi
integerrima ^{bis} tripolliciana ^{racemi} in apice ramo
suo basi bipollicares basi ramo uno altero a capite pedicel
longi squarosi, sparsi pedicelli unguiculares glabri. Foliole
calycis lanceolata

~~Spinosum Zygophyllum foliis conjugatis sessilibus: foliolis linearibus carnosis supra planis,
caule fruticoso L. S. V. p. 400 *~~

~~Tabago tenuifolia spinosa, fructu rotundo Burm. afr. 5. t. 2. pag. 2.~~

~~Habitat in Othopia 5 L. L. f. 101~~

~~caulis erectus: Ramis alternis longitudine plantae. Stipula verticillata quatuor, fu-
bulata, breviflora, rigidula. folia quaternaria, linearia, carnosae, supra plana,
subtus cretiuscula, acuta. Corolla nutantes luteo L. L. c.~~

lanceolata Gomphia integerrima
Ochna foliis lanceolatis obtusis pinnis emarginatis, nervis
verminali.

Habitat in India orientali Hong

et Gaberrima, Rami alterni cortice cinereo

Petala brevissima petiolata, sparsa, basi acuta, apice obtusis-
sima, emarginata, bipolluciana, nitida, areolis subcircularibus,
unquem lata

Petiolus brevis semus ad insertionem intrasatus.

Panicula terminalis, ^{facie Ochn. squarrosa} ~~pedunculata~~ patentissima, rami reflexi,
purpurascens, et facie Ochn. squarrosa

Soliola Calycina lanceolata.

Corolla spatulata longitudine calycis
filamina 10.

Basis receptaculo superimposita.

Secundum
v. 54
p. 4
Corolla

Character generici in Ochna in edit. Generum Plant. Schimper
tam in squarrosa quam in
lat Petala Corolla; ^{v. 54} Petala 10 ad esp. ^{v. 54} et esto certum
affert est, in ~~specimen~~ non in unio.
sed plurimis specimenibus vidi. Petala decidua. ~~calyx~~
~~calyx~~ calyx persistit

Dichonis