


Gymnanthes

- ♂ Amentum nudum. Stamina pedicelli tripartiti l. 3 thotomae, antheriferi
♀ Cementum l. germen pedicellatum. Stylus trifidus. Capsula trilocula

1. lucida g. monica, staminibus trichotomis, femineis solitariis pedicellatis
2. elliptica g. dioica, staminibus tripartitis, femineis solitariis amentaceis



a amantum cum foemina ad basin magnetis.
 naturalis b. Hamnia aucta. i. Pappula
 d. eadem trans respectu e. f. eadem cum anillo
 f. eadem absque anillo



Lucida Gymnanthes. monoxia, staminibus trichotomis, femineis solitariis, pedicella
his Sw. prod. pag. 96.

Habitat in Montserrat In Ryan.

Rami teretes, alterni, remote ciliati, crassitie filii impositi, glabri, cor
que cinereo.

Folia versus apices ramorum, distantia, petiolata, alterna, bi- tri polli
cario, elliptico-lanceolata, inferne versus partem angustata, basi
acuta, apice obtusa, saepe emarginata, remote serrulata, utrinque
glaberrima, tenuispmi venosa, supra nitida.

Petioles semiunguiculari, purpureo fientes.

Amenta axillaria, solitaria, subsessilia, bipollicaria. Flos femineus
inferne ad basin eodem amento, pedunculatus solitarius, pedunculo
unguiculari

Germe oblongum, glabrum. Stigma Stylus vix ullus. Stigmata tria reflexa
lancearia.

♂ Stamina distantia receptaculo subovato, insidentis. filamenta stria
bifida ~~et~~ ^{supra} quadrifida, basi saepe lanceolata. Anthe
ra oblonga

Capsula triocula, trilocularis; loculis monoplocis. Semina oblonga
basi articulo impleto

a amentum b. Stamina receptaculo infidenter magnitudine aucta. c) capsula
magnitudine naturali d) transversim secta, e) semen cum anillo f) semen

opaca Gymnanthes foliis oblongis mucronatis integerrimis

Habitat in Bonaria Commesor herb Jusf.

Rami teretes, glabri, cortice cinereo. Folia petiolata, pollicaria, obtusiuscula,
basia acuta, axillaria. Stipula setacea. Pedunculus terminalis, flos foemineus
inferne pedicellatus; masculi dichotome ramosi.

Gymnanthes

Habitat in America herb. Desfont.

Folia lanceolato-oblonga, utrinque acutiuscula. Pedunculi inferne, uti & folia junceorum hirsuti. Pedunculi superne villosi. Flores foeminei inferne, 1-2 alterni cum masculis mixti, subsessiles.

1. *Cucida G. monovica*, staminibus trichotomis; femineis solitariis pedicella
tis. W. prodr. pag. 96.

Habitat in Hispaniola;

Rami teretes cortice cinereo ramuli ter verticillati (an semper.) Folia
petiolatis alterna lanceolato-elliptica, in sedum obovata, obtu-
re superne subpirata inferne integerrima, glabra venosa supra
nubida bipollicaria. Amenta Petiolis purpurascens unguicu-
lari. Amenta ^{ma} axillaria sessile solitaria in divifera ramo-
rum laxe ternis folio brevioris

2 elliptica G. dioica staminibus tripartitis, femineis amentaceis Sw. prodr
nuz. 96.
Habitat in America

non vidi