

# Hedera

1. helix H. foliis ovatis lobatisque
  2. pendula H. foliis ovato-lanceolatis integris; pedunculis longissimis pendulis
  3. nutans H. foliis ellipticis zoniatis, umbellis nutantibus hemisphaericis
  4. capitellata H. foliis ellipticis integris, racemis compositis terminalibus; floribus sp.  
libus capitellatis.
- The rebintinaea

capitata Hedera foliis ellipticis integris, ramis compositis terminalibus,  
floribus sessilibus capitellatis Smith. ii. part. 4. t. 4. \*

Aralia capitata Jacq. Amer. Og. t. 61.

Aralia capitata: arboracea, foliis simplicibus ellipticis integerrimis, paniculis  
terminalibus, floribus sessilibus capitatis Sw. prodr. pag. 55

Habitat in Martinica et Jamaica sylvis

pendula # foliis ovato-lanceolatis integris, pedunculis longissimis pendulis lvs.

prodr. p. 51.

Habitat in Jamaica

non vidi

nutans # Folis alleptis conacis, umbellis nutantibus hemisphaericis Sw.

prodr. p. 51.

Habitat in Jamaica.

non vidi

Hedera

7 terebinthinaea Hedera foliis septenatis ellipticis integerrimis  
Arbor Zeylanica & ita dicta, resinam terebinthinae similem fundens  
Herm. mus. zeyl. pag. 50. Burm. Thesaur. zeyl. pag. 28.  
Itawol. Lin. fl. zeyl. n. 624.

Habitat in sylvis et nemorosis zeylona König 5  
~~folia glabra~~  
Rami teretes, glabri, subherbacei, crassitie penna Cygni

Folia petiolata, septena. <sup>lateralia ~~antip.~~ minoras</sup> soligla umbellata, petiolata, elliptica  
3-4 polliciana, elliptica, obtusa vel acuta, interdum acumi  
ne basi terminata, integerrima, glabra, subvenosa.

Petioles puthamosi, teretes, basi dilatati amplexantibus: pariter  
et in medio bipollicares, laterales minores

Racemus terminalis, ad latus rami foliosi, compositus. Pedun-  
culus universalis tripollicaris, crassitie penna asperna,

Pedunculi pariter alter quini, alterni, ultimi duo  
conjugati, semipedales, patentes. Flores umbellati: umbella  
pedunculata, alterna 16-17 in singulo ~~radice~~ <sup>pedunculo pariter</sup>, pedunculus  
unquiculus ~~in~~ <sup>in</sup> umbella 6 vel 8 pedunculato  
radice umbella 6. 8 semiquiculares, glabri ~~ut~~



*Helix* G. foliis ovatis lobatisque Lin. S. V

- Hedera arborea* Bauh. pin. 305  
β. *Hedera poetica* Bauh. pin. 305  
γ. *Hedera major* *fertili* Bauh. pin. 305  
δ. *Hedera humi* *repens* Bauh. pin. 305

Habitat in Europa arboribus putrescentibus, inque sepibus & Rami teretes, hinc inde inferne fibillis radiceantibus emittentes glabri. Folia in *fertili* cordato tri vel quinquangulata in *fertili* ovato <sup>subtriangulari</sup> ~~rhomboides~~ attenuata, glaberrima supra nitida a subcoriacea inteserrima venosa. Racemi terminalis ramorum tomentosi. Umbella racemosi alterni patenti sumi

*Pseudophyllum* A. arborens, petiolis pedatis, foliis umbellatis terminalibus  
petiolatis, foliolis aliis centralibus, *Species Conyosporis Sw. prod. p. 58.*

*Pseudophyllum* Brown. *Jam. 195. t. 19. fig. 1. 2.*

Habitat in Jamaica

Rami glabri folia petiolata. foliola circiter 10 petiolata  
canaliculata elliptica bipollicaria integerrima utrinque glabra  
puberula inaequalia rostrata pedales pedicelli sparsi

~~non in di~~

Species heder. uti et *Asalea capitata* secund. Dr. Schumier