

Hedwigia

1. balsamifera H.
- 2 - varia
- 3 - pubescens

varia et pubescens quae sub nomine Hedwigia inter Am. et in Systemate

balsamifera Hedwigia

altissima Lucia. Publ. fl. Gu. pag. 342 Tab. 122.

Habitat in Monserrat de Ryan, in insula St. Crucis legit West
de Rooy

Rami teretes, glabri.

alterna vel

Folia in apice ramorum, petiolata, pedalis, pennata cum impari,
trijuga. Foliola breviter petiolata, opposita, quinque pollicana, inter non
minora, sequi pollicem lato, oblongo-lanceolata, acuminata, acuta,
integerrima, venosa, utriusque glabra.

Petioli triquetri, glabri, purpurascenti; supra plani ^{sive laevi} ~~absolite striati~~

subtus striati: parthales femiungui ulari.

Panicula paniculi ^{ramum} infra folia, ^{pedunculati} approximati, alterni, glabri, striato-angu-
lati, purpurascentes, longitudine fere foliorum: parthales oppositi
pedicelli ultimi triflori; flores pedicellati.

Calyx monophyllus cyathiformis, parvus, quadridentatus, glaber.

Corolla tetrapetala, calycis duplo longior: Petalae oblongae, obtusae.

Stamina filamenta vix ulla, receptaculo inserta. Anthera oblonga,
parva, corolla brevior.

Germen pyramidatum, longitudine staminum, glabrum, basi octo crena-
tatum. Stylus nullus. Stigma sesple acutum.

(Corolla primum monopetala videtur, demum vero in quinque petala distincta
resolvitur, et stamina omnino receptaculo inserta sunt. Criblatis *Frax* huius
pertinere videtur et ^{arbor} planta jam descripta, quae non differt videtur ab *Frax*
altissima, obstat vero Capsula bacata *Frax*, de qua tacet Sw. asperit vero
Dr. Ryan Capsulam huius materiam resinofam continere, ideoque *Hedwigia*
genere ab *Frax* non differt

1 Balsamifera Hedwigia Schw. prodr. p. 62.

Bos Cochon Nicof. h. St. Doming. 169.

Habitat in Hispaniola

Rami teretes ~~canaliculati~~ ^{in minutis cuneis adpressi} punctati. Folia petiolata alterna, impari
pinnata, sempervivalia, 3juga. Foliola ^{opposita} ~~ovata~~ petiolata, ovata, acu-
minata glabra, venosa, membranacea trijocollicaria, undulata, glabra.
Petiolus supra planus, subtus convexus. Panicula terminalis magna
diffusa. Pedunculis alternis spithameis compressis. Pedicelli ultimi
oppositi. Calyx campanulatus parvis glaber quadridentatus. Corolla
tubulosa, calyce duplo longior ^{duplo brevior} ~~longior~~ ^{duplo brevior} ~~longior~~ laciniis oblongis obtusis. Stamina
tantum 7 ^{inveni} ~~inveni~~ ^{filamenta brevia, recaptatulo infixa?} ~~recaptatulo infixa?~~ anthera oblonga germi-
ni adpresso. Germen ovatum acutum falcatum longitudine staminum. Stylus
nullus

Ficus altissima

1 actosima *Tecia* foliis lanceolato-ellipticis, racemis axillaribus
Tecia actosima; foliis amplissimis, impari pinnatis; fructu racemoso
Habitat in Sylvis guianis

Arbor actosima. Folio alterna, impari pinnata bipuga pedali
et ultra. Foliole opposita brevissime petiolata semi pedalia acumi
ne semipollicari terminata, integerrima, venosa, glabra
Capsula subovata magna, bi^{3/4} valvis Valvulis carnosiss
extus verrucosis nupersentibus. Semina ospecula duo, 3-4.
nigra pulpa alba dulci obvoluta