

Helminthia

Calyx duplex. Semina obovata, transversim striata. Receptaculum tuberculatum

echinoides 1/2

aculeata 2/4

Sprengeana 3/4.

Pappus striatus

Echinosides P. peranthus extenonibus preta phyllis interiore aristato majoribus Lin. Spec.
pl. pag. 1114x. S. v.

Buglossum echinoides luteum hieracis cognatum Lob. ii. 577.

Hieracium echinoides capitulis carni benedicti Bauh. pin. 120

ps. *Hieracium* capituli carni benedicti Herm. parad 105. t. 105.

Habitat in Anglia, Gallia, Italia sylvis caducis, argenteis

Caulis erectus, pubescentibus sub ramosis pilis rigidis adpressis. Folia lanceolata
inferiora basi in petiolum amplectuntur, caulibus attenuata dentata, ramae
superiora basi aequalia amplectuntur, in tegmina omnia pilis rigidis adpressis
pedunculis ^{ad marginem} terminalibus ramosi. Radicelli hepatici. Calyx exteriore interiore
major folioli ovatis acuminatis, setula terminatis, margine stimulis
rigidis ciliatis. Interiore octangulatis, folioli totidem carinatis, infra
apicem aristatis, his exteriore foliola totidem minuta locis. Semina
munita. Pappus stipitatus plumosus

aculeata Picus calycibus laxis, ^{Picus} foliis dentatis pedunculis subaphyllis, foliis ellipti-
co-lanceolatis cauleque frugoso aculeatis foliis sinuatis.

Arenarium ethioides, capitulis cardui Benedicti, majus, arabicum, Journes.

Inst. pag. 470

Legi in montosis ~~montibus~~ ^{montibus} ad Zouan urbe Tunetana haud frequens
Caulis pedalis erectus, ^{cras} simplex, striatus, adpressus spinulis totus ~~asperus~~
spinulis parvis rigidis pungentibus.

Folia radicalia plurima radicalia, oblonga, in petiolum deprentia,
bipollicaria et ultra, obtusa, dentato-subsinuata, margini et
in pagina utraque adpressis aculeis parvis, numerosi inaequalibus
albis adpressa. Caulis paucus, semiamplexicaulis, lanceo-
latus, remotus, ~~non~~ dentatus, aculeis aculeatque inferiori,
sesquipollicaria, sensim minora, rudimenta ramulorum in axillis.

Pedunculi versus summitatem, ramosi, hispidi, foliolo uno alterove
spinuloso ciliato.

Foliola calycina in ^{linear-} calyx calyculatus: foliola lanceolata, dorso spinuloso-ciliata
Pappus plumosus. Coma sub¹strata & transversum sub²strata. Pappus pelosus
(Differt a Pico ethioides ab Pentanthia pentaphylli ex tenore
vate

Picus calycibus simplicibus perianthio
maximo distalibus, foliis profunde sinu-
tis honor. fl. quadr. n. 243

~~Pennis calycibus simplicibus perianthis maximis obvallatis, foliis profunde
sinuatis Gron: fl. Orient pag. 100~~

~~Differt ab Echioides ab sentio perianthii reflexionis pentaphylli~~

¹¹⁷¹
japonica P. *penanthus* *hispida*, foliis *dentatis*, pedunculis *subaphyllis* *hispidis* Thunb. Fl. 1791. p. 297

Habitat in Japonia montibus

Caulis *terres*, *stratus*, *hispidus*, *erectus* *paniculatus*, *byudalis*. Folia *infera* *subpetiolata* *attenuata* *digitalia*; *superiora* *sessilia* *pollicaria*: *omnia* *concolorata*, *dentata* *utroque* *verso*. *Panicula* *floris* *paniculata*, *fastigiata*. *Pedunculi* *longi*, *radicelli* *subscissis*, *rufi*, *aphylli* *cernui*. *Peranth.* *duplex* *erectum*. *Semina* *oblonga* *rugosa*

1819

vellosum E. foliis oppositis delusatis ~~et~~ ovatis acutis subserratis subtus villosis
tomentosis, calycibus 2, floribus, caule fruticoso Sw. prod III
Conyza fruticosa L. in D. D. in Swan hist. 1. 257. t. 161. fig. 2.
Habitat in Jamaica, Domingo &

Sprengerianum. H. caule ramoso foliis, foliis semper amplexicaulibus oblongis ve
rardis hispida. Her. I. V. p. 7104

Hieracium puberulum ramosum Sprengerianum. Bauh. hist. 2. p. 1025.
Mon. f. hist. 3. p. 65. f. 7. t. 5. fig. 17

Habitat in Lusitania

obj. vide in Hierac. Sprengerianum

Est ~~Oxyd. specios.~~ ^{ut Hier.} Descript. in Reuch. Diversa planta a Bauh.
Descript. hinc aliam plantam ac Bauh. hist. indicare videtur