

Peucedanum

1. americana *St.* caulis nudi, foliis radicalibus septem lobatis
2. Dichotoma *St.* caule ramoso pedunculatis axillaribus bifloris foliis
lineari lanceolatis integris

villosa Heuchera

Habitat in Carolina Herb. Pursh.

Folia petiolata, cordata-subrotunda, bipolliciana, squelota: lobi ovati acuti, obtuse inaequaliter dentata, utrinque pilosa. Petioli hirsuti. Scopus semi-pedalis, inferne hirsutus, squamis nonnullis remotissimis adpressis. Panicula terminalis. Flores 3-4 ple. minores quam in Americana. Calycis villosi

americana H. scapis subnudis, thyso elongato, foliis radicalibus longe petiolatis septem
lobis, bis acute crenatis Murr. in Com. Gott 1772. pag. 66

Cottusa americana, flore squalide purpureo Herm. parad. 131.

t. 131.

Sanicula f. *cottusa americana* fructu, flori bus squalide purpureis
Pursh. alm. 332. t. 50. f. 3.

~~*H. cuthera*~~ americana Lin. Sp. pl. pag. 320

Habitata in Virginia &

Folia radicalia subrotunda cordata et angulata septem
nervi venosa margine piloso, inaequaliter dentata
dentibus acuminatis, bipollicaria et ultra. Scapus
pedalis erectus simplex superne in ramos floriferos
divisus. Pedunculi alterni pubescentes uti scapus
bractea subulata ad basin per ramos et ramulorum

— scapo paniculato foliis subseptemlobis acute crenatis