

Holotheum

herpetum *Holosteum folio orbiculato herpetis* ~~am. aind 2 p.~~

L.V. p. 129.

~~Algae folio orbiculato herpetis~~

Habitat in malabaria

~~Petala alba subtrifida, calycis mucosa L.~~

delectatur, nec enim aliud est quam genus *Crotodi* qd. hab. Rozen: und
descriptionem fersit L. Rozenii sequentem synonyma.

add. S. Spegula affinis *Pearnell* Donii a folio subrotundis pubescentibus
act. Philof. angl.

cordatum 4 foliis subcordatis

Alpne americana, nummulariifolium Herm. parad. 2. Tab. 11.

Polypodium foliis orbiculatis oppositis, axillis laxis terminalibus remotis

Brown. jam. 139.

Habitat in Jamaica, Surinamo

Caules filiformes glabri prostrumbeati. Subramosi teretibus
foliis petiolata opposita semiconvoluta mucronata
nervosa venosa glabra. Stylis utraque geminis bipartitis albis lacini
filulatis. Puncicula axillaris palmari. Dichotoma. bractea min
to filulata ad ramificatione.

pp proprie generis sic. Richard. spec. Stipulida dictum

Diandrum H. caulebus procumbentibus rigidioribus, foliis subtendens
floribus diandris Sw. prodr. p. 27.
Habitat in Jamaica C

non vidi

Fucculentum Holostem foliis ellipticis carnosis ~~Am. Acad. 3. p. 21~~ S. V. p. 129.
alpine foliis ellipticis fucculentis Cold. Nov. 9.
Habitat 7 in Novoborac'o
Petala alba subtrifida calyce minore Lin.

umbellatum # floribus umbellatis Cof. d. 1204

Caryophyllus arvensis umbellatus, foliis glabro Bauh. pin. 210.

Caryophyllus arvensis Bauh. hist. 3. p. 361.

Holosteum caryophyllum arvensis Tabern. ic. 233.

Lycnius graminis hispida umbellifera Monf. hist. 2. p. 446. p. 5. v. 22. 1746

Habitat in Europa australis arvis C.
fata Perast. Caules plures erecti tenues digitales vel palmari-
inferne foliis simplicissimis. folia sessile opposita lanceolata
acuta integerrima avenae glaucis vixia basi petiolo ornata semina
quicquid Pedunculi & geminales 4-5 umbellati sive pollicares
inaequales uniflori biflori reflexi.