

Homalium

1. racemosum H. foliis serratis, racemis axillaribus terminalibusque
floribus pedicellatis
2. Racoubea H. foliis dentatis cuneatis, racemis axillaribus ter-
minalibusque, floribus seplatis

Huc pertinet Perida River Guss

rademofum Homaliūm foliis serratis, racemis axillaribus & terminalibusque,
floribus pedunculatis h. prod. pag. 86.

Homaliūm rademofum Galqu. Amer. hist. 170. t. 103. fig. 72.

Halibati in Jamaica. 5

Rami teretes glabri, ramosi.

Folia petiolata, alterna, ovato-lanceolata, utroque glabra, acumi-
nata, venosa, palmata.

Racemi terminales et axillares, foliis brevioris, simplices, 6-7 flori.

Pedicelli alterni, unus vel alter interdum oppositus, vix semicongi-
gularis.

Calyx et Corolla semisuperi, Calycis foliola cum petala alternan-
ter breviora, ovata, oculo armato subvellosa, canescentia.

Paragermen inferum turbinatum, superum conica villosa.

Styli tres

3 Peti-floris G. ~~Caulis fruticuloso ^{toment pullos incano} pullos ^{tomento sericans} erectior~~
~~foliis obovatis plicatis.~~
~~Glomeris peti-floris~~

~~Habitat in Arabia &~~

~~Structura reliquorum ut supra limite invenuntur~~
~~Tota planta a pulis cineris incano, pulis longioris quam~~
~~in glomeride. caulis ^{hodie hodie} ramosissimus, rami diffusis alternis~~
~~geniculis ^{in raris} articulis. spinae quatuor rigidae minutae. & basi pedunculorum~~
~~persistentes. Folia ad genicula 2-4. basi~~
~~acuta subretusa plicata undulata, unguiculata~~
~~seorsim aggregata sessiles vel brevipedunculati. Calycis~~
~~brevioris foliis magis ovatis. Capsula ut in glomeride~~

pariflorum Homalium

Acoma La Marti

Habitat in Madagascariæ Retanensis herb. Mus. G.

Rami teretes, glabri, apicè & orulo armato leviter villosi. Folia petiolata, alterna bis.
In pollicana, lanceolato-oblonga, basi acuta, apicè acuminata, obtusa, remote crenata,
interdum integerrima, basi apicèque versus integerrima, membranacea,
glaberrima, supra tenuia, suba venia, subtus pallidiora, venosa. Racemi ex
apicè ramorum, composti, bis. In pollicana. Pediculi conestantcs. Bra-
ctea parva, lanceolata. Flores parvi

Homalium?

Habitat Cayenne herb. Thaurini

Rami glabri, teretes. Folia petiolata, alterna, 6 pollicaria, pennata cum impari
si que supra. foliola pinnata, et infra repanda emota. Petioli angulati.
Racemi axillares et terminales, compositi, canescentes

racemosum Homaleum, foliis serratis racemis axillaribus Verminalibus,
floribus pedicellatis

Homaleum racemosum Jacq. Am. 170. Tab. 183. fig. 72.

Habitat in America meridionali 5

Rami virides glabri, pro uacro punctati. Folia petiolato alterna
~~ovata lance ovata~~ lanceolata ovata membranacea, venosa utrin-
que glabra 4-5 pollicaria attenuata, serrata. Petiolus brevis
Racemi ^{pedunculatus filiformis} ~~ax~~ foliis paucis brevioribus ^{racematis} remotis ^{racematis} ~~palpiter~~
conspicuis
aucto anonymo Plum (in herb. Guis)

Raioubea

Dentata Coraceu

2 racemifera fl. foliis serratis, racemis caulis terminalibusque, floribus
pedicellatis
pedunculatis in prodr. pag 86.

~~Homalium racemifera L'Her. bot 170 & 103 f. 72~~

~~Habitat in Jamaica~~

Raioubea guianensis Aub. ff. in pag 591. t. 236

Habitat in Guiana

Frutex quadrupedalis. Rami vel opedale. folia alterna,
retiolata, glabra, rigida. Corolla lutea, flor. in spica
remota Aub.